

# World Journal of *Gastrointestinal Oncology*

*World J Gastrointest Oncol* 2024 April 15; 16(4): 1091-1675



## EDITORIAL

- 1091 Parallel pathways: A chronicle of evolution in rectal and breast cancer surgery  
*Pesce A, Fabbri N, Iovino D, Feo CV*
- 1097 Hepatitis B virus genotypes in precision medicine of hepatitis B-related hepatocellular carcinoma: Where we are now  
*Sukowati CHC, Jayanti S, Turyadi T, Muljono DH, Tiribelli C*

## REVIEW

- 1104 Novel milestones for early esophageal carcinoma: From bench to bed  
*Qi JH, Huang SL, Jin SZ*
- 1119 Colorectal cancer screening: A review of current knowledge and progress in research  
*Lopes SR, Martins C, Santos IC, Teixeira M, Gamito É, Alves AL*
- 1134 New avenues for the treatment of immunotherapy-resistant pancreatic cancer  
*Silva LGO, Lemos FFB, Luz MS, Rocha Pinheiro SL, Calmon MDS, Correa Santos GL, Rocha GR, de Melo FF*

## MINIREVIEWS

- 1154 Present situation of minimally invasive surgical treatment for early gastric cancer  
*Li CY, Wang YF, Luo LK, Yang XJ*
- 1166 Mixed neuroendocrine non-neuroendocrine neoplasms in gastroenteropancreatic tract  
*Díaz-López S, Jiménez-Castro J, Robles-Barraza CE, Ayala-de Miguel C, Chaves-Conde M*
- 1180 Esophageal cancer screening, early detection and treatment: Current insights and future directions  
*Qu HT, Li Q, Hao L, Ni YJ, Luan WY, Yang Z, Chen XD, Zhang TT, Miao YD, Zhang F*

## ORIGINAL ARTICLE

## Retrospective Cohort Study

- 1192 Pre-operative enhanced magnetic resonance imaging combined with clinical features predict early recurrence of hepatocellular carcinoma after radical resection  
*Chen JP, Yang RH, Zhang TH, Liao LA, Guan YT, Dai HY*
- 1204 Clinical analysis of multiple primary gastrointestinal malignant tumors: A 10-year case review of a single-center  
*Zhu CL, Peng LZ*

**Retrospective Study**

- 1213** Predictive model for non-malignant portal vein thrombosis associated with cirrhosis based on inflammatory biomarkers  
*Nie GL, Yan J, Li Y, Zhang HL, Xie DN, Zhu XW, Li X*
- 1227** Predictive modeling for postoperative delirium in elderly patients with abdominal malignancies using synthetic minority oversampling technique  
*Hu WJ, Bai G, Wang Y, Hong DM, Jiang JH, Li JX, Hua Y, Wang XY, Chen Y*
- 1236** Efficacy and predictive factors of transarterial chemoembolization combined with lenvatinib plus programmed cell death protein-1 inhibition for unresectable hepatocellular carcinoma  
*Ma KP, Fu JX, Duan F, Wang MQ*
- 1248** Should we perform sigmoidoscopy for colorectal cancer screening in people under 45 years?  
*Leong W, Guo JQ, Ning C, Luo FF, Jiao R, Yang DY*
- 1256** Computed tomography-based radiomics diagnostic approach for differential diagnosis between early- and late-stage pancreatic ductal adenocarcinoma  
*Ren S, Qian LC, Cao YY, Daniels MJ, Song LN, Tian Y, Wang ZQ*
- 1268** Prognostic analysis of related factors of adverse reactions to immunotherapy in advanced gastric cancer and establishment of a nomogram model  
*He XX, Du B, Wu T, Shen H*

**Clinical Trials Study**

- 1281** Safety and efficacy of a programmed cell death 1 inhibitor combined with oxaliplatin plus S-1 in patients with Borrmann large type III and IV gastric cancers  
*Bao ZH, Hu C, Zhang YQ, Yu PC, Wang Y, Xu ZY, Fu HY, Cheng XD*

**Observational Study**

- 1296** Computed tomography radiogenomics: A potential tool for prediction of molecular subtypes in gastric stromal tumor  
*Yin XN, Wang ZH, Zou L, Yang CW, Shen CY, Liu BK, Yin Y, Liu XJ, Zhang B*
- 1309** Application of texture signatures based on multiparameter-magnetic resonance imaging for predicting microvascular invasion in hepatocellular carcinoma: Retrospective study  
*Nong HY, Cen YY, Qin M, Qin WQ, Xie YX, Li L, Liu MR, Ding K*
- 1319** Causal roles of gut microbiota in cholangiocarcinoma etiology suggested by genetic study  
*Chen ZT, Ding CC, Chen KL, Gu YJ, Lu CC, Li QY*
- 1334** Is recovery enhancement after gastric cancer surgery really a safe approach for elderly patients?  
*Li ZW, Luo XJ, Liu F, Liu XR, Shu XP, Tong Y, Lv Q, Liu XY, Zhang W, Peng D*
- 1344** Establishment of a cholangiocarcinoma risk evaluation model based on mucin expression levels  
*Yang CY, Guo LM, Li Y, Wang GX, Tang XW, Zhang QL, Zhang LF, Luo JY*

- 1361** Effectiveness of fecal DNA syndecan-2 methylation testing for detection of colorectal cancer in a high-risk Chinese population

*Luo WF, Jiao YT, Lin XL, Zhao Y, Wang SB, Shen J, Deng J, Ye YF, Han ZP, Xie FM, He JH, Wan Y*

#### Clinical and Translational Research

- 1374** Clinical and socioeconomic determinants of survival in biliary tract adenocarcinomas

*Sahyoun L, Chen K, Tsay C, Chen G, Protiva P*

- 1384** Risk factors, prognostic factors, and nomograms for distant metastasis in patients with diagnosed duodenal cancer: A population-based study

*Shang JR, Xu CY, Zhai XX, Xu Z, Qian J*

- 1421** NOX4 promotes tumor progression through the MAPK-MEK1/2-ERK1/2 axis in colorectal cancer

*Xu YJ, Huo YC, Zhao QT, Liu JY, Tian YJ, Yang LL, Zhang Y*

#### Basic Study

- 1437** Curcumin inhibits the growth and invasion of gastric cancer by regulating long noncoding RNA AC022424.2

*Wang BS, Zhang CL, Cui X, Li Q, Yang L, He ZY, Yang Z, Zeng MM, Cao N*

- 1453** MicroRNA-298 determines the radio-resistance of colorectal cancer cells by directly targeting human dual-specificity tyrosine(Y)-regulated kinase 1A

*Shen MZ, Zhang Y, Wu F, Shen MZ, Liang JL, Zhang XL, Liu XJ, Li XS, Wang RS*

- 1465** Human  $\beta$ -defensin-1 affects the mammalian target of rapamycin pathway and autophagy in colon cancer cells through long non-coding RNA TCONS\_00014506

*Zhao YX, Cui Y, Li XH, Yang WH, An SX, Cui JX, Zhang MY, Lu JK, Zhang X, Wang XM, Bao LL, Zhao PW*

- 1479** FAM53B promotes pancreatic ductal adenocarcinoma metastasis by regulating macrophage M2 polarization

*Pei XZ, Cai M, Jiang DW, Chen SH, Wang QQ, Lu HM, Lu YF*

- 1500** Transcriptome sequencing reveals novel biomarkers and immune cell infiltration in esophageal tumorigenesis

*Sun JR, Chen DM, Huang R, Wang RT, Jia LQ*

- 1514** Construction of CDKN2A-related competitive endogenous RNA network and identification of GAS5 as a prognostic indicator for hepatocellular carcinoma

*Pan Y, Zhang YR, Wang LY, Wu LN, Ma YQ, Fang Z, Li SB*

- 1532** Two missense STK11 gene variations impaired LKB1/adenosine monophosphate-activated protein kinase signaling in Peutz-Jeghers syndrome

*Liu J, Zeng SC, Wang A, Cheng HY, Zhang QJ, Lu GX*

- 1547** Long noncoding RNAs HAND2-AS1 ultrasound microbubbles suppress hepatocellular carcinoma progression by regulating the miR-873-5p/tissue inhibitor of matrix metalloproteinase-2 axis

*Zou Q, Wang HW, Di XL, Li Y, Gao H*

- 1564** Upregulated lncRNA PRNT promotes progression and oxaliplatin resistance of colorectal cancer cells by regulating HIPK2 transcription

*Li SN, Yang S, Wang HQ, Hui TL, Cheng M, Zhang X, Li BK, Wang GY*

### **SYSTEMATIC REVIEWS**

- 1578** Prognosis value of heat-shock proteins in esophageal and esophagogastric cancer: A systematic review and meta-analysis

*Nakamura ET, Park A, Pereira MA, Kikawa D, Tustumi F*

- 1596** Risk factors for hepatocellular carcinoma associated with hepatitis C genotype 3 infection: A systematic review

*Farooq HZ, James M, Abbott J, Oyibo P, Divall P, Choudhry N, Foster GR*

### **META-ANALYSIS**

- 1613** Effectiveness and tolerability of programmed cell death protein-1 inhibitor + chemotherapy compared to chemotherapy for upper gastrointestinal tract cancers

*Zhang XM, Yang T, Xu YY, Li BZ, Shen W, Hu WQ, Yan CW, Zong L*

- 1626** Success rate of current human-derived gastric cancer organoids establishment and influencing factors: A systematic review and meta-analysis

*Jiang KL, Wang XX, Liu XJ, Guo LK, Chen YQ, Jia QL, Yang KM, Ling JH*

### **CASE REPORT**

- 1647** Pathologically successful conversion hepatectomy for advanced giant hepatocellular carcinoma after multidisciplinary therapy: A case report and review of literature

*Chu JH, Huang LY, Wang YR, Li J, Han SL, Xi H, Gao WX, Cui YY, Qian MP*

- 1660** Clinical pathological characteristics of “crawling-type” gastric adenocarcinoma cancer: A case report

*Xu YW, Song Y, Tian J, Zhang BC, Yang YS, Wang J*

- 1668** Primary pancreatic peripheral T-cell lymphoma: A case report

*Bai YL, Wang LJ, Luo H, Cui YB, Xu JH, Nan HJ, Yang PY, Niu JW, Shi MY*

**ABOUT COVER**

Peer Reviewer of *World Journal of Gastrointestinal Oncology*, Lie Zheng, Director, Professor, Department of Gastroenterology, Shaanxi Provincial Hospital of Traditional Chinese Medicine, Xi'an 730000, Shaanxi Province, China. xinliwen696@126.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Gastrointestinal Oncology* (WJGO, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, *etc.*

**INDEXING/ABSTRACTING**

The WJGO is now abstracted and indexed in PubMed, PubMed Central, Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGO as 3.0; IF without journal self cites: 2.9; 5-year IF: 3.0; Journal Citation Indicator: 0.49; Ranking: 157 among 241 journals in oncology; Quartile category: Q3; Ranking: 58 among 93 journals in gastroenterology and hepatology; and Quartile category: Q3. The WJGO's CiteScore for 2022 is 4.1 and Scopus CiteScore rank 2022: Gastroenterology is 71/149; Oncology is 197/366.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Xiang-Di Zhang; Production Department Director: Xiang Li; Cover Editor: Jia-Ru Fan.

**NAME OF JOURNAL**

*World Journal of Gastrointestinal Oncology*

**ISSN**

ISSN 1948-5204 (online)

**LAUNCH DATE**

February 15, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Monjur Ahmed, Florin Burada

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

**PUBLICATION DATE**

April 15, 2024

**COPYRIGHT**

© 2024 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Risk factors for hepatocellular carcinoma associated with hepatitis C genotype 3 infection: A systematic review

Hamzah Z Farooq, Michael James, Jane Abbott, Patrick Oyibo, Pip Divall, Naheed Choudhry, Graham R Foster

**Specialty type:** Infectious diseases

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

**P-Reviewer:** Du H, China

**Received:** November 17, 2023

**Peer-review started:** November 17, 2023

**First decision:** December 11, 2023

**Revised:** December 21, 2023

**Accepted:** January 23, 2024

**Article in press:** January 23, 2024

**Published online:** April 15, 2024



**Hamzah Z Farooq, Michael James, Jane Abbott, Naheed Choudhry, Graham R Foster,** Blizzard Institute, Queen Mary University of London, London E1 2AT, United Kingdom

**Patrick Oyibo,** School of Health and Psychological Sciences, University of London, London EC1V 0HB, United Kingdom

**Pip Divall,** University Hospitals of Leicester Library, University Hospitals of Leicester NHS Trust, Leicester LE3 9QP, United Kingdom

**Corresponding author:** Hamzah Z Farooq, MBChB, MRCP, MSc, Doctor, Research Fellow, Blizzard Institute, Queen Mary University of London, 4 Newark Street, London E1 2AT, United Kingdom. [hamzah.farooq@qmul.ac.uk](mailto:hamzah.farooq@qmul.ac.uk)

### Abstract

#### BACKGROUND

Hepatitis C virus (HCV) is a blood-borne virus which globally affects around 79 million people and is associated with high morbidity and mortality. Chronic infection leads to cirrhosis in a large proportion of patients and often causes hepatocellular carcinoma (HCC) in people with cirrhosis. Of the 6 HCV genotypes (G1-G6), genotype-3 accounts for 17.9% of infections. HCV genotype-3 responds least well to directly-acting antivirals and patients with genotype-3 infection are at increased risk of HCC even if they do not have cirrhosis.

#### AIM

To systematically review and critically appraise all risk factors for HCC secondary to HCV-G3 in all settings. Consequently, we studied possible risk factors for HCC due to HCV-G3 in the literature from 1946 to 2023.

#### METHODS

This systematic review aimed to synthesise existing and published studies of risk factors for HCC secondary to HCV genotype-3 and evaluate their strengths and limitations. We searched Web of Science, Medline, EMBASE, and CENTRAL for publications reporting risk factors for HCC due to HCV genotype-3 in all settings, 1946-2023.

#### RESULTS

Four thousand one hundred and forty-four records were identified from the four databases with 260 records removed as duplicates. Three thousand eight hundred



and eighty-four records were screened with 3514 excluded. Three hundred and seventy-one full-texts were assessed for eligibility with seven studies included for analysis. Of the seven studies, three studies were retrospective case-control trials, two retrospective cohort studies, one a prospective cohort study and one a cross-sectional study design. All were based in hospital settings with four in Pakistan, two in South Korea and one in the United States. The total number of participants were 9621 of which 167 developed HCC (1.7%). All seven studies found cirrhosis to be a risk factor for HCC secondary to HCV genotype-3 followed by higher age (five-studies), with two studies each showing male sex, high alpha fetoprotein, directly-acting antivirals treatment and achievement of sustained virologic response as risk factors for developing HCC.

## CONCLUSION

Although, studies have shown that HCV genotype-3 infection is an independent risk factor for end-stage liver disease, HCC, and liver-related death, there is a lack of evidence for specific risk factors for HCC secondary to HCV genotype-3. Only cirrhosis and age have demonstrated an association; however, the number of studies is very small, and more research is required to investigate risk factors for HCC secondary to HCV genotype-3.

**Key Words:** Hepatocellular carcinoma; Hepatitis C; Genotype 3; Systematic review; Blood-borne viruses; Liver cancer

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Hepatitis C virus (HCV) genotype-3 accounts for 17.9% of HCV infections with an increased risk of hepatocellular carcinoma (HCC) globally. In this systematic review and meta-analysis, we screened 4144 records to find only seven studies which study risk factors for HCC. Conducted primarily in Global South hospital settings, the studies encompassed 9621 participants, revealing cirrhosis and age as consistent risk factors for HCC. While cirrhosis and age emerge as contributors, the scarcity of studies underscores the urgent need for expanded research. Limited evidence exists on other factors, emphasising the need for further research to understand specific risk contributors to HCC secondary to HCV Genotype-3.

**Citation:** Farooq HZ, James M, Abbott J, Oyibo P, Divall P, Choudhry N, Foster GR. Risk factors for hepatocellular carcinoma associated with hepatitis C genotype 3 infection: A systematic review. *World J Gastrointest Oncol* 2024; 16(4): 1596-1612

**URL:** <https://www.wjgnet.com/1948-5204/full/v16/i4/1596.htm>

**DOI:** <https://dx.doi.org/10.4251/wjgo.v16.i4.1596>

## INTRODUCTION

Hepatitis C virus (HCV) is a blood-borne virus which globally affects around 79 million people[1] and is associated with high morbidity and mortality. Chronic infection leads to cirrhosis in a large proportion of patients after 30 years of asymptomatic infection and often causes hepatocellular carcinoma (HCC) in people with cirrhosis. HCV has six genotypes (G1-G6) globally with G1 accounting for 49.1% of all HCV infections, followed by G3 (17.9%), G4 (16.8%), G2 (11.0%), G5 (2.0%), and G6 (1.4%)[2].

The highest prevalence of G3 in Western Europe is Norway (50%), England (47%), and Finland (43%) with 10% in North America (22% in Canada) with 26.9% in South America[3-5]. However, the greatest burden of G3 is in South and Central Asia with 71.6% of HCV infections being of this genotype which is very common in Pakistan and India[2,6-8].

G3 infection is not susceptible to the first generation of direct-acting antiviral (DAA) protease inhibitors and has reduced susceptibility to Sofosbuvir[6,9-13], particularly in patients with cirrhosis. The efficacy of next generation protease inhibitor-based regimens (glecaprevir/pibrentasvir) may also be reduced[14-16]. However even in patients with this genotype viral clearance rates are well over 90% and these effective, affordable oral antiviral treatments are widely available. However, in patients with HCV induced cirrhosis, viral clearance does not abolish the risk of HCC[17].

HCC is a feared complication of HCV and of all the genotypes, patients with G3 infection have the highest incidence [18,19]. In most patients, cancer is linked to cirrhosis but in subjects infected with G3 even those without cirrhosis are at increased risk[20,21]. The only effective strategy to manage liver cancer is early detection of asymptomatic tumours by screening followed by loco-regional or immunomodulatory/kinase inhibitor combination therapies. Current recommendations are to screen all cirrhotic patients by ultrasound 6-monthly. However, in G3, where cirrhosis is not an adequate risk factor, we need to screen more subjects[22] and require epidemiological risk assessment tools to determine which subjects require surveillance.

There is therefore a need to identify and evaluate risk factors for HCC secondary to HCV-G3 to assist in identification of people at high risk. However, although there are risk factors identified for the most common genotype, G1; this is not the case for G3.

To address this, we aimed to systematically review and critically appraise all risk factors for HCC secondary to HCV-G3 in all settings. Consequently, we studied possible risk factors for HCC due to HCV-G3 in the literature from 1946 to 2023.



## MATERIALS AND METHODS

### Search strategy, selection criteria, screening process

**Literature search:** We searched the following four databases for articles: Web of Science, Medline, EMBASE, and CENTRAL; utilising the search strategy pre-defined by an expert librarian ([Supplementary Table 1](#)) for studies published between 1<sup>st</sup> January 1946 to 17<sup>th</sup> December 2022.

The search aimed to include all relevant studies reporting original data for the comparison of HCC risk of patients with HCV G3, from inception up until December 2022. The following keywords: “Hepatocellular carcinoma”, “hepatitis C”, and “genotype 3” were combined with other search terms, using Boolean operators and truncation. Secondly, the reference lists of all included articles were manually reviewed to identify any unidentified publications and grey literature was searched. No restrictions were set for publication year and status, or geographical area.

**Selection criteria:** We applied the following inclusion criteria to studies: (1) Participants/population: Patients in primary care, hospital settings and national databases; (2) exposure: Risk factor for HCC secondary to HCV-G3; (3) comparison: Risk factor for HCC secondary to non-HCV-G3 or control; and (4) outcome: Development of HCC.

We included randomized control trials and observational studies (case-control, cohort, and cross-sectional) and excluded any studies which did not fit the above criteria, mathematical modelling studies or were not published in English.

Studies were eligible for the meta-analysis if they fulfilled the following criteria: (1) Study design: Cohort studies, case-control studies or randomized controlled trials based on original data; (2) study population and exposure: For cohort studies, both HCV G3 infected group and a comparison group of HCV non-G3 infected patients in the same study, with at least 10 patients in each group, and for case-control studies at least 10 patients in each group of HCV-G3 HCC cases and non-HCC as controls. In studies where there were all HCV genotypes, we included those which had data (in the main results or supplementary appendices) of individual patients with HCV-G3 who developed HCC; (3) methods: Studies reporting odds ratios (OR), relative risks (RR), or hazard ratios (HR), or sufficient data to calculate the effect size (ES); (4) outcome: The number of HCC in each patient group is stated; and (5) the manuscript is published as a full paper in a peer-reviewed journal.

The following studies were excluded: (1) Animal or *in vitro* studies; (2) studies without clearly reported control or comparison group; (3) studies with unclear HCC outcome; and (4) letters to the editor, review articles, guidelines, and conference abstracts (not peer-reviewed) were excluded.

To ensure the exposure (HCV G3 infection) was present prior to the development of HCC, we excluded all studies where there was combined data in HCV genotypes and where we could not extract data (in the main results or supplementary appendices) of individual patients with HCV-G3 who developed HCC.

**Screening process:** We planned for two reviewers (HZF and MJ) to screen all abstracts to ensure a robust screening process with each abstract reviewed by at least two reviewers utilising the Rayyan QCRI programme. Any conflicting decisions were discussed and referred to a third reviewer if required.

Post primary screening, two reviewers (HZF and MJ) screened the full texts to ensure the papers fully fit the criteria with conflicting decisions discussed and referred to a third reviewer if needed.

**Data extraction (selection and coding):** Two reviewers (HZF and MJ) independently screened the full text of the included papers and extracted the following data for each included study: (1) Setting of study: Country and whether primary or secondary care; (2) characteristics of study population: Age and sex; (3) study design; (4) number of study participants in study; (5) type of HCV; (6) number participants who developed HCC; (7) risk factors identified for HCC; (8) proportion of participants with particular risk factor: Number and percentage; (9) odds ratio of risk factor; (10) hazards ratio of risk factor; and (11) number of participants who cleared HCV or were actively infected ([Supplementary Table 2](#)).

Study characteristics, context, quality, and findings were captured and summarized with similarities and differences compared across the studies in a tabular form, using appropriate subgroup analysis with comparison of the performance of different risk factors. All data were captured with a spreadsheet (MS Excel) and validated by an independent reviewer (MJ).

**Risk of bias (quality) assessment:** Two reviewers (HZF and MJ) utilised a standardised data extraction form based on the criteria for assessing the quality of risk factor studies. We utilised the Newcastle-Ottawa Scale (NOS) to assess the quality of the studies, judging studies based on points awarded for selection of study groups, comparability of groups and exposure/outcome ascertainment ([Supplementary material](#)). Any conflicting decisions were discussed and referred to a third reviewer if required. Studies with scores of < 5, 5-7, and > 7 points were considered to be of low, sufficient, and high quality, respectively. Any conflicting decisions were discussed and referred to a third reviewer (JA) if required.

### Data synthesis and statistical analysis

**Statistical analysis:** We manually extracted the crude number of patients who developed HCC in patients with HCV-G3 and utilised these data for pooled ES and 95% confidence intervals (CIs) were estimated. As the outcome of HCV G3 HCC is rare with the worldwide HCC incidence of 9.5 cases per 100000 person-years[23] odds ratios (ORs), relative risk (RR), and hazard ratios (HRs) were deemed to be equivalent. For studies which calculated HRs, we captured these for analysis. Correspondingly, and for those that had no calculated HRs we extracted the crude number of patients who developed HCC and calculated HRs.

**Meta-analysis and assessment of heterogeneity:** We carried out meta-analysis of hazard ratios in Jamovi version 2.2.5 using the “meta-analysis” package minimally adjusting for age and sex reported in the studies.

We calculated pooled summary effect estimates using the restricted-maximum likelihood model (random effects model) weighting of HRs on the natural logarithmic scale and quantified between-study heterogeneity using the  $I^2$  statistic; significance of heterogeneity was investigated using Cochran’s  $Q$  test ( $P$  threshold = 0.05). Where  $I^2$  was > 0 and heterogeneity were significant, we present random-effects summary estimates. We undertook multiple sensitivity analyses whereby analyses were restricted to studies adjusting for various additional confounders, and stratified by percentage of G3, to investigate robustness of observed associations.

**Publication bias:** Funnel plots were utilised to assess for publication bias with Egger’s regression for small-study effects used to assess the degree of asymmetry, with statistical significance level of  $P < 0.05$ .

**Funding:** The funders had no role in the study design, data collection, data analysis, data interpretation, or writing of this report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit it for publication.

## RESULTS

### Literature search

We identified 4144 records from the four databases with 260 records removed as duplicates *via* manual reviewing, 3884 records were screened with 3514 excluded (Figure 1).

Three hundred and seventy-one full texts were retrieved and assessed for eligibility of which 2 reports could not be retrieved even with contacting the study authors. Of these, 15 studies were initially included with 348 excluded. Post-preliminary analysis, seven studies were included with 363 excluded due to the defined reasons (Figure 1; Supplementary Figure 1).

### Study selection

Data were initially extracted from 15 selected studies[18,20,21,22-35] which provided information from a total of 12674 participants (Supplementary Table 3). Preliminary analysis of the studies showed that these studies combined genotype data even with their primary focus on G3. To ensure robustness of the data with particular reference to G3, the decision was made to exclude those which did not fully categorise G3 (*i.e.*, inability to extract individualised G3 participant data) and thus only seven studies were included in the final data analysis[18,26,28,29,31,33,34] (Table 1).

Of these seven studies, two studies reported on only G3, two studies had participants which were > 90% G3 and two studies had > 5% G3 participants for which de-aggregated individual data could be collected[18,26,28,29,31,33,34] (Table 2).

### Study characteristics

Of the seven studies, three studies were retrospective case-control trials, two retrospective cohort studies, one a prospective cohort study and one a cross-sectional study design (Table 2). All were based in hospital settings with four in Pakistan, two in South Korea and one in the United States. All studies required HCV RNA sequencing with genotyping *via* ISO accredited standards to demonstrate HCV infection with diagnosis of HCC based on a combination of serum alpha fetoprotein (AFP) and either imaging and/or histological classification. Only one study utilised ICD-coding which was based in the Global North[18].

One study involved a national database cohort (Military Veterans) with the other six studies involving single-centre hospital centres with participants enrolled between October 1999 and December 2016 (Tables 1 and 2).

### Study population characteristics

The total number of participants included in the analysis were 113160 with 9541 HCV-G3 of which 162 developed HCC (Table 2). There were 8826 male participants with 796 female participants showing a preponderance of male (91.7%) participants. The demographics of participants are shown in Table 2. The mean duration of follow-up was 19.93 months ranging from six to 59.6 months. Four studies enrolled Pakistani participants (100.0%)[26,29,31,33], with two Korean (100.0%)[28,34] and one study recruiting primarily White and African-American participants (85.8%)[18].

The mean age of participants was 49.77 across the seven studies. The prevalence of cirrhosis was shown in 6 out of the 7 studies with an average of 51% ranging from 12% to 100%. The majority of the studies ensured the removal of the confounding effect of co-infection with HIV and/or hepatitis B virus (HBV) infection due to their exclusion criteria as part of their protocol with only one study (Kanwal *et al*[19]) including 242 out of 8337 HCV-G3 participants (2.9%) and two studies (Cha *et al*[29] and Khan *et al*[30]) including participants with HBV and HCV-G3 [4/98 (4.1%) and 5/147 (3.4%), respectively].

All seven studies demonstrated data of HCV infection status with an average of 83.05% participants having active HCV-G3 infection (ranging from 8.27% to 100.00%) with 37.71% participants clearing HCV-G3 (range 0-92%).

### Quality assessment and risk of bias

The majority of the studies were published in Q1 or Q2 quartile journals as per SJR with only one published in a Q3

**Table 1 Study level characteristics, risk estimates of hepatocellular carcinoma and adjusted covariates of included studies**

Ref.	Country	Journal	SJR ranking quartile	Study design	Enrolment period	Study setting	Average follow-up (months)	HCV diagnosis	HCC diagnosis	Risk estimates of HCC	Covariates adjusted for
Aziz <i>et al</i> [27], 2019	Pakistan	<i>Pak J Med Sc</i>	Q3	Cross-sectional	June 2016 to January 2018	Hospital	6.0	HCV Ab, HCV RNA, and genotyping	USS abdomen, serum AFP, CT abdomen	Crude numbers	Exclusions: HBV, HIV, age < 18 yr or > 70 yr, pregnancy, previous liver lesion, “extremely fragile”, low bodyweight (not defined), known mental health issues, patients who were taking phenytoin, rifampicin, carbamazepine, patients with pancytopenia
Cha <i>et al</i> [29], 2016	Korea	<i>Medicine</i>	Q3	Retrospective case-control	January 2005 to December 2014	Hospital	59.6	HCV Ab, HCV RNA, and genotyping	USS abdomen, serum AFP, CT abdomen, histological examination	HR with 95% confidence interval	Patients with < 6 months of follow-up or patients with HCC diagnosed within 6 months of enrolment in the study
Khan <i>et al</i> [30], 2009	Pakistan	<i>Journal of Medical Virology</i>	Q1	Retrospective case-control	January 2006 to September 2007	Hospital	6.0	HCV Ab, HCV RNA, and genotyping	2 of 3 criteria: Serum AFP > 400 IU/mL, CT/MRI or liver biopsy	Crude numbers	Co-infection with HBV or HDV
Kanwal <i>et al</i> [19], 2014	United States	<i>Journal of Hepatology</i>	Q1	Retrospective cohort study	October 1999 to September 2009	Hospital	12.0	HCV Ab, HCV RNA, and genotyping	HCC (ICD-9 code 155.1)	HR with 95% confidence interval	< 1 yr of follow-up
Maryam <i>et al</i> [32], 2018	Pakistan	<i>Journal of Medical Virology</i>	Q1	Retrospective case-control	ND	Hospital	ND	HCV RNA and genotyping	Liver biopsy	Crude numbers	Nil
Park <i>et al</i> [35], 2019	Korea	<i>BMC Cancer</i>	Q1, Q2	Retrospective cohort study	January 2005 to December 2016	Hospital	24.0	HCV Ab, HCV RNA, and genotyping	USS abdomen, serum AFP, CT Abdomen, histological examination	Crude numbers	People with HIV and/or HBV, < 6 months of follow-up
Tayyab <i>et al</i> [34], 2020	Pakistan	<i>BMC Gastroenterology</i>	Q2	Prospective cohort	October 2014 to March 2017	Hospital	12.0	HCV Ab, HCV RNA, and genotyping	USS abdomen, serum AFP, CT abdomen	HR with 95% confidence interval	HBV co-infection

HCV: Hepatitis C virus; Ab: Antibody; MRI: Magnetic resonance imaging; CT: Computed tomography; USS: Ultrasound sonography; AFP: Alpha fetoprotein; HBV: Hepatitis B virus; HIV: Human immunodeficiency virus; SJR: Scimago Journal & Country Rank; Q: Quartile; HR: Hazard ratio; HCC: Hepatocellular carcinoma.

journal. The quality of the studies was moderate with majority of studies NOS scores ranging from 6 to 8 (out of maximum score of 8) with only one study scoring a very low score of two (Table 3). The study with a low score of two primarily investigated a unique genomic marker for HCC and had a small sample size and thus was included in the analysis. Five of the studies ensured good methodological quality with two of relatively low quality. All but one study had a specified enrolment period with good data on follow-up of participants.

Table 2 Patients level characteristics for studies included in the meta-analysis

Ref.	Total number of participants	Percentage genotype 3, n (%)	Number of HCV GT3 participants	Age, yr, median or mean	Sex (M/F)	Number of HCC, n (%)	Patients without HCC, n (%)	Risk factor	Number with risk factor who developed HCC, n (%)	Cirrhosis (%)	Active HCV (%)	Cleared HCV, n (%)	HIV, n (%)	HBV, n (%)	Hazards ratio of risk factor	OR/HR/RR calculation (in study or calculated independently)
Aziz <i>et al</i> [27], 2019	300	300 (100.00)	300	55.08 +/- 5.62	179/121	10 (3.33)	290 (96.67)	DAA treatment (SOF + DAC +/- RBV)	10 (3.33)	100	100	276 (92.00)	0 (0)	0 (0)		Independent calculation
								Child Pugh A (compensated cirrhosis) and SVR not achieved	2 (0.67)	100	100	276 (92.00)	0 (0)	0 (0)		Independent calculation
								Child Pugh B (compensated cirrhosis) and SVR achieved	5 (1.67)	100	100	276 (92.00)	0 (0)	0 (0)		Independent calculation
								Child Pugh B (Decompensated cirrhosis) and SVR not achieved	3 (1.00)	100	100	276 (92.00)	0 (0)	0 (0)		Independent calculation
								Male	7 (2.33)	100	100	276 (92.00)	0 (0)	0 (0)		Independent calculation
								Female	3 (1.00)	100	100	276 (92.00)	0 (0)	0 (0)		Independent calculation
Cha <i>et al</i> [29], 2016	1335	98 (7.30)	98	41.8 +/- 10.5	79/19	4 (4.10)	94 (95.92)	Age > 40 yr	ND	25.50	100	34 (34.70)	0 (0)	4 (4.1)	2.697 (0.436-16.683), <i>P</i> = 0.286	Calculated in study
								Cirrhosis at enrolment	25 (25.50)	25.50	100	34 (34.70)	0 (0)	4 (4.1)	33.834 (2.088-548.269), <i>P</i> = 0.013	Calculated in study
								Alcohol intake > 40 g/d	53 (54.60)	25.50	100	34 (34.70)	0 (0)	4 (4.1)	8.556 (0.693-105.623), <i>P</i> = 0.094	Calculated in study
								SVR	34 (34.70)	25.50	100	34 (34.70)	0 (0)	4 (4.1)	0.848 (0.063-11.445), <i>P</i> = 0.901	Calculated in study

									Decompensated cirrhosis and achieved SVR*	1	25.50	100	34 (34.70)	0 (0)	4 (4.1)		Independent calculation
									Did not achieve SVR*	1	25.50	100	34 (34.70)	0 (0)	4 (4.1)		Independent calculation
									Low platelet count	ND	25.50	100	34 (34.70)	0 (0)	4 (4.1)	1.00 (1.00-1.00), <i>P</i> = 0.872	Calculated in study
Khan <i>et al</i> [30], 2009	158	147 (93.00)	147	47.3 +/- 12.5	102/56	65 (44.20)	82 (55.78)	Male	51	17.69	87.10	30 (18.99)	0 (0)	5			Independent calculation
								Female	14	17.69	87.10	30 (18.99)	0 (0)	5			Independent calculation
								Age > 46.9 yr	65	17.69	87.10	30 (18.99)	0 (0)	5			Independent calculation
								High AFP	65	17.69	87.10	30 (18.99)	0 (0)	5			Independent calculation
								High HCV VL	65	17.69	87.10	30 (18.99)	0 (0)	5			Independent calculation
								ALP > 68	65	17.69	87.10	30 (18.99)	0 (0)	5			Independent calculation
								Anti-HBc*	46	17.69	87.10	30 (18.99)	0 (0)	5			Independent calculation
								HCV viraemia*	58	17.6	87.10	30 (18.99)	0 (0)	5			Independent calculation
Kanwal <i>et al</i> [19], 2014	110484	8337 (7.54)	8337	50.2 +/- 6.4	8095/242	ND	ND	Cirrhosis	ND	12	86	1167 (14.00)	242 (2.9)	0 (0)	1.44 (1.23-1.68)		Calculated in study
								Diabetes	ND	12	86	1167 (14.00)	242 (2.9)	0 (0)	1.30 (1.88-1.90)		Calculated in study
								Age > 50 yr	ND	12	86	1167 (14.00)	242 (2.9)	0 (0)	1.79 (1.53-2.11)		Calculated in study
								Age < 50 yr	ND	12	86	1167 (14.00)	242 (2.9)	0 (0)	1.86 (1.56-2.22)		Calculated in study
Maryam <i>et al</i> [32], 2018	50	50 (100.00)	50	58 (47-73)	37/23	27 (54.00)	23 (46.00)	NRAS oncogene	27 (54.00)	ND	100	0	ND	ND			Independent calculation
								Male*	22	ND	100	0	ND	ND			Independent calculation
								Female*	5	ND	100	0	ND	ND			Independent

															calculation	
Park <i>et al</i> [35], 2019	180	16 (8.88)	16	46 (40-53)	45306	16 (100.00)	0 (0)	Male	15 (93.80)	100	100	2 (12.50)	0 (0)	0 (0)	Independent calculation	
								Diabetes	6 (40.00)	100	100	2 (12.50)	0 (0)	0 (0)	Independent calculation	
								Cirrhosis	16 (100.00)	100	100	2 (12.50)	0 (0)	0 (0)	Independent calculation	
								Alcohol intake > 60 g/d	3 (18.80)	100	100	2 (12.50)	0 (0)	0 (0)	Independent calculation	
								High HCV VL	6 (37.70)	100	100	2 (12.50)	0 (0)	0 (0)	Independent calculation	
								MELD-score > 9.5	16 (100.00)	100	100	2 (12.50)	0 (0)	0 (0)	Independent calculation	
								Female	1 (6.25)	100	100	2 (12.50)	0 (0)	0 (0)	Independent calculation	
								High AFP	16 (100.00)	100	100	2 (12.50)	0 (0)	0 (0)	Independent calculation	
								Not achieved SVR*	2	100	100	2 (12.50)	0 (0)	0 (0)	Independent calculation	
Tayyab <i>et al</i> [34], 2020	653	593 (90.81)	593	50 (41-56)	319/334	40 (6.13)	613 (93.87)	Age, per 10-yr increase	ND	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	1.71 (1.25-2.33), <i>P</i> = 0.001	Calculated in study
								Use of SOF/DCV/RBV	9 (22.50)	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	17.05 (2.09-139.47), <i>P</i> = 0.01	Calculated in study
								Cirrhosis	40 (6.13)	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation
								Male*	18	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation
								Female*	22	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation
								High BMI*	5	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation
								Hypertension*	3	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation
								Diabetes*	19	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation

HBV Co-infection*	12	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation
Achieved SVR*	35	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation
Not achieved SVR*	5	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation
SOF/RBV use*	29	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation
SOF/RBV/PEG- IFN use*	1	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation
SOF/DCV use*	1	49.31	54 (8.27)	599 (91.78)	ND	0 (0)	ND	Independent calculation

NB: \*Risk factor not deemed statistically significant in the study's own analysis. Crude numbers were extracted to pool and calculate hazard ratios. HR: Hazard ratio; OR: Odds ratio; RR: Relative risks; ND: Not determined; SOF: Sofosbuvir; RBV: Ribavirin; PEG-IFN: Pegylated interferon; DCV: Daclatasvir; SVR: Sustained virologic response; HBV: Hepatitis B virus; HCV: Hepatitis C virus; Anti-HBc: Total antibody to hepatitis B core antigen; AFP: Alpha feto protein; MELD: Model for end-stage liver disease; VL: Viral load; HCC: Hepatocellular carcinoma.

### **Risk of HCC secondary to HCV-G3**

Overall, 162 participants (1.7%) developed HCC during the follow-up period. The risk factors studied by the seven studies can be categorised as either participant background factors, biochemical factors or treatment factors ([Supplementary Figure 1](#)). The majority of the studies investigated the potential risk factors of gender at birth (male/female = five studies), cirrhosis (seven), age (five) from a participant background perspective. For treatment factors, the risk factors studied related to achievement of SVR (seven) or use of DAAs (five) with studied biochemical risk factors of high AFP (two), high HCV viral load (two) with one study each on ALP, low platelets levels, Child-Pugh Score (B or C), and high Model for End-Stage Liver Disease (MELD) score.

From their primary analysis, a total of seven studies assessed demonstrated cirrhosis to be a risk factor for HCC secondary to HCV-G3 followed by higher age (5), with two studies each showing male sex, high AFP, DAA treatment and achievement of SVR as risk factors for developing HCC.

A total of seven studies assessed demonstrated cirrhosis to be a risk factor for HCC secondary to HCV-G3 followed by higher age (5), with two studies each showing male sex, high AFP, DAA treatment and achievement of SVR as risk factors for developing HCC.

Utilising the individual participant data ([Table 4](#)) and pooling the data from the seven studies, we only found a strong association between age > 50 (HR: 1.86, 95%CI: 1.05-2.55). We also found a relatively moderate association with cirrhosis (HR: 1.44, 95%CI: 0.78-2.10), high AFP (HR: 0.97, 95%CI: 0.57-1.37), male gender (HR: 0.93, 95%CI: 0.45-1.41) and weight gain (HR: 0.84, 95%CI: 0.37-1.31), high HCV VL (HR: 0.43, 95%CI: 0.03-0.83), ALP > 68 (HR: 0.43, 95%CI: 0.03-0.83), and alcohol intake > 40 g/dL (HR: 0.24, 95%CI: 0.17-0.34) ([Figure 2](#)).

Some studies also showed an association between DAA use, MELD Score >9.5, Female, Diabetes, NRAS Oncogene ([Figure 2](#)). However, high statistical heterogeneity ( $I^2 = 79.84\%$ , with  $P < 0.001$ ) was observed. As the heterogeneity was high, the factors were not fully combined for a pooled HR: to demonstrate an appropriate view of the data. The funnel plot did demonstrate asymmetry (Egger's test = 4.936,  $P < 0.001$ ) did not indicate for small-study effects ([Figure 3](#)).



Table 3 The Newcastle-Ottawa Scale assessment for included studies

Ref.	Type of Study	Selection								Compara- bility	Exposure					Total score	
		Adequate case defini- tion <sup>1</sup>	Represe- ntative- ness of cases <sup>1</sup>	Selection of controls <sup>1</sup>	Definition of controls <sup>1</sup>	Represe- ntative of exposed cohort <sup>2</sup>	Selection of non- exposed cohort <sup>2</sup>	Ascertain- ment of expo- sure <sup>2</sup>	Demonst- ration that outcome of interest was not present at start of study <sup>2</sup>	Compara- bility based on the design or analysis	Ascertain- ment of expo- sure <sup>1</sup>	Same method of ascertain- ment for partici- pants <sup>1</sup>	Nonres- ponse rate <sup>1</sup>	Assess- ment of outcome <sup>2</sup>	Was follow-up long enough for out- comes to occur <sup>2</sup>	Adequ- acy of follow up of cohorts <sup>2</sup>	
Aziz <i>et al</i> [27], 2019	Cross- Sectional	NA	NA	NA	NA	*	*	*	*	*	NA	NA	NA	*		*	7
Cha <i>et al</i> [29], 2016	Cohort	NA	NA	NA	NA	*	*	*	*	*	NA	NA	NA	*	*	*	8
Khan <i>et al</i> [30], 2009	Case- control	*			*	NA	NA	NA	NA	*	*	*	*	NA	NA	NA	6
Kanwal <i>et al</i> [19], 2014	Cohort	NA	NA	NA	NA	*	*	*	*	*	NA	NA	NA	*	*	*	8
Maryam <i>et al</i> [32], 2018	Case- control				*	NA	NA	NA	NA			*	*	NA	NA	NA	3
Park <i>et al</i> [35], 2019	Cohort	NA	NA	NA	NA	*	*	*	*	*	NA	NA	NA	*	*	*	8
Tayyab <i>et al</i> [34], 2020	Cohort	NA	NA	NA	NA	*	*	*	*	*	NA	NA	NA	*	*	*	8

<sup>1</sup>For case-control studies.<sup>2</sup>For cohort or cross-sectional studies.

NA: Not applicable.

### Sub-group analysis

We performed a sub-analysis of risk factors where there were more than three studies studying the uniform risk factor. We pooled the HRs of to show an overall effect size, utilising the random effects model (Figure 4).

When exclusively pooling the studies, the combined HR: for cirrhosis is 0.49 (95%CI: 0.02-0.96,  $I^2 = 98.96\%$ ,  $P \leq 0.001$ ,  $n = 3$ ), for age 1.43 (95%CI: 0.73-2.13,  $I^2 = 96.44\%$ ,  $n = 4$ ) and for male gender 0.41 (95%CI: -0.11 to 0.94,  $I^2 = 99.45\%$ ,  $P \leq 0.001$ ,  $n = 3$ ).

**Table 4 Pooled individual participant data for all participants with hepatitis C genotype 3 who developed hepatocellular carcinoma**

Risk factor	Number of participants
Patient-dependent factors	
Cirrhosis	66
Male	118
Female	62
Age > 40 yr	65
Alcohol intake > 40 g/d	56
Anti-HBc	46
Diabetes	25
Age < 50 yr	0
NRAS oncogene	27
Age, per 10-yr increase	0
High BMI	5
Hypertension	3
HBV co-infection	12
Treatment dependent factors	
DAA treatment	66
SVR achieved	74
SVR not achieved	13
Decompensated cirrhosis and achieved SVR	1
Use of SOF/DCV/RBV	9
Biochemical factors	
Low platelet count	16
High AFP	65
High HCV VL	71
ALP > 68	65
HCV viraemia	58
MELD-score > 9.5	16


HBc: Hepatitis B virus core protein; BMI: Body mass index; DAA: Direct-acting antiviral; SVR: Sustained virologic response; SOF: Sofosbuvir; DCV: Daclatasvir; RBV: Ribavirin; HBV: Hepatitis B virus; HCV: Hepatitis C virus; AFP: Alpha feto protein; MELD: Model for end-stage liver disease; VL: Viral load.

## DISCUSSION

HCV infection represents a significant global health burden, with millions of individuals affected worldwide. Considered a “viral time bomb”[36], the World Health Organization’s (WHO) ambitious target of eliminating HCV as a public health threat by 2030 has spurred unprecedented efforts to increase screening, diagnosis, and treatment access. While advances in DAA therapy have led to remarkable rates of viral clearance, the emergence of HCC in post-treatment patients has raised concerns and new challenges.

Among the various HCV genotypes, G3 has attracted particular attention due to its distinctive association with HCC development. Notably, patients infected with G3 have a higher predisposition to developing HCC, even in the absence of cirrhosis. This unique genotype’s enhanced hepatocarcinogenic potential warrants further exploration and the need to investigate risk factors associated with the development to HCC.

This systematic review of seven HCV-G3 studies with 9541 HCV-G3 participants shows that cirrhosis and age greater than 40 are principal risk factors for developing HCC in people with HCV-G3. It is the largest study focusing on HCC secondary to HCV-G3 of which 162 developed HCC.




**Figure 1** Study selection process for studies from 1<sup>st</sup> January 1946 to 17<sup>th</sup> December 2022. HCC: Hepatocellular carcinoma; HBV: Hepatitis B virus; G3: Genotype 3.

This study shows that there are few published studies on HCV-G3 and HCC and the majority of the studies are observational studies of retrospective design which do not have the ability to fully investigate confounding factors. It also demonstrates that the data is very heterogenous in HCV-G3 studies with a lack of high-quality studies and randomised control trials with a focus on HCV-G3. Of note, there is a lack of data and association with diabetes, HBV co-infection and/or high BMI especially with the increasing prevalence of metabolic dysfunction-associated steatotic liver disease.


This may be due to the lack of G3 patients and participants in the countries where the majority of HCV and HCC clinical trials occur. The highest global prevalence is of G3 in South and Central Asia (71.6% of HCV infection), contrasting to 24.8% in Western Europe and 10%-12% in the United States[5], where the highest number of HCV and HCC clinical trials occur. Without adequate G3 participants in the Global North, it is difficult to power studies to demonstrate appropriate risk factors for HCC in HCV-G3.

Correspondingly, in this study we have noted that there is only a moderate association of HCV-G3 with cirrhosis leading to HCC. This contrasts to G1 where there has been established a high association of cirrhosis with HCC[37-39] with some studies demonstrating a significant HR of 6.686 (4.319-10.350)[40]. Similar significant associations with cirrhosis and HCC were noted in G4[24] and G6[41], with a lack of data for G5 due to its low global prevalence. Majority of HCC predictive scores aim to quantify HCC risk in the presence of cirrhosis due to the high association with HCC[42-45]. However, these scores have been developed and validated on a predominance of G1 and G2 participants with a low percentage of G3 participants, warranting further studies for HCC in G3-predominant populations.

Efforts to eliminate HCV, especially in regions with high endemicity of G3, such as in India and Pakistan, face substantial challenges. The efficacy of treatment strategies in curbing HCV transmission must be supported by surveillance for potential risk of subsequent HCC development in patients with G3 mono-infection and those with co-infection with HBV and/or HIV. The evolving epidemiological landscape demands careful surveillance and long-term follow-up of patients treated for HCV, particularly those belonging to high-risk populations.



**Figure 2** Meta-analysis with hazard ratios of all included studies on risk factors for hepatocellular carcinoma associated with Hepatitis C Genotype 3. SOF: Sofosbuvir; DAC: Daclatasvir; RBV: Ribavirin; AFP: Alpha fetoprotein; ALP: Alkaline phosphatase; HCV: Hepatitis C virus.




**Figure 3** Funnel plot of all included studies.

To supplement the WHO's ambitious HCV elimination goals and reduce the burden of associated disease, it is imperative to implement proactive measures for identifying and managing HCC risk in patients post viral clearance. Strategies may include intensified surveillance, targeted risk stratification, and tailored treatment approaches based on HCV genotype and individual patient characteristics.

There is currently a lack of data in the literature regarding the risk factors for HCC secondary to hepatitis HCV-G3. Yet, no confirmed risk factors have been identified. To better understand the risk factors for HCC secondary to HCV-G3, a case-control trial is needed. Such a trial would allow for a more in-depth investigation of the risk factors associated with this condition.

## CONCLUSION

The global initiative to eliminate HCV by 2030 represents a remarkable public health undertaking. However, the emergence of HCC as a significant concern in patients post viral clearance, particularly in HCV-G3 infections, demands



**Figure 4 Forest plot.** A: Forest plot of studies studying age as a risk factor for hepatocellular carcinoma associated with hepatitis C genotype 3 (HCV-G3); B: Forest plot of studies studying male sex as a risk factor for hepatocellular carcinoma associated with HCV-G3; C: Forest plot of studies studying cirrhosis as a risk factor for hepatocellular carcinoma associated with HCV-G3.

careful consideration. Collaborative efforts between healthcare providers, researchers, and policymakers are essential to develop effective risk mitigation strategies while ensuring the successful elimination of HCV on a global scale. Continued research into the mechanistic basis of HCC development in HCV-G3 infections will be crucial in shaping preventive and therapeutic interventions to safeguard the progress made towards an HCV-free future.

## ARTICLE HIGHLIGHTS

**Research background**

Neglected hepatitis C genotype 3 (HCV-G3) is a global health concern as it is more oncogenic than other genotypes.

**Research motivation**

It leads to hepatocellular carcinoma (HCC) in people without cirrhosis and HCV-G3 HCC risk factors are currently unknown with no validated risk assessment tools.

**Research objectives**

To systematically review and critically appraise all risk factors for HCC secondary to HCV-G3 in all settings. Consequently, we studied possible risk factors for HCC due to HCV-G3 in the literature from 1946 to 2023.

**Research methods**

We searched the following four databases for articles: Web of Science, Medline, EMBASE, and CENTRAL; for studies published between 1<sup>st</sup> January 1946 to 17<sup>th</sup> December 2022.

**Research results**

Cirrhosis, higher age, and male gender were found to be strongly associated with HCC due to HCV-G3.

**Research conclusions**

There is currently a lack of data in the literature regarding the risk factors for HCC secondary to HCV-G3. As of yet, no confirmed risk factors have been identified.

**Research perspectives**

With limited studies on HCV-G3 and HCC, further research is needed to provide a risk assessment tool for HCC secondary to HCV-G3.

## FOOTNOTES

**Author contributions:** Farooq HZ, Choudhry N, and Foster GR contributed to conceptualization; Farooq HZ, James M, and Abbott J contributed to data curation; Farooq HZ and Oyibo P contributed to formal analysis; Farooq HZ and Foster GR contributed to funding acquisition; Farooq HZ and James M contributed to investigation; Farooq HZ and Foster GR contributed to methodology; Foster GR contributed to supervision; Farooq HZ and Oyibo P contributed to validation; Farooq HZ and Oyibo P contributed to visualization; Farooq HZ and Foster GR contributed to roles/writing-original draft; all authors contributed to writing-review and editing.

**Supported by** the Clinical Research Fellowship Grant from the Wellcome Trust, United Kingdom, No. 227516/Z/23/Z.

**Conflict-of-interest statement:** The authors declare that they have no conflict of interest.

**PRISMA 2009 Checklist statement:** The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** United Kingdom

**ORCID number:** Hamzah Z Farooq 0000-0002-1361-9692; Graham R Foster 0000-0002-3704-386X.

**S-Editor:** Chen YL

**L-Editor:** A

**P-Editor:** Zhang XD

## REFERENCES

- 1 Mahmud S, Al Kanaani Z, Abu-Raddad LJ. Characterization of the hepatitis C virus epidemic in Pakistan. *BMC Infect Dis* 2019; **19**: 809 [PMID: 31521121 DOI: 10.1186/s12879-019-4403-7]
- 2 Petruzzello A, Marigliano S, Loquercio G, Cozzolino A, Cacciapuoti C. Global epidemiology of hepatitis C virus infection: An up-date of the distribution and circulation of hepatitis C virus genotypes. *World J Gastroenterol* 2016; **22**: 7824-7840 [PMID: 27678366 DOI: 10.3748/wjg.v22.i24.7824]



- 10.3748/wjg.v22.i34.7824]
- 3 **Chan CP**, Uemura H, Kwan TH, Wong NS, Oka S, Chan DPC, Lee SS. Review on the molecular epidemiology of sexually acquired hepatitis C virus infection in the Asia-Pacific region. *J Int AIDS Soc* 2020; **23**: e25618 [PMID: 32969173 DOI: 10.1002/jia2.25618]
- 4 **Gower E**, Estes C, Blach S, Razavi-Shearer K, Razavi H. Global epidemiology and genotype distribution of the hepatitis C virus infection. *J Hepatol* 2014; **61**: S45-S57 [PMID: 25086286 DOI: 10.1016/j.jhep.2014.07.027]
- 5 **Messina JP**, Humphreys I, Flaxman A, Brown A, Cooke GS, Pybus OG, Barnes E. Global distribution and prevalence of hepatitis C virus genotypes. *Hepatology* 2015; **61**: 77-87 [PMID: 25069599 DOI: 10.1002/hep.27259]
- 6 **Chan A**, Patel K, Naggie S. Genotype 3 Infection: The Last Stand of Hepatitis C Virus. *Drugs* 2017; **77**: 131-144 [PMID: 28074358 DOI: 10.1007/s40265-016-0685-x]
- 7 **Goossens N**, Negro F. Is genotype 3 of the hepatitis C virus the new villain? *Hepatology* 2014; **59**: 2403-2412 [PMID: 24155107 DOI: 10.1002/hep.26905]
- 8 **Erman Daloğlu A**, Parkan ÖM, Erdoğan A, Peker BO, Can Sarınoğlu R, Sağlık İ, İnan D, Kuloğlu MM, Mutlu D, Öngüt G, Çolak D. [Distribution of Hepatitis C Virus (HCV) Genotypes Among Intravenous Drug and Non-Drug User Patients]. *Mikrobiyol Bul* 2021; **55**: 30-40 [PMID: 33590979 DOI: 10.5578/mb.20108]
- 9 **Butt AS**, Abbas Z, Jafri W. Hepatocellular carcinoma in pakistan: where do we stand? *Hepat Mon* 2012; **12**: e6023 [PMID: 23166534 DOI: 10.5812/hepatmon.6023]
- 10 **Young J**, Wong S, Janjua NZ, Klein MB. Comparing direct acting antivirals for hepatitis C using observational data - Why and how? *Pharmacol Res Perspect* 2020; **8**: e00650 [PMID: 32894643 DOI: 10.1002/prp2.650]
- 11 **Afzal MS**. Hepatitis C Virus and Interferon-Free Antiviral Therapeutics Revolution: Implications for Pakistan. *Viral Immunol* 2017; **30**: 252-257 [PMID: 28118096 DOI: 10.1089/vim.2016.0164]
- 12 **Akhtar S**, Moatter T, Azam SI, Rahbar MH, Adil S. Prevalence and risk factors for intrafamilial transmission of hepatitis C virus in Karachi, Pakistan. *J Viral Hepat* 2002; **9**: 309-314 [PMID: 12081609 DOI: 10.1046/j.1365-2893.2002.00350.x]
- 13 **Amougou-Atsama M**, Jean Adrien Atangana P, Noah Noah D, Fewou Moundipa P, Pineau P, Njouom R. The role of hepatitis C virus genotypes and core mutations in hepatocellular carcinoma in Cameroon. *J Viral Hepat* 2020; **27**: 880-885 [PMID: 32301239 DOI: 10.1111/jvh.13303]
- 14 **Chhatwal J**, Chen Q, Ayer T, Bethea ED, Kanwal F, Kowdley KV, Wang X, Roberts MS, Gordon SC. Hepatitis C virus re-treatment in the era of direct-acting antivirals: projections in the USA. *Aliment Pharmacol Ther* 2018; **47**: 1023-1031 [PMID: 29377245 DOI: 10.1111/apt.14527]
- 15 **Zeuzem S**, Foster GR, Wang S, Asatryan A, Gane E, Feld JJ, Asselah T, Bourlière M, Ruane PJ, Wedemeyer H, Pol S, Flisiak R, Poordad F, Chuang WL, Stedman CA, Flamm S, Kwo P, Dore GJ, Sepulveda-Arzola G, Roberts SK, Soto-Malave R, Kaita K, Puoti M, Vierling J, Tam E, Vargas HE, Bruck R, Fuster F, Paik SW, Felizarta F, Kort J, Fu B, Liu R, Ng TI, Pilot-Matias T, Lin CW, Trinh R, Mensa FJ. Glecaprevir-Pibrentasvir for 8 or 12 Weeks in HCV Genotype 1 or 3 Infection. *N Engl J Med* 2018; **378**: 354-369 [PMID: 29365309 DOI: 10.1056/NEJMoa1702417]
- 16 **Wei L**, Wang G, Alami NN, Xie W, Heo J, Xie Q, Zhang M, Kim YJ, Lim SG, Fredrick LM, Lu W, Liu W, Kalluri HV, Krishnan P, Tripathi R, Mobashery N, Burroughs M, Asatryan A, Jia J, Hou J. Glecaprevir-pibrentasvir to treat chronic hepatitis C virus infection in Asia: two multicentre, phase 3 studies- a randomised, double-blind study (VOYAGE-1) and an open-label, single-arm study (VOYAGE-2). *Lancet Gastroenterol Hepatol* 2020; **5**: 839-849 [PMID: 32682494 DOI: 10.1016/S2468-1253(20)30086-8]
- 17 **Yamana Y**, Kanda T, Matsumoto N, Honda M, Kumagawa M, Sasaki R, Kanezawa S, Mizutani T, Yamagami H, Masuzaki R, Ishii T, Nirei K, Moriama M. Efficacy of Glecaprevir/Pibrentasvir for Real-World HCV Infected Patients in the Northern Part of Tokyo, Japan. *J Clin Med* 2021; **10** [PMID: 34884231 DOI: 10.3390/jcm10235529]
- 18 **Dash S**, Aydin Y, Widmer KE, Nayak L. Hepatocellular Carcinoma Mechanisms Associated with Chronic HCV Infection and the Impact of Direct-Acting Antiviral Treatment. *J Hepatocell Carcinoma* 2020; **7**: 45-76 [PMID: 32346535 DOI: 10.2147/JHC.S221187]
- 19 **Kanwal F**, Kramer JR, Ilyas J, Duan Z, El-Serag HB. HCV genotype 3 is associated with an increased risk of cirrhosis and hepatocellular cancer in a national sample of U.S. Veterans with HCV. *Hepatology* 2014; **60**: 98-105 [PMID: 24615981 DOI: 10.1002/hep.27095]
- 20 **Lee SS**, Kim CY, Kim BR, Cha RR, Kim WS, Kim JJ, Lee JM, Kim HJ, Ha CY, Kim TH, Jung WT, Lee OJ. Hepatitis C virus genotype 3 was associated with the development of hepatocellular carcinoma in Korea. *J Viral Hepat* 2019; **26**: 459-465 [PMID: 30516858 DOI: 10.1111/jvh.13047]
- 21 **Nkontchou G**, Ziol M, Aout M, Lhabadie M, Baazia Y, Mahmoudi A, Roulot D, Ganne-Carrie N, Grando-Lemaire V, Trinchet JC, Gordien E, Vicaud E, Baghad I, Beaugrand M. HCV genotype 3 is associated with a higher hepatocellular carcinoma incidence in patients with ongoing viral C cirrhosis. *J Viral Hepat* 2011; **18**: e516-e522 [PMID: 21914071 DOI: 10.1111/j.1365-2893.2011.01441.x]
- 22 **McMahon BJ**, Bruden D, Townshend-Bulson L, Simons B, Spradling P, Livingston S, Gove J, Hewitt A, Plotnik J, Homan C, Espera H, Negus S, Snowball M, Barbour Y, Bruce M, Gounder P. Infection With Hepatitis C Virus Genotype 3 Is an Independent Risk Factor for End-Stage Liver Disease, Hepatocellular Carcinoma, and Liver-Related Death. *Clin Gastroenterol Hepatol* 2017; **15**: 431-437.e2 [PMID: 27765729 DOI: 10.1016/j.cgh.2016.10.012]
- 23 **Rumgay H**, Arnold M, Ferlay J, Lesi O, Cabaasag CJ, Vignat J, Laversanne M, McGlynn KA, Soerjomataram I. Global burden of primary liver cancer in 2020 and predictions to 2040. *J Hepatol* 2022; **77**: 1598-1606 [PMID: 36208844 DOI: 10.1016/j.jhep.2022.08.021]
- 24 **El-Serag HB**, Kanwal F, Richardson P, Kramer J. Risk of hepatocellular carcinoma after sustained virological response in Veterans with hepatitis C virus infection. *Hepatology* 2016; **64**: 130-137 [PMID: 26946190 DOI: 10.1002/hep.28535]
- 25 **Adnan F**, Khan NU, Iqbal A, Ali I, Petruzzello A, Sabatino R, Guzzo A, Loquercio G, Botti G, Khan S, Naeem M, Khan MI. Interleukin-6 polymorphisms in HCC patients chronically infected with HCV. *Infect Agent Cancer* 2020; **15**: 21 [PMID: 32266003 DOI: 10.1186/s13027-020-00285-9]
- 26 **Asim M**, Sarma MP, Kar P. Etiological and molecular profile of hepatocellular cancer from India. *Int J Cancer* 2013; **133**: 437-445 [PMID: 23233429 DOI: 10.1002/ijc.27993]
- 27 **Aziz B**, Nazar T, Akhlaq S. The frequency of occurrence of Hepatocellular Carcinoma after direct antiviral therapy in Hepatitis C virus patients. *Pak J Med Sci* 2019; **35**: 101-105 [PMID: 30881405 DOI: 10.12669/pjms.35.1.109]
- 28 **Ul Abideen Z**, Siddique S, Nasrullah I, Khan JS, Rehman S, Tong Y, Qamar R, Butt AM. A 2-year retrospective study of viral and host-associated risk factors in Pakistani hepatocellular carcinoma patients. *Eur J Gastroenterol Hepatol* 2019; **31**: 1103-1109 [PMID: 30829691 DOI: 10.1097/MEG.0000000000001384]
- 29 **Cha RR**, Lee SS, Lee CM, Ji SB, Jung HC, Cho HC, Kim JJ, Lee JM, Kim HJ, Ha CY, Kim TH, Jung WT, Lee OJ. Clinical Features and Outcomes of Patients With Genotype 3 Hepatitis C Virus Infection in Korea: A Retrospective Observational Study. *Medicine (Baltimore)* 2016;



- 95: e2755 [PMID: 26871824 DOI: 10.1097/MD.0000000000002755]
- 30 **Khan A**, Tanaka Y, Azam Z, Abbas Z, Kurbanov F, Saleem U, Hamid S, Jafri W, Mizokami M. Epidemic spread of hepatitis C virus genotype 3a and relation to high incidence of hepatocellular carcinoma in Pakistan. *J Med Virol* 2009; **81**: 1189-1197 [PMID: 19475617 DOI: 10.1002/jmv.21466]
- 31 **Gulnaz A**, Sayyed AH, Amin F, Khan Au, Aslam MA, Shaikh RS, Ali M. Association of XRCC1, XRCC3, and XPD genetic polymorphism with an increased risk of hepatocellular carcinoma because of the hepatitis B and C virus. *Eur J Gastroenterol Hepatol* 2013; **25**: 166-179 [PMID: 23044807 DOI: 10.1097/MEG.0b013e328359a775]
- 32 **Maryam M**, Idrees M. Study of promoter hypomethylation profiles of RAS oncogenes in hepatocellular carcinoma derived from hepatitis C virus genotype 3a in Pakistani population. *J Med Virol* 2018; **90**: 1516-1523 [PMID: 29742286 DOI: 10.1002/jmv.25221]
- 33 **Wu N**, Rao HY, Yang WB, Gao ZL, Yang RF, Fei R, Gao YH, Jin Q, Wei L. Impact of hepatitis C virus genotype 3 on liver disease progression in a Chinese national cohort. *Chin Med J (Engl)* 2020; **133**: 253-261 [PMID: 31934936 DOI: 10.1097/CM9.0000000000000629]
- 34 **Tayyab GUN**, Rasool S, Nasir B, Rubi G, Abou-Samra AB, Butt AA. Hepatocellular carcinoma occurs frequently and early after treatment in HCV genotype 3 infected persons treated with DAA regimens. *BMC Gastroenterol* 2020; **20**: 93 [PMID: 32252635 DOI: 10.1186/s12876-020-01249-4]
- 35 **Park HK**, Lee SS, Im CB, Im C, Cha RR, Kim WS, Cho HC, Lee JM, Kim HJ, Kim TH, Jung WT, Lee OJ. Hepatitis C virus genotype affects survival in patients with hepatocellular carcinoma. *BMC Cancer* 2019; **19**: 822 [PMID: 31429755 DOI: 10.1186/s12885-019-6040-3]
- 36 **Littera R**, Zamboni F, Tondolo V, Fantola G, Chessa L, Orrù N, Sanna M, Valentini D, Cappai L, Mulargia M, Caocci G, Arras M, Floris A, Orrù S, La Nasa G, Carcassi C. Absence of activating killer immunoglobulin-like receptor genes combined with hepatitis C viral genotype is predictive of hepatocellular carcinoma. *Hum Immunol* 2013; **74**: 1288-1294 [PMID: 23756163 DOI: 10.1016/j.humimm.2013.05.007]
- 37 Detonating a viral time bomb--the hepatitis C pandemic. *Lancet* 2013; **381**: 178 [PMID: 23332946 DOI: 10.1016/S0140-6736(13)60070-7]
- 38 **Silini E**, Bottelli R, Asti M, Bruno S, Candusso ME, Brambilla S, Bono F, Iamoni G, Tinelli C, Mondelli MU, Ideo G. Hepatitis C virus genotypes and risk of hepatocellular carcinoma in cirrhosis: a case-control study. *Gastroenterology* 1996; **111**: 199-205 [PMID: 8698200 DOI: 10.1053/gast.1996.v111.pm8698200]
- 39 **Silini E**, Bono F, Cividini A, Cerino A, Bruno S, Rossi S, Belloni G, Brugnetti B, Civardi E, Salvaneschi L. Differential distribution of hepatitis C virus genotypes in patients with and without liver function abnormalities. *Hepatology* 1995; **21**: 285-290 [PMID: 7843695]
- 40 **Nousbaum JB**, Pol S, Nalpas B, Landais P, Berthelot P, Bréchet C. Hepatitis C virus type 1b (II) infection in France and Italy. Collaborative Study Group. *Ann Intern Med* 1995; **122**: 161-168 [PMID: 7810932 DOI: 10.7326/0003-4819-122-3-199502010-00001]
- 41 **Shiha G**, Mikhail NNH, Soliman R, Hassan A, Eslam M. Predictive performance and clinical utility of HCC risk scores in chronic hepatitis C: a comparative study. *Hepatol Int* 2022; **16**: 159-170 [PMID: 35034266 DOI: 10.1007/s12072-021-10284-6]
- 42 **Lee MH**, Hsiao TI, Subramaniam SR, Le AK, Vu VD, Trinh HN, Zhang J, Jin M, Wong VW, Wong GL, Nguyen MH. HCV Genotype 6 Increased the Risk for Hepatocellular Carcinoma Among Asian Patients With Liver Cirrhosis. *Am J Gastroenterol* 2017; **112**: 1111-1119 [PMID: 28440303 DOI: 10.1038/ajg.2017.123]
- 43 **Ioannou GN**, Green PK, Beste LA, Mun EJ, Kerr KF, Berry K. Development of models estimating the risk of hepatocellular carcinoma after antiviral treatment for hepatitis C. *J Hepatol* 2018; **69**: 1088-1098 [PMID: 30138686 DOI: 10.1016/j.jhep.2018.07.024]
- 44 **Sharma SA**, Kowgier M, Hansen BE, Brouwer WP, Maan R, Wong D, Shah H, Khalili K, Yim C, Heathcote EJ, Janssen HLA, Sherman M, Hirschfield GM, Feld JJ. Toronto HCC risk index: A validated scoring system to predict 10-year risk of HCC in patients with cirrhosis. *J Hepatol* 2017 [PMID: 28844936 DOI: 10.1016/j.jhep.2017.07.033]
- 45 **Flemming JA**, Yang JD, Vittinghoff E, Kim WR, Terrault NA. Risk prediction of hepatocellular carcinoma in patients with cirrhosis: the ADDRESS-HCC risk model. *Cancer* 2014; **120**: 3485-3493 [PMID: 25042049 DOI: 10.1002/cncr.28832]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [office@baishideng.com](mailto:office@baishideng.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

