

World Journal of *Clinical Pediatrics*

Quarterly Volume 13 Number 1 March 9, 2024



EDITORIAL

Jain S. 'Prediabetes' as a practical distinctive window for workable fruitful wonders: Prevention and progression alert as advanced professionalism. *World J Clin Pediatr* 2024; 13(1): 89201 [DOI: [10.5409/wjcp.v13.i1.89201](https://doi.org/10.5409/wjcp.v13.i1.89201)]

MINIREVIEWS

Hudson AS, Wahbeh GT, Zheng HB. Imaging and endoscopic tools in pediatric inflammatory bowel disease: What's new? *World J Clin Pediatr* 2024; 13(1): 89091 [DOI: [10.5409/wjcp.v13.i1.89091](https://doi.org/10.5409/wjcp.v13.i1.89091)]

Elghoudi A, Zourob D, Al Atrash E, Alshamsi F, Alkatheeri M, Narchi H, Bitar R. Evolving strategies: Enhancements in managing eosinophilic esophagitis in pediatric patients. *World J Clin Pediatr* 2024; 13(1): 89580 [DOI: [10.5409/wjcp.v13.i1.89580](https://doi.org/10.5409/wjcp.v13.i1.89580)]

ORIGINAL ARTICLE

Case Control Study

Murillo Galvis M, Ortegon Ochoa S, Plata García CE, Valderrama Junca MP, Inga Ceballos DA, Mora Gómez DM, Granados CM, Rondón M. Exclusive breastfeeding greater than 50%, success of education in a university hospital in Bogotá: Case-control study. *World J Clin Pediatr* 2024; 13(1): 87713 [DOI: [10.5409/wjcp.v13.i1.87713](https://doi.org/10.5409/wjcp.v13.i1.87713)]

Atef Abdelsattar Ibrahim H, Mohsen M, Salep Aziz Hanna B, Mahmoud D, Mohamed Abdelhamid El-Khashab K. Childhood asthma biomarkers including zinc: An exploratory cross-sectional study. *World J Clin Pediatr* 2024; 13(1): 87866 [DOI: [10.5409/wjcp.v13.i1.87866](https://doi.org/10.5409/wjcp.v13.i1.87866)]

Metwali WA, Elmashad AM, Hazzaa SME, Al-Beltagi M, Hamza MB. Salivary C-reactive protein and mean platelet volume as possible diagnostic markers for late-onset neonatal pneumonia. *World J Clin Pediatr* 2024; 13(1): 88645 [DOI: [10.5409/wjcp.v13.i1.88645](https://doi.org/10.5409/wjcp.v13.i1.88645)]

Kalashnikova E, Isupova E, Gaidar E, Sorokina L, Kaneva M, Masalova V, Dubko M, Kornishina T, Lubimova N, Kuchinskaya E, Chikova I, Raupov R, Kalashnikova O, Kostik M. BCD020 rituximab bioanalogue compared to standard treatment in juvenile systemic lupus erythematosus: The data of 12 months case-control study. *World J Clin Pediatr* 2024; 13(1): 89049 [DOI: [10.5409/wjcp.v13.i1.89049](https://doi.org/10.5409/wjcp.v13.i1.89049)]

Retrospective Cohort Study

Isa HM, Isa AJ, Alnasheet MA, Mansoor MM. Fever assessment in children under five: Are we following the guidelines? *World J Clin Pediatr* 2024; 13(1): 88864 [DOI: [10.5409/wjcp.v13.i1.88864](https://doi.org/10.5409/wjcp.v13.i1.88864)]

Belozerov KE, Solomatina NM, Isupova EA, Kuznetsova AA, Kostik MM. Systemic juvenile idiopathic arthritis-associated lung disease: A retrospective cohort study. *World J Clin Pediatr* 2024; 13(1): 88912 [DOI: [10.5409/wjcp.v13.i1.88912](https://doi.org/10.5409/wjcp.v13.i1.88912)]

Observational Study

Suksantilerd S, Thawatchai R, Rungrojjananon N. Prevalence of vitamin D deficiency in exclusively breastfed infants at Charoenkrung Pracharak Hospital. *World J Clin Pediatr* 2024; 13(1): 86693 [DOI: [10.5409/wjcp.v13.i1.86693](https://doi.org/10.5409/wjcp.v13.i1.86693)]

Zein MM, Arafa N, El-Shabrawi MHF, El-Koofy NM. Effect of nutrition-related infodemics and social media on maternal experience: A nationwide survey in a low/middle income country. *World J Clin Pediatr* 2024; 13(1): 89139 [DOI: [10.5409/wjcp.v13.i1.89139](https://doi.org/10.5409/wjcp.v13.i1.89139)]

Manokaran K, Spaan J, Cataldo G, Lyons C, Mitchell PD, Sare T, Zimmerman LA, Rufo PA. Inpatient management of iron deficiency anemia in pediatric patients with inflammatory bowel disease: A single center experience. *World J Clin Pediatr* 2024; 13(1): 89318 [DOI: [10.5409/wjcp.v13.i1.89318](https://doi.org/10.5409/wjcp.v13.i1.89318)]

El Mouzan M, Al Sarkhy A, Assiri A. Gut microbiota predicts the diagnosis of ulcerative colitis in Saudi children. *World J Clin Pediatr* 2024; 13(1): 90755 [DOI: [10.5409/wjcp.v13.i1.90755](https://doi.org/10.5409/wjcp.v13.i1.90755)]

SYSTEMATIC REVIEWS

Al-Beltagi M, Saeed NK, Bediwy AS, Elbeltagi R, Hamza MB. Gastrointestinal tolerability of organic infant formula compared to traditional infant formula: A systematic review. *World J Clin Pediatr* 2024; 13(1): 88783 [DOI: [10.5409/wjcp.v13.i1.88783](https://doi.org/10.5409/wjcp.v13.i1.88783)]

Avendaño-Vásquez CJ, Villamizar-Osorio ML, Niño-Peñaranda CJ, Medellín-Olaya J, Reina-Gamba NC. Sociodemographic determinants associated with breastfeeding in term infants with low birth weight in Latin American countries. *World J Clin Pediatr* 2024; 13(1): 89086 [DOI: [10.5409/wjcp.v13.i1.89086](https://doi.org/10.5409/wjcp.v13.i1.89086)]

LETTER TO THE EDITOR

Grauslund AC, Lindkvist EB, Thorsen SU, Ballegaard S, Faber J, Svensson J, Berg AK. Pressure pain sensitivity: A new stress measure in children and adolescents with type 1 diabetes? *World J Clin Pediatr* 2024; 13(1): 89619 [DOI: [10.5409/wjcp.v13.i1.89619](https://doi.org/10.5409/wjcp.v13.i1.89619)]

ABOUT COVER

Peer Reviewer of *World Journal of Clinical Pediatrics*, Fateen Ata, MSc, MBBS, ABHS (Internal Medicine) Academic Editor, Academic Fellow, Academic Research, Doctor, Endocrinology, Hamad Medical Corporation, Qatar. docfateenata@gmail.com

AIMS AND SCOPE

The primary aim of the *World Journal of Clinical Pediatrics* (WJCP, *World J Clin Pediatr*) is to provide scholars and readers from various fields of pediatrics with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCP mainly publishes articles reporting research results and findings obtained in the field of pediatrics and covering a wide range of topics including anesthesiology, cardiology, endocrinology, gastroenterology, hematology, immunology, infections and infectious diseases, medical imaging, neonatology, nephrology, neurosurgery, nursing medicine, perinatology, pharmacology, respiratory medicine, and urology.

INDEXING/ABSTRACTING

The WJCP is now abstracted and indexed in PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The WJCP's CiteScore for 2022 is 1.7 and Scopus CiteScore rank 2022: Pediatrics, perinatology and child health is 176/306.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Yu-Qing Zhao*; Production Department Director: *Xiang Li*; Editorial Office Director: *Xu Guo*.

NAME OF JOURNAL

World Journal of Clinical Pediatrics

ISSN

ISSN 2219-2808 (online)

LAUNCH DATE

June 8, 2012

FREQUENCY

Quarterly

EDITORS-IN-CHIEF

Toru Watanabe, Consolato M Sergi, Elena Daniela Serban, Surjit Singh

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2219-2808/editorialboard.htm>

PUBLICATION DATE

March 9, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Observational Study

Gut microbiota predicts the diagnosis of ulcerative colitis in Saudi children

Mohammad El Mouzan, Ahmed Al Sarkhy, Asaad Assiri

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Ding X, China

Received: December 12, 2023

Peer-review started: December 12, 2023

First decision: December 19, 2023

Revised: January 1, 2024

Accepted: February 6, 2024

Article in press: February 6, 2024

Published online: March 9, 2024



Mohammad El Mouzan, Ahmed Al Sarkhy, Asaad Assiri, Department of Pediatrics, Gastroenterology Unit, King Saud University, Riyadh 11461, Saudi Arabia

Corresponding author: Mohammad El Mouzan, MD, Full Professor, Department of Pediatrics, King Saud University, 1, King Abdullah Street, Riyadh 11461, Saudi Arabia.
melmouzan@ksu.edu.sa

Abstract

BACKGROUND

Ulcerative colitis (UC) is an immune-mediated chronic inflammatory condition with a worldwide distribution. Although the etiology of this disease is still unknown, the understanding of the role of the microbiota is becoming increasingly strong.

AIM

To investigate the predictive power of the gut microbiota for the diagnosis of UC in a cohort of newly diagnosed treatment-naïve Saudi children with UC.

METHODS

The study population included 20 children with a confirmed diagnosis of UC and 20 healthy controls. Microbial DNA was extracted and sequenced, and shotgun metagenomic analysis was performed for bacteria and bacteriophages. Biostatistics and bioinformatics demonstrated significant dysbiosis in the form of reduced alpha diversity, beta diversity, and significant difference of abundance of taxa between children with UC and control groups. The receiver operating characteristic curve, a probability curve, was used to determine the difference between the UC and control groups. The area under the curve (AUC) represents the degree of separability between the UC group and the control group. The AUC was calculated for all identified bacterial species and for bacterial species identified by the random forest classification algorithm as important potential biomarkers of UC. A similar method of AUC calculation for all bacteriophages and important species was used.

RESULTS

The median age and range were 14 (0.5-21) and 12.9 (6.8-16.3) years for children with UC and controls, respectively, and 40% and 35% were male for children with UC and controls, respectively. The AUC for all identified bacterial species was 89.5%. However, when using the bacterial species identified as important by

random forest classification algorithm analysis, the accuracy increased to 97.6%. Similarly, the AUC for all the identified bacteriophages was 87.4%, but this value increased to 94.5% when the important bacteriophage biomarkers were used.

CONCLUSION

The very high to excellent AUCs of fecal bacterial and viral species suggest the potential use of noninvasive microbiota-based tests for the diagnosis of unusual cases of UC in children. In addition, the identification of important bacteria and bacteriophages whose abundance is reduced in children with UC suggests the potential of preventive and adjuvant microbial therapy for UC.

Key Words: Ulcerative colitis; Microbiota; Area under the curve; Children; Saudi Arabia

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This study reports the predictive power of fecal microbiota, bacteria and bacteriophages, in predicting the diagnosis of ulcerative colitis in children. This was demonstrated by the calculation of the area under the receiver operating characteristic curve (AUC). High values of the AUC up to 97.6% and 94.5% for bacteria and bacteriophage, respectively, indicate excellent predictive power in differentiating children with ulcerative colitis (UC) from controls. This finding may lead to the development of noninvasive microbiota-based test for the diagnosis of unusual cases of UC in children.

Citation: El Mouzan M, Al Sarkhy A, Assiri A. Gut microbiota predicts the diagnosis of ulcerative colitis in Saudi children. *World J Clin Pediatr* 2024; 13(1): 90755

URL: <https://www.wjgnet.com/2219-2808/full/v13/i1/90755.htm>

DOI: <https://dx.doi.org/10.5409/wjcp.v13.i1.90755>

INTRODUCTION

Ulcerative colitis (UC) is an immune-mediated inflammatory bowel disease. Although the incidence of this disease is highest in Western populations, it is increasing globally[1-3]. The etiology of UC is unknown; however, multifactorial factors involving interactions between genetics, host immunity, the mucosal barrier, and the gut microbiome are highly suspected[4-6]. The role of the microbiota has been extensively reported mainly in Western populations, with strong evidence of an association with UC.

In Saudi Arabia, a developing country in transition, the incidence and clinical patterns of UC have been reported[7-10]. In addition, the microbiota profile of Saudi children with Crohn's disease (CD) has been reported to be significantly associated with not only the presence of bacteria but also the high area under curve (AUC) for bacteria in fecal samples, suggesting high accuracy in predicting the diagnosis of CD[11,12]. However, there are no reports on the predictive power of the microbiota for the diagnosis of UC. The objective of this study was to evaluate the role of the microbiota in predicting the diagnosis of UC in Saudi children.

MATERIALS AND METHODS

Study population

Children with a confirmed diagnosis of UC were enrolled in the study. The children were recruited from multiple hospitals in Riyadh, Kingdom of Saudi Arabia. The inclusion criteria included new-onset and untreated disease, as well as no antibiotic exposure for at least 6 months before stool collection. Fecal samples from the children with UC were collected before bowl preparation. Healthy school children were randomly selected as controls. Stool samples from children with UC and controls were collected in cryovials without fixatives or stabilizers and immediately stored at -80°C until analysis.

DNA extraction and sequencing

Bacterial and viral DNA from fecal samples was isolated using the QIAGEN DNeasy PowerSoil Pro Kit according to the manufacturer's protocol. DNA libraries were prepared using the Nextera XT DNA Library Preparation Kit (Illumina) and IDT Unique Dual Indexes with a total DNA input of 1ng. Library were subsequently sequenced on an Illumina NovaSeq S4 platform.

Statistical and bioinformatics analysis

Shannon alpha diversity metrics were calculated in R using the R package "vegan". Wilcoxon rank-sum tests were performed between groups using the R package ggsignif[13,14]. Bray-Curtis dissimilarity was calculated in R using the

vegan package with the function `vegdist`, and PCoA tables were generated using the `ape` function `pcoa`. PERMANOVA tests for each distance matrix were generated using the `vegan`'s `function adonis2`, and beta dispersion was calculated and compared using the ANOVA method for the `betadisper` function from `vegan`[15]. DESeq2 was used to estimate differential abundance between cohorts based on count data[16]. The random forest classification algorithm was applied to the relative abundance data to predict bacterial and viral species biomarkers that might improve prediction[17].

The receiver operating characteristic (ROC) curve was used to determine the difference between the UC and control groups. The area under the curve (AUC) represents the degree of separability between the UC group and the control group. The AUC was calculated for all identified bacterial and bacteriophage species in this study and for bacterial and bacteriophage species identified by the random forest classification algorithm as important potential biomarkers of UC [18].

Ethical aspects: The study was approved by the Institutional Board Review of the College of Medicine, King Saud University in Riyadh, Kingdom of Saudi Arabia [No: 10/2647/IRB,26/6/2010]. Guardians and/or children signed informed consent and/or assent before enrollment in the study.

RESULTS

The median age and range were 14 (0.5-21) and 12.9 (6.8-16.3) years for children with UC and controls, respectively, and 40% and 35% were male for children with UC and controls, respectively. A high number of significant bacterial and bacteriophage dysbiosis events were found (unpublished data). Among these, 11 bacterial species biomarkers were identified. These included the *Bifidobacterium angulatum*, *Alistipes putredinis*, *Bacteroides caccae*, and *Bifidobacterium adolescentis* (Table 1). Similarly, among the high number of bacteriophages, four were identified as biomarkers. These included the *Salmonella* phage SEN4, *Streptococcus* phage YMC-2011, and uncultured *crAssphage* (Table 2).

The AUC for all identified bacterial species was 89.5% (79.1%-100.0%), but when based on the biomarkers, the accuracy increased to 97.6% (94.2%-100.0%) indicating very good to excellent predictive power (Figure 1). Similarly, the AUC for all the identified bacteriophages was 87.4% (75.9%-98.8%), but the AUC increased to 94.5% (87.8%-100%), when the identified important species were used, indicating very good to excellent predictive power (Figure 2).

DISCUSSION

Shotgun metagenomic analysis of bacterial and viral bacteriophage species in fecal samples of children with new-onset untreated UC revealed significant differential abundances between the UC group and the control group, indicating significant dysbiosis (unpublished data). The AUC of the ROC curve represents the degree of separability between the UC group and the control group, indicating the predictive power of the ROC curve for UC diagnosis.

In this study, we calculated the AUC based not only on the entire bacterial species and bacteriophages but also on important species identified by the random forest classification algorithm. The calculated AUC based on the abundance of all the bacterial species increased from 89.5% to 97.6% when only 11 bacterial species biomarkers were considered, indicating increased predictive power of the important bacterial species biomarkers. Similarly, the AUC calculated based on the bacteriophages increased from 87.4% to 94.5% when only four biomarkers were considered, indicating that the use of these bacteriophage biomarkers has greater predictive power for distinguishing UC patients from controls. The excellent predictive power of these biomarkers indicates the potential for the development of microbiota-based diagnostic tests. Among the bacteria and bacteriophages, *Bifidobacterium angulatum* and uncultured *crAssphage* had the highest median importance scores. *Bifidobacterium angulatum* is a species that belongs to the *Bifidobacterium* genus that is known to modulate the immune system and may be considered protective against UC[19,20]. *Uncultured crAssphage* is the most abundant human-associated virus and is found in the gut virome in approximately 50% of humans. This virus infects species of *Bacteroides* with mostly beneficial effects on health. Accordingly, *Bifidobacterium angulatum* and uncultured *crAssphage* could constitute the basis of prophylactic or therapeutic options[21-24].

The excellent predictive diagnostic power for UC in this report is slightly greater but consistent with the 93% accuracy for UC diagnosis reported within a multiclass disease in an adult study in Hong Kong[25] and the 91% accuracy in a group of children with UC in which shotgun metagenomic bacterial species-level abundance was used[26]. Finally, the 84.4% to 95% predictive power of the bacteriophage species identified in this study has not been reported thus far and deserves further study.

Study limitations: This study had a relatively small sample size, but it may be acceptable for this is the first study to use metagenomic analysis in a non-Western childhood population to determine the accuracy of the microbiota in predicting the diagnosis of UC.

CONCLUSION

The very high to excellent AUCs of fecal bacterial and viral species indicate the potential for the development of noninvasive microbiota-based tests for the diagnosis of UC and for preventive and adjuvant microbial therapy for UC. In addition, the identification of important bacteria and bacteriophages whose abundance is reduced in children with UC suggests the potential of preventive and adjuvant microbial therapy.

Table 1 Bacterial microbiota biomarkers importance score

S. No.	Bacterial species	Mean	Median	Minimum	Maximum	Decision
1	<i>Alistipes communis</i>	3.199	3.222	1.528	5.055	Confirmed
2	<i>Alistipes putredinis</i>	6.748	7.094	3.605	8.565	Confirmed
3	<i>Bacteroides caccae</i>	5.914	6.28	2.717	7.552	Confirmed
4	<i>Bifidobacterium adolescentis</i>	5.843	6.123	3.073	7.578	Confirmed
5	<i>Bifidobacterium angulatum</i>	8.89	9.47	4.265	10.827	Confirmed
6	<i>Bifidobacterium bifidum</i>	4.138	4.293	1.512	5.794	Confirmed
7	<i>Bifidobacterium catenulatum</i>	5.544	5.823	2.246	7.352	Confirmed
8	<i>Dialister succinatiphilus</i>	3.418	3.594	-0.47	4.86	Confirmed
9	<i>Peptostreptococcus stomatis</i>	3.367	3.411	1.358	4.983	Confirmed
10	<i>Prevotella copri</i>	3.826	3.812	1.463	5.595	Confirmed
11	<i>Streptococcus_u_s</i>	3.987	3.93	1.595	6.232	Confirmed

Table 2 Viral microbiota biomarkers scores

S. No.	Bacteriophage	Mean	Median	Minimum	Maximum	Decision
1	<i>Salmonella phage SEN4</i>	5.311	5.474	2.349	8.294	Confirmed
2	<i>Siphoviridae_u_s</i>	7.224	7.591	3.16	10.1	Confirmed
3	<i>Streptococcus phage YMC-2011</i>	7.989	8.611	3.409	11.18	Confirmed
4	<i>uncultured crAssphage</i>	18.35	20.11	6.433	23.25	Confirmed




Figure 1 The predictive power of fecal bacteriome. A: Area under the curve (AUC) based on the entire bacterial species shows 89.5% (79.1%–100%CI) accuracy in predicting ulcerative colitis (UC); B: Random forest algorithm was performed on the entire dataset to identify important features significantly predictive of UC increased the AUC to 97.6% (94.2–100%CI).




Figure 2 The predictive power of fecal bacteriophage. A: Area under the curve (AUC) based on the entire bacteriophage shows of 87.4% (75.9%–98.8%) in predicting ulcerative colitis (UC) in stool samples; B: Random forest algorithm was performed on the entire dataset to identify important features significantly predictive of UC increased the AUC to 94.5% (87.8%–100%CI).

ARTICLE HIGHLIGHTS

Research background

Microbiota dysbiosis has been reported in patients with ulcerative colitis (UC).

Research motivation

The role of the microbiota in predicting UC has rarely been reported.

Research objectives

To evaluate the predictive power of fecal bacteria and bacteriophages for diagnosing UC in children.

Research methods

Metagenomic analysis of bacterial and bacteriophage DNA in the stool of children with newly diagnosed UC. The area under the curve (AUC) was calculated to evaluate the predictive power of the total bacteria and bacteriophages, and random forest analysis was used to identify important microbes for distinguishing UC patients from controls.

Research results

The discriminatory power of the entire bacterial species (AUC: 89.5%) and bacteriophages (AUC: 87.4%) was very high. The random forest classification algorithm analysis revealed the excellent predictive power of important bacterial species (AUC: 97.6%) and bacteriophages (AUC: 94.5%).

Research conclusions

The very high to excellent AUCs of fecal bacterial and viral species indicate the potential for the development of noninvasive microbiota-based tests for the diagnosis of UC in children. In addition, the identification of important bacteria and bacteriophages whose abundance is reduced in children with UC suggests the potential of preventive and adjuvant microbial therapy for UC.

Research perspectives

Future research in this area with larger sample sizes is needed to clarify the role of the microbiota in the diagnosis, prevention, and treatment of UC.

FOOTNOTES

Author contributions: El Mouzan M designed and supervised the study and wrote the manuscript; Al Sarkhy A and Assiri A participated equally in recruiting participants and revising the manuscript draft; All authors have read and approved the final manuscript.

Supported by Researchers Supporting Project, King Saud University, Riyadh, Saudi Arabia, No. RSPD2024R864.

Institutional review board statement: The study was approved by the Institutional Board Review of the College of Medicine, King Saud University in Riyadh, Kingdom of Saudi Arabia [No: 10/2647/IRB,26/6/2010]. Guardians and/or children signed informed consent and/or assent before enrollment in the study.

Informed consent statement: Guardians and/or children signed informed consent and/or assent before enrollment in the study.

Conflict-of-interest statement: All authors have no conflicts of interest to disclose.

Data sharing statement: Datasets are available from the corresponding author at email: drmouzan@gmail.com.

STROBE statement: The authors have read the STROBE Statement – checklist of items, and the manuscript was prepared and revised according to the STROBE Statement – checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Saudi Arabia

ORCID number: Mohammad El Mouzan 0000-0001-8699-3143; Ahmed Al Sarkhy 0000-0002-1424-5784; Asaad Assiri 0000-0003-3357-5794.

S-Editor: Liu JH

L-Editor: A

P-Editor: Zhao YQ

REFERENCES

- Molodecky NA, Soon IS, Rabi DM, Ghali WA, Ferris M, Chernoff G, Benchimol EI, Panaccione R, Ghosh S, Barkema HW, Kaplan GG. Increasing incidence and prevalence of the inflammatory bowel diseases with time, based on systematic review. *Gastroenterology* 2012; **142**: 46-54.e42; quiz e30 [PMID: 22001864 DOI: 10.1053/j.gastro.2011.10.001]
- Malaty HM, Fan X, Opekun AR, Thibodeaux C, Ferry GD. Rising incidence of inflammatory bowel disease among children: a 12-year study. *J Pediatr Gastroenterol Nutr* 2010; **50**: 27-31 [PMID: 19934770 DOI: 10.1097/MPG.0b013e3181b99baa]
- Khan R, Kuenzig ME, Benchimol EI. Epidemiology of Pediatric Inflammatory Bowel Disease. *Gastroenterol Clin North Am* 2023; **52**: 483-496 [PMID: 37543395 DOI: 10.1016/j.gtc.2023.05.001]
- Turpin W, Goethel A, Bedrani L, Croitoru Mdc K. Determinants of IBD Heritability: Genes, Bugs, and More. *Inflamm Bowel Dis* 2018; **24**: 1133-1148 [PMID: 29701818 DOI: 10.1093/ibd/izy085]
- Kayama H, Okumura R, Takeda K. Interaction Between the Microbiota, Epithelia, and Immune Cells in the Intestine. *Annu Rev Immunol* 2020; **38**: 23-48 [PMID: 32340570 DOI: 10.1146/annurev-immunol-070119-115104]
- Zihni C, Mills C, Matter K, Balda MS. Tight junctions: from simple barriers to multifunctional molecular gates. *Nat Rev Mol Cell Biol* 2016; **17**: 564-580 [PMID: 27353478 DOI: 10.1038/nrm.2016.80]
- El Mouzan MI, Saadah O, Al-Saleem K, Al Edreesi M, Hasosah M, Alanazi A, Al Mofarreh M, Asery A, Al Qourain A, Nouli K, Al Hussaini A, Telmesani A, AlReheili K, Alghamdi S, Alrobiaa N, Alzaben A, Mehmadi A, Al Hebbi H, Al Sarkhy A, Al Mehaidib A, Al Saleem B, Assiri A, Wali S. Incidence of pediatric inflammatory bowel disease in Saudi Arabia: a multicenter national study. *Inflamm Bowel Dis* 2014; **20**: 1085-1090 [PMID: 24788219 DOI: 10.1097/MIB.0000000000000048]
- Saadah OI, El Mouzan M, Al Mofarreh M, Al Mehaidib A, Al Edreesi M, Hasosah M, Al-Hussaini A, AlSaleem K. Characteristics of Pediatric Crohn's Disease in Saudi Children: A Multicenter National Study. *Gastroenterol Res Pract* 2016; **2016**: 7403129 [PMID: 26858752 DOI: 10.1155/2016/7403129]
- AlSaleem K, El Mouzan MI, Saadah OI, AlSaleem B, Al-Hussaini A, Hassosa M, Ali AM, Banemai MO, Halaby H, El Edreesi M. Characteristics of pediatric ulcerative colitis in Saudi Arabia: a multicenter national study. *Ann Saudi Med* 2015; **35**: 19-22 [PMID: 26142933 DOI: 10.5144/0256-4947.2015.19]
- Al-Hussaini A, El Mouzan M, Hasosah M, Al-Mehaidib A, AlSaleem K, Saadah OI, Al-Edreesi M. Clinical Pattern of Early-Onset Inflammatory Bowel Disease in Saudi Arabia: A Multicenter National Study. *Inflamm Bowel Dis* 2016; **22**: 1961-1970 [PMID: 27104817 DOI: 10.1097/MIB.0000000000000796]
- El Mouzan MI, Winter HS, Assiri AA, Korolev KS, Al Sarkhy AA, Dowd SE, Al Mofarreh MA, Menon R. Microbiota profile in new-onset pediatric Crohn's disease: data from a non-Western population. *Gut Pathog* 2018; **10**: 49 [PMID: 30519287 DOI: 10.1186/s13099-018-0276-3]
- El Mouzan MI, Winter HS, Al Sarkhy AA, Korolev K, Menon R, Assiri AA. Bacterial dysbiosis predicts the diagnosis of Crohn's disease in Saudi children. *Saudi J Gastroenterol* 2021; **27**: 144-148 [PMID: 33642351 DOI: 10.4103/sjg.SJG_409_20]
- Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlenn D, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH,

- Szoecs E, Wagner H. *vegan: Community Ecology Package* 2019. R package version 2.5-6. Accessed May 15, 2023. Available from: <https://CRAN.R-project.org/package=vegan>
- 14 **Ahlmann-Eltze C.** *ggsignif: Significance Brackets for 'ggplot2'* 2019. R package version 0.6.0. Accessed May 15, 2023. Available from: <https://CRAN.R-project.org/package=ggsignif>
- 15 **Wickham H.** *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York, 2016
- 16 **Love MI, Huber W, Anders S.** Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014; **15**: 550 [PMID: 25516281 DOI: 10.1186/s13059-014-0550-8]
- 17 **Kursa MB, Rudnicki WR.** "Feature Selection with the Boruta Package. *J Stat Softw* 2010; **36**: 1-13 [DOI: 10.18637/jss.v036.i11]
- 18 **Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, Müller M.** pROC: an open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics* 2011; **12**: 77 [PMID: 21414208 DOI: 10.1186/1471-2105-12-77]
- 19 **Henrick BM, Rodriguez L, Lakshmikanth T, Pou C, Henckel E, Arzoomand A, Olin A, Wang J, Mikes J, Tan Z, Chen Y, Ehrlich AM, Bernhardtsson AK, Mugabo CH, Ambrosiani Y, Gustafsson A, Chew S, Brown HK, Prambs J, Bohlin K, Mitchell RD, Underwood MA, Smilowitz JT, German JB, Frese SA, Brodin P.** Bifidobacteria-mediated immune system imprinting early in life. *Cell* 2021; **184**: 3884-3898.e11 [PMID: 34143954 DOI: 10.1016/j.cell.2021.05.030]
- 20 **de Vrese M, Winkler P, Rautenberg P, Harder T, Noah C, Laue C, Ott S, Hampe J, Schreiber S, Heller K, Schrezenmeier J.** Effect of Lactobacillus gasseri PA 16/8, Bifidobacterium longum SP 07/3, B. bifidum MF 20/5 on common cold episodes: a double blind, randomized, controlled trial. *Clin Nutr* 2005; **24**: 481-491 [PMID: 16054520 DOI: 10.1016/j.clnu.2005.02.006]
- 21 **Aldars-García L, Chaparro M, Gisbert JP.** Systematic Review: The Gut Microbiome and Its Potential Clinical Application in Inflammatory Bowel Disease. *Microorganisms* 2021; **9** [PMID: 33946482 DOI: 10.3390/microorganisms9050977]
- 22 **Wiredu Ocansey DK, Hang S, Yuan X, Qian H, Zhou M, Valerie Olovo C, Zhang X, Mao F.** The diagnostic and prognostic potential of gut bacteria in inflammatory bowel disease. *Gut Microbes* 2023; **15**: 2176118 [PMID: 36794838 DOI: 10.1080/19490976.2023.2176118]
- 23 **Manandhar I, Alimadadi A, Aryal S, Munroe PB, Joe B, Cheng X.** Gut microbiome-based supervised machine learning for clinical diagnosis of inflammatory bowel diseases. *Am J Physiol Gastrointest Liver Physiol* 2021; **320**: G328-G337 [PMID: 33439104 DOI: 10.1152/ajpgi.00360.2020]
- 24 **Koonin EV, Yutin N.** The crAss-like Phage Group: How Metagenomics Reshaped the Human Virome. *Trends Microbiol* 2020; **28**: 349-359 [PMID: 32298613 DOI: 10.1016/j.tim.2020.01.010]
- 25 **Su Q, Liu Q, Lau RI, Zhang J, Xu Z, Yeoh YK, Leung TWH, Tang W, Zhang L, Liang JQY, Yau YK, Zheng J, Liu C, Zhang M, Cheung CP, Ching JYL, Tun HM, Yu J, Chan FKL, Ng SC.** Faecal microbiome-based machine learning for multi-class disease diagnosis. *Nat Commun* 2022; **13**: 6818 [PMID: 36357393 DOI: 10.1038/s41467-022-34405-3]
- 26 **Zuo W, Wang B, Bai X, Luan Y, Fan Y, Michail S, Sun F.** 16S rRNA and metagenomic shotgun sequencing data revealed consistent patterns of gut microbiome signature in pediatric ulcerative colitis. *Sci Rep* 2022; **12**: 6421 [PMID: 35440670 DOI: 10.1038/s41598-022-07995-7]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

