

World Journal of *Gastroenterology*

World J Gastroenterol 2024 April 14; 30(14): 1934-2067



EDITORIAL

- 1934 Topic highlight on texture and color enhancement imaging in gastrointestinal diseases
Toyoshima O, Nishizawa T, Hata K
- 1941 Immune checkpoint inhibitor-associated gastritis: Patterns and management
Lin J, Lin ZQ, Zheng SC, Chen Y
- 1949 Liver biopsy in the post-hepatitis C virus era in Japan
Ikura Y, Okubo T, Sakai Y
- 1958 Current status of liver transplantation for human immunodeficiency virus-infected patients in mainland China
Tang JX, Zhao D
- 1963 Bowel function and inflammation: Is motility the other side of the coin?
Panarese A

REVIEW

- 1968 Necroptosis contributes to non-alcoholic fatty liver disease pathoetiology with promising diagnostic and therapeutic functions
Sun HJ, Jiao B, Wang Y, Zhang YH, Chen G, Wang ZX, Zhao H, Xie Q, Song XH

MINIREVIEWS

- 1982 Omics-based biomarkers as useful tools in metabolic dysfunction-associated steatotic liver disease clinical practice: How far are we?
Trinks J, Mascardi MF, Gadano A, Marciano S

ORIGINAL ARTICLE

Retrospective Study

- 1990 Characteristics of early gastric tumors with different differentiation and predictors of long-term outcomes after endoscopic submucosal dissection
Zhu HY, Wu J, Zhang YM, Li FL, Yang J, Qin B, Jiang J, Zhu N, Chen MY, Zou BC

Observational Study

- 2006 Preoperative albumin-bilirubin score and liver resection percentage determine postoperative liver regeneration after partial hepatectomy
Takahashi K, Gosho M, Miyazaki Y, Nakahashi H, Shimomura O, Furuya K, Doi M, Owada Y, Ogawa K, Ohara Y, Akashi Y, Enomoto T, Hashimoto S, Oda T

Basic Study

- 2018** *Fusobacterium nucleatum*-induced imbalance in microbiome-derived butyric acid levels promotes the occurrence and development of colorectal cancer
Wu QL, Fang XT, Wan XX, Ding QY, Zhang YJ, Ji L, Lou YL, Li X
- 2038** Comparative transcriptomic analysis reveals the molecular changes of acute pancreatitis in experimental models
Zheng P, Li XY, Yang XY, Wang H, Ding L, He C, Wan JH, Ke HJ, Lu NH, Li NS, Zhu Y

CASE REPORT

- 2059** Outcomes of endoscopic sclerotherapy for jejunal varices at the site of choledochojejunostomy (with video): Three case reports
Liu J, Wang P, Wang LM, Guo J, Zhong N

ABOUT COVER

Editorial Board Member of *World Journal of Gastroenterology*, Sandro Contini, MD, Former Associate Professor, Department of Surgical Sciences, University of Parma, Parma 43123, Italy. sandrocontini46@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Gastroenterology* (WJG, *World J Gastroenterol*) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE), MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJG as 4.3; Quartile category: Q2. The WJG's CiteScore for 2021 is 8.3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*, Production Department Director: *Xiang Li*, Cover Editor: *Jia-Ru Fan*.

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Andrzej S Tarnawski

EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF

Xian-Jun Yu (Pancreatic Oncology), Jian-Gao Fan (Chronic Liver Disease), Hou-Bao Liu (Biliary Tract Disease)

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

PUBLICATION DATE

April 14, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

PUBLISHING PARTNER

Shanghai Pancreatic Cancer Institute and Pancreatic Cancer Institute, Fudan University
Biliary Tract Disease Institute, Fudan University

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/bpg/GerInfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

PUBLISHING PARTNER's OFFICIAL WEBSITE

<https://www.shca.org.cn>
<https://www.zs-hospital.sh.cn>



Current status of liver transplantation for human immunodeficiency virus-infected patients in mainland China

Jian-Xin Tang, Dong Zhao

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Majhi S, India

Received: December 28, 2023

Peer-review started: December 28, 2023

First decision: February 9, 2024

Revised: February 18, 2024

Accepted: March 21, 2024

Article in press: March 21, 2024

Published online: April 14, 2024



Jian-Xin Tang, Dong Zhao, Department of Liver Surgery and Organ Transplantation Center, Shenzhen Third People's Hospital, The Second Affiliated Hospital of Southern University of Science and Technology, National Clinical Research Center for Infectious Disease, Shenzhen 518000, Guangdong Province, China

Corresponding author: Dong Zhao, MD, Chief Doctor, Department of Liver Surgery and Organ Transplantation Center, Shenzhen Third People's Hospital, The Second Affiliated Hospital of Southern University of Science and Technology, National Clinical Research Center for Infectious Disease, No. 29 Bulan Road, Longgang District, Shenzhen 518000, Guangdong Province, China. zdong1233@126.com

Abstract

According to the report from the Chinese Center for Disease Control and Prevention, the prevalence of human immunodeficiency virus (HIV) infection exceeded 1.2 million individuals by the year 2022, with an annual increase of about 80000 cases. The overall prevalence of hepatitis B surface antigen among individuals co-infected with HIV reached 13.7%, almost twice the rate of the general population in China. In addition to the well-documented susceptibility to opportunistic infections and new malignancies, HIV infected patients frequently experience liver-related organ damage, with the liver and kidneys being the most commonly affected. This often leads to the development of end-stage liver and kidney diseases. Therefore, organ transplantation has emerged as an important part of active treatment for HIV infected patients. However, the curative effect is not satisfactory. HIV infection has been considered a contraindication for organ transplantation. Until the emergence of highly active anti-retroviral therapy in 1996, the once intractable replication of retrovirus was effectively inhibited. With prolonged survival, the failure of important organs has become the main cause of death among HIV patients. Therefore, transplant centers worldwide have resumed exploration of organ transplantation for HIV-infected individuals and reached a positive conclusion. This study provides an overview of the current landscape of HIV-positive patients receiving liver transplantation (LT) in mainland China. To date, our transplant center has conducted LT for eight end-stage liver disease patients co-infected with HIV, and all but one, who died two months postoperatively due to sepsis and progressive multi-organ failure, have survived. Comparative analysis with hepatitis B virus-infected patients during the same period revealed no statistically significant differences in acute rejection reactions, cytomegalovirus infection, bacteremia, pulmonary infections, acute kidney injury, new-onset cancers, or vascular and biliary complications.

Key Words: Liver transplantation; Human immunodeficiency virus; Infection; Hepatitis B virus; End-stage liver disease; Mainland China

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: In mainland China, human immunodeficiency virus (HIV) infection has long been considered an absolute contraindication for liver transplantation (LT). Until June 2004, Chinese scholars performed LT on two HIV patients, but these cases were only followed up for 24 and 22 months, respectively. Subsequently, there were no reported cases of LT in mainland China for patients with HIV infection until April 2019, when our transplant center performed ABO-incompatible LT for a patient with liver failure co-infected with HIV and hepatitis B virus. This study provides an overview of the current landscape of HIV-positive patients receiving liver transplantation in mainland China.

Citation: Tang JX, Zhao D. Current status of liver transplantation for human immunodeficiency virus-infected patients in mainland China. *World J Gastroenterol* 2024; 30(14): 1958-1962

URL: <https://www.wjgnet.com/1007-9327/full/v30/i14/1958.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v30.i14.1958>

INTRODUCTION

Human immunodeficiency virus (HIV) remains a significant global health concern, with the latest data from the Joint United Nations Programme on HIV/acquired immunodeficiency syndrome (AIDS) indicating a staggering 38 million individuals infected worldwide in 2021. Approximately 4000 people contract HIV daily, resulting in an annual increase of 1.5 million new infections[1]. According to a report from the Chinese Center for Disease Control and Prevention, it is projected that by 2023, over 1.2 million individuals in China will be living with HIV, with an annual increment of around 80000 cases (<https://www.chinacdc.cn>. Accessed 30 November 2023). Since the advent of highly active antiretroviral therapy (HAART) in 1996, substantial progress has been made in controlling HIV infection, managing opportunistic infections, reducing mortality rates, and prolonging life expectancy among infected individuals[2]. HIV/AIDS has become a chronic, manageable condition. However, liver-related mortality remains a formidable challenge, particularly for HIV-positive individuals undergoing HAART, who are increasingly susceptible to complications associated with co-infections, notably hepatitis C virus (HCV) and hepatitis B virus (HBV), leading to hepatocellular carcinoma, end-stage liver disease (ESLD), or alcohol-related liver disease[3,4]. Given the shared transmission routes, co-infections with chronic HCV and chronic HBV are prevalent[5].

GLOBAL TRENDS IN LIVER TRANSPLANTATION FOR HIV-POSITIVE PATIENTS

Liver transplantation (LT) is the only effective method for treating various ESLDs. Initially, HIV-positive patients were excluded from LT due to concerns about their compromised immune system, potential acceleration of HIV progression with immunosuppressive drugs, and limited organ resources[6,7]. However, by the year 2000, as the number of HIV-positive patients requiring LT significantly increased, the focus shifted towards addressing ESLD and end-stage renal disease in these individuals. The United States and the United Kingdom initiated two pilot trials for LT in HIV-positive individuals, yielding favorable outcomes without evidence of immunosuppressive drugs exacerbating the progression of HIV[8]. Then, a prospective multicenter study confirmed that LT in HIV patients exhibited excellent patient and graft outcomes[9].

In 2020, a large retrospective study enrolling LT recipients from the Organ Procurement and Transplantation Network and the Scientific Registry of Transplant Recipients from 2008 to 2015 elucidated the survival trends and characteristics of HIV-infected LT recipients[10]. Among 73206 LT patients, 658 (0.9%) were HIV-infected. From 2008 to 2011, patients survival rates at 1 year and 3 years for the HIV-infected group were 78.2% and 64.1%, respectively. From 2012 to 2015, the 1-year and 3-year survival rates for the HIV-infected group improved significantly to 86.2% and 75.4%, respectively. The cumulative graft survival rates during 2012-2015 also showed substantial improvement, likely attributed to the rational use of anti-HIV drugs, immunosuppressants, and advancements in surgical techniques. Currently, LT outcomes and survival for HIV-positive recipients are comparable to those of HIV-negative recipients[11,12]. However, during the years 2012-2015, the 3-year graft survival rates for HIV-infected recipients with and without concurrent HCV infection were 62.6% and 84.7%, respectively, with similar results observed in patient survival rates. HIV/HCV co-infected recipients exhibited significantly lower survival rates than HCV mono-infected patients, possibly due to a higher rate of post-transplant HCV recurrence leading to graft failure and recipient mortality[13,14]. Cohort studies from France and Spain indicated a 5-year survival rate below 55% for HIV/HCV co-infected LT recipients[15,16]. Therefore, the question of whether LT should be pursued for this patient population remains contentious. To comprehensively evaluate the long-term postoperative survival of HIV-positive LT recipients, a retrospective cohort study with a follow-up exceeding 10

years examined 180 cases in the United States from 2002 to 2011[17]. At 5 years after transplantation (64.8% for HIV+/HCV- *vs* 51.8% for HIV+/HCV+, $P = 0.15$) and 10 years after transplantation (43.9% for HIV+/HCV- *vs* 44.1% for HIV+/HCV+, $P = 0.20$), there was no statistically significant difference in long-term survival rates between mono-infected (HIV+/HCV-) and co-infected (HIV+/HCV+) recipients. However, when compared to the general transplant population with HCV-positive recipients, the long-term survival and graft survival rates for HIV/HCV co-infected recipients after transplantation were suboptimal. Therefore, further research is necessary to investigate the long-term postoperative outcomes for HIV/HCV co-infected LT recipients.

In contrast to co-infection with HCV, HIV-infected patients concurrently affected by HBV appear to have a more favorable prognosis after LT when an appropriate HBV recurrence prevention strategy is implemented. Optimal prevention for recurrent HBV infection in HIV/HBV co-infected patients seems to involve the combination of antiviral therapy with hepatitis B immunoglobulin (HBIG)[18]. In a prospective multicenter study in the United States, with passive prophylaxis using HBIG and antiviral treatment, there was no significant difference in graft and patient survival rates between HIV/HBV co-infected and HBV mono-infected LT recipients, with 3-year survival rates reaching 85%[11]. Studies by Anadol *et al*[19] demonstrated a 5-year survival rate of 80% for HIV/HBV co-infected patients, with no clinically relevant HBV-related ESLD observed after LT. Tateo *et al*[20] reported a cumulative graft and patient survival rate of 100% at an average follow-up of 32 months for 13 HIV/HBV co-infected LT recipients. To assess the postoperative prognosis and the risk of complications for HIV/HBV co-infected patients, we reviewed the literature on LT in HBV/HIV co-infected patients, revealing 1-year and 3-year cumulative survival rates of 85.9% and 77.3%, respectively. Moreover, due to the high recurrence rate of HBV viremia, lifelong prophylactic medication is recommended[21].

LIVER TRANSPLANTATION FOR HIV-POSITIVE PATIENTS IN MAINLAND CHINA

According to a cross-sectional study in China, the hepatitis B surface antigen infection rate among HIV-infected individuals can reach 14.5%, nearly three times higher than the general population[22]. In mainland China, HIV infection has long been considered an absolute contraindication for LT. The advent of HAART in 1996 improved the prognosis for HIV-infected individuals and encouraged some transplant centers in China to accept organ transplant candidates who were HIV positive. Until June 2004, mainland Chinese scholars performed LT on two HIV patients[23], but these cases were only followed up for 24 months and 22 months, respectively. Subsequently, there were no reported cases of LT in mainland China for patients with concomitant HIV infection until April 2019, when our transplant center performed LT for a patient with liver failure co-infected with HIV and HBV. This case is the first reported instance of ABO-incompatible LT for HIV/HBV co-infection[21]. As of the latest follow-up, the patient has survived healthily for over 4 years, with normal liver enzyme levels and no transplant rejection reactions. To date, our transplant center has conducted LT for eight ESLD patients co-infected with HIV, and all but one, who died two months postoperatively due to sepsis and progressive multi-organ failure, have survived. There were no statistically significant differences in acute rejection reactions, cytomegalovirus infection, bacteremia, pulmonary infections, acute kidney injury, new-onset cancers, and vascular or biliary complications compared to HBV-infected patients during the same period. The preoperative CD4(+) T cell count for the first ABO-incompatible patient in our preliminary report was only 42 cells/ μ L, while the preoperative CD4(+) T cell counts for the remaining patients were all below 200 cells/ μ L. With the exception of one case resulting in mortality, all other patients have maintained good health postoperatively.

CONCLUSION

Our study findings suggest that the criteria for preoperative CD4(+) T cell counts in LT recipients co-infected with HIV still require further exploration. Also, a comprehensive assessment beyond preoperative CD4(+) T cell counts is essential to ensure the safety of LT recipients in this category. Despite the relatively low number of cases of LT in HIV-infected individuals in mainland China, postoperative survival rates are comparable to those in Western countries. Given the limited number of cases included in the study, our next step will involve expanding the sample size and conducting more in-depth clinical research.

FOOTNOTES

Author contributions: Zhao D contributed to study conception and design, and administrative support; Zhao D and Tang JX contributed to provision of the study materials, and collection and assembly of the data; and all authors contributed to the writing and final approval of the manuscript.

Supported by the Shenzhen Science and Technology R&D Fund, No. JCYJ20220530163011026; and Shenzhen Third People's Hospital, No. G2022008 and No. G2021008.

Conflict-of-interest statement: No conflict-of-interest to disclose.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to

distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Jian-Xin Tang 0000-0003-4416-5336; Dong Zhao 0000-0003-3773-721X.

S-Editor: Chen YL

L-Editor: A

P-Editor: Yuan YY

REFERENCES

- Ostankova YV**, Shchemelev AN, Thu HHK, Davydenko VS, Reingardt DE, Serikova EN, Zueva EB, Totolian AA. HIV Drug Resistance Mutations and Subtype Profiles among Pregnant Women of Ho Chi Minh City, South Vietnam. *Viruses* 2023; **15** [PMID: 37896785 DOI: 10.3390/v15102008]
- Rowell-Cunsolo TL**, Hu G, Bellerose M, Liu J. Trends in Comorbidities Among Human Immunodeficiency Virus-Infected Hospital Admissions in New York City from 2006-2016. *Clin Infect Dis* 2021; **73**: e1957-e1963 [PMID: 33245318 DOI: 10.1093/cid/ciaa1760]
- Smith CJ**, Ryom L, Weber R, Morlat P, Pradier C, Reiss P, Kowalska JD, de Wit S, Law M, el Sadr W, Kirk O, Friis-Moller N, Monforte Ad, Phillips AN, Sabin CA, Lundgren JD; D:A:D Study Group. Trends in underlying causes of death in people with HIV from 1999 to 2011 (D:A:D): a multicohort collaboration. *Lancet* 2014; **384**: 241-248 [PMID: 25042234 DOI: 10.1016/S0140-6736(14)60604-8]
- Puri P**, Kumar S. Liver involvement in human immunodeficiency virus infection. *Indian J Gastroenterol* 2016; **35**: 260-273 [PMID: 27256434 DOI: 10.1007/s12664-016-0666-8]
- Abdi M**, Ahmadi A, Mokarizadeh A. Biomarkers for Assessment of Human Immunodeficiency Virus and its Co-Infection with Hepatitis B and Hepatitis C Viruses: A Comprehensive Review. *Iran J Pathol* 2023; **18**: 230-243 [PMID: 37942194 DOI: 10.30699/IJP.2023.1972384.3009]
- Stock P**, Roland M, Carlson L, Freise C, Hirose R, Terrault N, Frassetto L, Coates T, Roberts J, Ascher N. Solid organ transplantation in HIV-positive patients. *Transplant Proc* 2001; **33**: 3646-3648 [PMID: 11750549 DOI: 10.1016/s0041-1345(01)02569-6]
- Roland ME**, Stock PG. Liver transplantation in HIV-infected recipients. *Semin Liver Dis* 2006; **26**: 273-284 [PMID: 16850377 DOI: 10.1055/s-2006-947297]
- Mohsen AH**, Easterbrook P, Taylor CB, Norris S. Hepatitis C and HIV-1 coinfection. *Gut* 2002; **51**: 601-608 [PMID: 12235089 DOI: 10.1136/gut.51.4.601]
- Roland ME**, Barin B, Huprikar S, Murphy B, Hanto DW, Blumberg E, Olthoff K, Simon D, Hardy WD, Beatty G, Stock PG; HIVTR Study Team. Survival in HIV-positive transplant recipients compared with transplant candidates and with HIV-negative controls. *AIDS* 2016; **30**: 435-444 [PMID: 26765937 DOI: 10.1097/QAD.0000000000000934]
- Campos-Varela I**, Dodge JL, Berenguer M, Adam R, Samuel D, Di Benedetto F, Karam V, Belli LS, Duvoux C, Terrault NA. Temporal Trends and Outcomes in Liver Transplantation for Recipients With HIV Infection in Europe and United States. *Transplantation* 2020; **104**: 2078-2086 [PMID: 32969987 DOI: 10.1097/TP.00000000000003107]
- Coffin CS**, Stock PG, Dove LM, Berg CL, Nissen NN, Curry MP, Ragni M, Regenstein FG, Sherman KE, Roland ME, Terrault NA. Virologic and clinical outcomes of hepatitis B virus infection in HIV-HBV coinfecting transplant recipients. *Am J Transplant* 2010; **10**: 1268-1275 [PMID: 20346065 DOI: 10.1111/j.1600-6143.2010.03070.x]
- Jacob JS**, Shaikh A, Goli K, Rich NE, Benhammou JN, Ahmed A, Kim D, Rana A, Goss JA, Naggie S, Lee TH, Kanwal F, Cholaneril G. Improved Survival After Liver Transplantation for Patients With Human Immunodeficiency Virus (HIV) and HIV/Hepatitis C Virus Coinfection in the Integrase Strand Transfer Inhibitor and Direct-Acting Antiviral Eras. *Clin Infect Dis* 2023; **76**: 592-599 [PMID: 36221143 DOI: 10.1093/cid/ciac821]
- Kardashian AA**, Price JC. Hepatitis C virus-HIV-coinfecting patients and liver transplantation. *Curr Opin Organ Transplant* 2015; **20**: 276-285 [PMID: 25944240 DOI: 10.1097/MOT.0000000000000199]
- Manzardo C**, Londoño MC, Castells L, Testillano M, Luis Montero J, Peñafiel J, Subirana M, Moreno A, Aguilera V, Luisa González-Diéguez M, Calvo-Pulido J, Xiol X, Salcedo M, Cuervas-Mons V, Manuel Sousa J, Suarez F, Serrano T, Ignacio Herrero J, Jiménez M, Fernandez JR, Giménez C, Del Campo S, Esteban-Mur JJ, Crespo G, de la Rosa G, Rimola A, Miro JM; FIPSE LT-HIV investigators. Direct-acting antivirals are effective and safe in HCV/HIV-coinfecting liver transplant recipients who experience recurrence of hepatitis C: A prospective nationwide cohort study. *Am J Transplant* 2018; **18**: 2513-2522 [PMID: 29963780 DOI: 10.1111/ajt.14996]
- Duclos-Vallée JC**, Féray C, Sebah M, Teicher E, Roque-Afonso AM, Roche B, Azoulay D, Adam R, Bismuth H, Castaing D, Vittecoq D, Samuel D; THEVIC Study Group. Survival and recurrence of hepatitis C after liver transplantation in patients coinfecting with human immunodeficiency virus and hepatitis C virus. *Hepatology* 2008; **47**: 407-417 [PMID: 18098295 DOI: 10.1002/hep.21990]
- Miro JM**, Montejó M, Castells L, Rafecas A, Moreno S, Agüero F, Abradelo M, Miralles P, Torre-Cisneros J, Pedreira JD, Cordero E, de la Rosa G, Moyano B, Moreno A, Perez I, Rimola A; Spanish OLT in HIV-Infected Patients Working Group investigators. Outcome of HCV/HIV-coinfecting liver transplant recipients: a prospective and multicenter cohort study. *Am J Transplant* 2012; **12**: 1866-1876 [PMID: 22471341 DOI: 10.1111/j.1600-6143.2012.04028.x]
- Locke JE**, Durand C, Reed RD, MacLennan PA, Mehta S, Massie A, Nellore A, DuBay D, Segev DL. Long-term Outcomes After Liver Transplantation Among Human Immunodeficiency Virus-Infected Recipients. *Transplantation* 2016; **100**: 141-146 [PMID: 26177090 DOI: 10.1097/TP.0000000000000829]
- Terrault NA**, Carter JT, Carlson L, Roland ME, Stock PG. Outcome of patients with hepatitis B virus and human immunodeficiency virus infections referred for liver transplantation. *Liver Transpl* 2006; **12**: 801-807 [PMID: 16628690 DOI: 10.1002/Lt.20776]
- Anadol E**, Beckebaum S, Radecke K, Paul A, Zoufaly A, Bickel M, Hitzenbichler F, Ganten T, Kittner J, Stoll M, Berg C, Manekeller S, Kalff JC, Sauerbruch T, Rockstroh JK, Spengler U. Orthotopic liver transplantation in human-immunodeficiency-virus-positive patients in Germany. *AIDS Res Treat* 2012; **2012**: 197501 [PMID: 22900154 DOI: 10.1155/2012/197501]

- 20 **Tateo M**, Roque-Afonso AM, Antonini TM, Medja F, Lombes A, Jardel C, Teicher E, Sebah M, Roche B, Castaing D, Samuel D, Duclos-Vallee JC. Long-term follow-up of liver transplanted HIV/hepatitis B virus coinfecting patients: perfect control of hepatitis B virus replication and absence of mitochondrial toxicity. *AIDS* 2009; **23**: 1069-1076 [PMID: 19417577 DOI: 10.1097/QAD.0b013e32832c2a37]
- 21 **Tang JX**, Zhang KJ, Fang TS, Weng RH, Liang ZM, Yan X, Jin X, Xie LJ, Zeng XC, Zhao D. Outcomes of ABO-incompatible liver transplantation in end-stage liver disease patients co-infected with hepatitis B and human immunodeficiency virus. *World J Gastroenterol* 2023; **29**: 1745-1756 [PMID: 37077518 DOI: 10.3748/wjg.v29.i11.1745]
- 22 **Xie J**, Han Y, Qiu Z, Li Y, Song X, Wang H, Thio CL, Li T. Prevalence of hepatitis B and C viruses in HIV-positive patients in China: a cross-sectional study. *J Int AIDS Soc* 2016; **19**: 20659 [PMID: 26979535 DOI: 10.7448/ias.19.1.20659]
- 23 **Shen ZY**, Zheng H, Deng YL, Zhu ZJ, Pan C. Two cases of liver transplantation in patients infected with human immunodeficiency virus. *Chin J Organ Transplant* 2007; **28**: 250 [DOI: 10.1097/mot.0b013e3282f1fb9]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

