



Contribution of the toxic AGEs (TAGE)-RAGE axis in NASH-related HCC



学术搜索

获得 1 条结果 (用时0.07秒)

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

英语网页

☒ 包括专利

☒ 包含引用

创建快讯

您是不是要找: Contribution of the toxic AGEs (**RAGE**)-RAGE axis in NASH-related HCC

[HTML] CARCINOMA-BASIC

Y Xu, X Qin, J Zhou, Z Tu, P Morales, Al Haza - 2012 - doko.vn

... This book is an important contribution to the field of hepatocellular carcinoma with an ... 55353, Figure 1) harboring this fragment was performed by Expressed Sequence Tag (EST) splicing, 5 ... grades, TNM staging, portal vein invasion and recurrence, but not with age, gender, viral ...

相关文章 引用 保存 更多

创建快讯

doko.vn 中的 [HTML]

您是不是要找: Contribution of the toxic AGEs (**RAGE**)-RAGE axis in NASH-related HCC

[关于 Google 学术搜索](#)

[隐私权](#)

[条款](#)

[提供反馈](#)

World Journal of Gastroenterology

ESPS Manuscript NO: 17674

Manuscript Type: REVIEW

Contribution of the toxic advanced glycation end-product-receptor axis in nonalcoholic steatohepatitis-related hepatocellular carcinoma

Takino J *et al.* Contribution of TAGE in HCC

Jun-ichi Takino, Kentaro Nagamine, Takamitsu Hori, Akiko Sakasai-Sakai, Masayoshi Takeuchi

Jun-ichi Takino, Kentaro Nagamine, Takamitsu Hori, Department of Biochemistry, Faculty of Pharmaceutical Sciences, Hiroshima International University, Hiroshima 737-0112, Japan

Akiko Sakasai-Sakai, Masayoshi Takeuchi, Department of Advanced Medicine, Medical

Match Overview

1	Internet 97 words crawled on 30-Mar-2015 www.wjnet.com	1%
2	CrossCheck 84 words Hideyuki Hyogo, "Advanced Glycation End Products (A ... Es) and their Involvement in Liver Disease", <i>Current Pha</i>	1%
3	Internet 80 words crawled on 13-Oct-2009 www.bentham.org	1%
4	CrossCheck 60 words Hideyuki Hyogo, "Serum Advanced Glycation End Pro ... cts Associated with NASH and Other Liver Diseases", <i>En</i>	1%
5	Internet 39 words crawled on 04-Oct-2010 www.hindawi.com	<1%
6	CrossCheck 30 words Takeuchi, Masayoshi, Jun-ichi Takino, Satomi Furuno, Hi kari Shirai, Mihoko Kawakami, Michiru Muramatsu, Yuka	<1%
7	CrossCheck 25 words M. Takeuchi, "Alternative Routes for the Formation of In ... munochemically Distinct Advanced Glycation End-produ	<1%
8	CrossCheck 24 words Jun-ichi Takino, "The formation of intracellular glycerali ... hyde-derived advanced glycation end-products and cytot	<1%
9	Internet 23 words crawled on 22-Apr-2014 www.kanazawa-med.ac.jp	<1%



Contribution of the toxic advanced glycation end-product-receptor axis in no



学术搜索

 我的著作引用情况



文章

您的搜索 - **Contribution of the toxic advanced glycation end-product-receptor axis in nonalcoholic steatohepatitis-related hepatocellular carcinoma** - 与所有文章均不相符。

我的图书馆

建议:

请检查输入字词有无错误。
请尝试其他的查询词
请改用较常见的字词。
请减少查询字词的数量。
[请向所有网络查询](#)

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

[关于 Google 学术搜索](#)

[隐私权](#)

[条款](#)

[提供反馈](#)

按日期排序


搜索所有网页

中文网页

英语网页

☒ 包括专利

☒ 包含引用

 创建快讯

[网页](#)[图片](#)[新闻](#)[购物](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 12,400 条结果 (用时 0.82 秒)

Role of receptor for advanced glycation end products ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - [翻译此页](#)

作者 : S Yamagishi - 2015 - [相关文章](#)

2015年2月11日 - Receptor for advanced glycation end products (RAGE) belongs to a ... non-ischemic liver injury, and hepatocellular carcinoma growth and ... contribute to the development and progression of numerous types of hepatic disorders [11,12]. ... non-progressive condition, to non-alcoholic steatohepatitis (NASH), ...

Involvement of the TAGE-RAGE system in non-alcoholic ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : M Takeuchi - 2014 - 被引用次数 : 1 - [相关文章](#)

2014年12月27日 - Here, we examine the contributions of RAGE and TAGE to various conditions and ... Core tip: Toxic advanced glycation end-products (TAGE) synthesis is increased by non-alcoholic steatohepatitis (NASH), and patients with ... which can result in cirrhosis, liver failure, and/or hepatocellular carcinoma[2-4].

Soluble receptor for advanced glycation end products and ...

onlinelibrary.wiley.com/doi/10.1002/hep.26264/pdf ▾ [翻译此页](#)

作者 : KA Moy - 2013 - 被引用次数 : 22 - [相关文章](#)

Binding of advanced glycation end products (AGEs) to their receptor (RAGE) ... sRAGE or N-(carboxymethyl)-lysine (CML)-AGE and hepatocellular carcinoma in a case- ... the AGE-RAGE axis in the development of HCC is ... sRAGE, soluble receptor for advanced glycation end product; NASH, nonalcoholic steatohepatitis.

[PDF] Toxic AGEs (TAGE) Theory in the Pathogenesis of NAFL...

clinmedjournals.org/articles/ijdc/ijdc-2-036.pdf ▾ [翻译此页](#)

2015年7月31日 - and hepatocellular carcinoma. ... derivatives termed "advanced glycation end-products, AGEs" (GA- ... AGEs, RAGE: Receptor for AGEs, NAFLD: