65736_Auto_Edited v3.docx

Quotes Excluded Bibliography Excluded

14%

iThenticate[®]

Name of Journal: World Journal of Gastroenterology

Manuscript NO: 65736

Manuscript Type: OPINION REVIEW

Screening for nonalcoholic fatty liver disease - when, who and how?

Screening for NAFLD

Christoph G Dietrich, Monika Rau, Andreas Geier





Screening for nonalcoholic fatty liver disease - when, who and how







ALL IMAGES VIDEOS

Add the Give with Bing extension >

48,400,000 Results

Screening for Nonalcoholic Fatty Liver Disease in ...

https://pubmed.ncbi.nlm.nih.gov/29889226

Any time ▼

Background: Inflammatory bowel disease (IBD) patients may be at risk for nonalcoholic fatty liver disease (NAFLD) due to chronic inflammation, hepatotoxic drugs, and alteration of the gut microbiota. Prospective data using accurate diagnostic methods are lacking. Methods: We prospectively investigated prevalence and predictors of NAFLD and liver fibrosis by transient elastography (TE) with ...

Cited by: 26 Author: Chiara Saroli Palumbo, Sophie Restellini,...

Publish Year: 2019

Screening for Nonalcoholic Fatty Liver Disease in the ...

https://pubmed.ncbi.nlm.nih.gov/31391806

Nonalcoholic **fatty liver disease** (NAFLD) is a spectrum of diseases ranging from simple hepatic steatosis to nonalcoholic steatohepatitis (NASH) to advanced fibrosis and cirrhosis as well as **liver** cancer. Despite the significant morbidity associated with NAFLD, there are no global consensus guideline

Cited by: 9

Author: Vijay Pandyarajan, Robert G Gish, Naim ...

Publish Year: 2019

PEOPLE ALSO ASK

How is the diagnosis of nonalcoholic fatty liver disease made?




V

How is ultrasonography used to diagnose fatty liver disease?

Fatty Liver

Medical Condition



A condition that occurs due to excessive build-up of fat in the liver.

- Very common (More than 3 million cases per year in US)
- No vaccine available
- A Requires lab test or imaging
- Treatable by a medical professional
- Can last source works or months