

# World Journal of *Clinical Cases*

*World J Clin Cases* 2023 December 26; 11(36): 8434-8605



**EDITORIAL**

- 8434 Post-trans-arterial chemoembolization hepatic necrosis and biliary stenosis: Clinical characteristics and endoscopic approach  
*Cocca S, Carloni L, Marocchi M, Grande G, Bianchini M, Colecchia A, Conigliaro R, Bertani H*

**MINIREVIEWS**

- 8440 Perioperative nursing care for hip arthroplasty patients with concomitant hypertension: A minireview  
*Ji CY, Yang LR*

**ORIGINAL ARTICLE****Retrospective Study**

- 8447 Evaluation of response to gemcitabine plus cisplatin-based chemotherapy using positron emission computed tomography for metastatic bladder cancer  
*Öztürk H, Karapolat İ*
- 8458 Functional magnetic resonance imaging study of group independent components underpinning item responses to paranoid-depressive scale  
*Stoyanov D, Paunova R, Dichev J, Kandilarova S, Khorev V, Kurkin S*

**EVIDENCE-BASED MEDICINE**

- 8475 Mendelian randomization provides evidence for a causal effect of serum insulin-like growth factor family concentration on risk of atrial fibrillation  
*Lin S, Tang J, Li X, Wu G, Lin YF, Li YF*

**SYSTEMATIC REVIEWS**

- 8486 Significance of fostering the mental health of patients with diabetes through critical time intervention  
*Eseadi C, Amedu AN, Aloh HE*

**META-ANALYSIS**

- 8498 Impact of angiotensin-converting enzyme inhibitors or angiotensin receptor blockers on the mortality in sepsis: A meta-analysis  
*Yang DC, Xu J, Jian L, Yu Y*

**CASE REPORT**

- 8507 Multiple sparganosis spinal infections mainly in the thoracic region: A case report  
*Wen GJ, Chen J, Zhang SF, Zhou ZS, Jiao GL*

- 8512** Iatrogenic flexor tendon rupture caused by misdiagnosing sarcoidosis-related flexor tendon contracture as tenosynovitis: A case report  
*Yan R, Zhang Z, Wu L, Wu ZP, Yan HD*
- 8519** Cholecystoenteric fistula in a patient with advanced gallbladder cancer: A case report and review of literature  
*Wang CY, Chiu SH, Chang WC, Ho MH, Chang PY*
- 8527** Intraperitoneal hyaline vascular Castleman disease: Three case reports  
*Gao JW, Shi ZY, Zhu ZB, Xu XR, Chen W*
- 8535** Iris metastasis from clear cell renal cell carcinoma: A case report  
*Wang TT, Chen XY, Min QY, Han YZ, Zhao HF*
- 8542** Spinal cord infarction attributed to SARS-CoV-2, with post-acute sequelae of COVID-19: A case report  
*Oleson CV, Olsen AC, Shermom S*
- 8551** Spontaneous gastric hematoma as a rare cause of acute abdomen: A case report  
*Budimir I, Žulec M, Eljuga K, Židak M, Lisek V*
- 8557** LiNA OperaScope™ for microwave endometrial ablation for endometrial polyps with heavy menstrual bleeding: A case report  
*Kakinuma K, Kakinuma T, Ueyama K, Shinohara T, Okamoto R, Yanagida K, Takeshima N, Ohwada M*
- 8563** Colonoscopy-induced acute appendicitis: A case report  
*Song XL, Ma JY, Zhang ZG*
- 8568** Post-laparotomy heterotopic ossification of the xiphoid process: A case report  
*Lee SS*
- 8574** Balloon displacement during caesarean section with pernicious placenta previa: A case report  
*Gu DF, Deng C*
- 8581** Synchronous carotid endarterectomy and coronary artery bypass graft: Four case reports  
*AlGhamdi FK, Altoijry A, AlQahtani A, Aldossary MY, AlSheikh SO, Iqbal K, Alayadhi WA*
- 8589** Intraoperative cardiogenic shock induced by refractory coronary artery spasm in a patient with myasthenia gravis: A case report  
*Hsu CW, Chang CC, Lin CS*
- 8595** Effects of video game-based therapy in an adolescent with cerebral palsy: A case report  
*Mohd Iqbal HA, Ho WS, Zamudin A, Hisham H, Mohd Nordin NA*

**LETTER TO THE EDITOR**

- 8603** Lyophilized recombinant human brain natriuretic peptide: A promising therapy in patients with chronic heart failure  
*Kourek C, Briasoulis A, Giamouzis G, Skoularis J, Xanthopoulos A*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Marco Infante, MD, PhD, Adjunct Professor, UniCamillus, Saint Camillus International University of Health Sciences, Rome 00131, Italy.  
marco.infante@unicamillus.org

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

*WJCC* mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJCC* as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Editorial Office Director: Jin-Lai Wang.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogorgos, Maurizio Serati

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

December 26, 2023

**COPYRIGHT**

© 2024 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

# Intraoperative cardiogenic shock induced by refractory coronary artery spasm in a patient with myasthenia gravis: A case report

Cheng-Wei Hsu, Chuen-Chau Chang, Chao-Shun Lin

**Specialty type:** Medicine, research and experimental

**Provenance and peer review:** Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): 0  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Liu YC, China

**Received:** October 26, 2023

**Peer-review started:** October 26, 2023

**First decision:** November 22, 2023

**Revised:** November 24, 2023

**Accepted:** December 7, 2023

**Article in press:** December 7, 2023

**Published online:** December 26, 2023



**Cheng-Wei Hsu, Chuen-Chau Chang, Chao-Shun Lin**, Department of Anesthesiology, Taipei Medical University Hospital, Taipei 110, Taiwan

**Chao-Shun Lin**, Department of Anesthesiology, School of Medicine, College of Medicine, Taipei Medical University, Taipei 110, Taiwan

**Corresponding author:** Chao-Shun Lin, MD, PhD, Associate Professor, Department of Anesthesiology, Taipei Medical University Hospital, No. 252 WuXing Street, Taipei 110, Taiwan. [lin.soon@gmail.com](mailto:lin.soon@gmail.com)

## Abstract

### BACKGROUND

Coronary artery spasm (CAS) is a rare but critical condition during surgery. Clinical manifestations can vary from only subtle electrocardiography change to sudden death. In this case report, we present the case of a patient with myasthenia gravis (MG) who developed refractory CAS-related cardiogenic shock during thymoma surgery.

### CASE SUMMARY

A 61-year-old man had a history of cigarette smoking and coronary artery disease with a bare metal stent placed. Three months ago, he suffered from coronary spasms, with three vessels involved, after surgery for cervical spine injury. He started having progressive dysphagia 4 wk prior and was diagnosed with MG *via* serologic tests, and computed tomography declared a thymoma in the anterior mediastinum. After the symptoms of MG subsided, he was referred for thymectomy. The operation was uneventful until the closing of the sternal wound. Electrocardiography showed sudden onset ST elevation, followed by ventricular tachycardia and severe hypotension. Cardiopulmonary cerebral resuscitation was initiated immediately with electrical defibrillation, extracorporeal membrane oxygenation was performed due to refractory cardiogenic shock, and the patient was transferred to an angiography room. Angiography showed diffuse CAS with three vessels involved. Intracoronary isosorbide dinitrate and adenosine were administered, and then the patient was transferred to the intensive care unit.

### CONCLUSION

Our case highlights the importance of being prepared for clinical situations such as the one described here and suggests the necessity of developing an appropriate anesthesia plan that includes proactive analgesia and preemptive coronary vasodilators.

**Key Words:** Coronary spasm; Myasthenia gravis; Thymectomy; Shock; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** In previous literature reviews, it has been noted that a correlation exists between myasthenia gravis (MG) and cardiac complications, such as coronary artery spasm (CAS), which frequently manifests as chest pain in affected patients. Nevertheless, when MG coincides with thymoma, surgical intervention is often necessary. The diagnosis of CAS while the patient is under general anesthesia poses a considerable challenge. Our case report aims to underscore scenarios of this nature and suggests an optimal anesthesia strategy in such cases.

**Citation:** Hsu CW, Chang CC, Lin CS. Intraoperative cardiogenic shock induced by refractory coronary artery spasm in a patient with myasthenia gravis: A case report. *World J Clin Cases* 2023; 11(36): 8589-8594

**URL:** <https://www.wjgnet.com/2307-8960/full/v11/i36/8589.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v11.i36.8589>

## INTRODUCTION

Coronary artery spasm (CAS) was recently recognized as a cause of myocardial infarction with nonobstructive coronary arteries[1]. The transient cessation of the coronary blood supply causes clinical manifestations that mimic ischemic heart disease. Perioperative CAS can be challenging for anesthesiologists, especially when patients are sedated or under general anesthesia. We present a case of myasthenia gravis (MG) that developed coronary spasm-related cardiogenic shock during thymoma surgery.

## CASE PRESENTATION

### Chief complaints

A 61-year-old man was diagnosed with MG and thymoma. He was scheduled for median sternotomy to undergo resection of the thymoma.

### History of present illness

The patient started having progressive dysphagia 4 wk prior, and the associated symptoms included diplopia, ptosis, general weakness, and easy choking. He denied chest pain, bloody sputum, and fever. Serologic tests were positive for antibodies against the acetylcholine receptor (serum level: 88.3 nmol/L). Computed tomography revealed a 5.4 cm enhanced lobular mass in the anterior mediastinum, which was declared to be a thymoma. He was diagnosed with MG, and intravenous immunoglobulin was administered at a dose of 600 mg/kg once-daily for 5 consecutive days. The symptoms of MG subsided after 20 d, and he was referred for surgical intervention. Preoperative echocardiography revealed preserved systolic function and normal wall motion without major valvular dysfunction. Thus, a medium sternotomy for anterior mediastinum tumor resection was arranged.


In the operating theater, general anesthesia was induced with lidocaine 60 mg, thiamylal sodium 300 mg, fentanyl 50 µg, and rocuronium 40 mg. A 37-Fr double lumen endotracheal tube was intubated, and then an arterial line and a central venous catheter were placed smoothly. The operation was uneventful until the closing of the sternal wound. Electrocardiography showed sudden onset ST elevation, followed by ventricular tachycardia and severe hypotension. Cardiopulmonary cerebral resuscitation was initiated immediately with electrical defibrillation (200 J), and the surgeon started to perform direct cardiac massage. However, ventricular tachycardia/fibrillation and hypotension persisted. Extracorporeal membrane oxygenation (ECMO) was performed 40 min after the initiation of cardiopulmonary cerebral resuscitation.

### History of past illness

The patient had a history of cigarette smoking and coronary artery disease with a bare metal stent placed 11 years prior. In January 2021, he suffered from a cervical spine injury with disc fracture at C5–6 and central cord syndrome from a traffic accident. He underwent cervical discectomy and interbody fusion with a cervical cage. After transfer to the intensive care unit, chest pain was mentioned. Then, sudden onset bradycardia following ventricular tachycardia and cardiac arrest occurred. Coronary angiography showed spasms in three vessels without obvious atherosclerotic lesions. He was transferred to a normal ward 10 d later and then discharged after rehabilitation was completed.

### Personal and family history

The patient denied any family history of MG.



DOI: 10.12998/wjcc.v11.i36.8589 Copyright ©The Author(s) 2023.

**Figure 1** Diffuse spasm involving the left anterior descending artery and left circumflex artery (arrows).

### **Physical examination**

After ECMO insertion, the patient's arterial blood pressure was approximately 90/50 mmHg. The heart rate was around 70 beats per minute and respiratory rate was set at 12 breaths per minute. Capnography showed a low end tidal CO<sub>2</sub> level (under 20 mmHg).

### **Laboratory examinations**

Arterial blood gas data showed acidosis (PH = 7.304) with elevated PaCO<sub>2</sub> (70.4 mmHg) and HCO<sub>3</sub><sup>-</sup> (35.3 mEq/L). Serum lactate level also increased (8.5 mmol/L). Serum creatine kinase (CK) and CK muscle and brain isoenzyme levels were in normal range but troponin-T level was elevated (0.039 ng/mL).

### **Imaging examinations**

Electrocardiography indicated a return to sinus rhythm with ST elevation within 10 min. Subsequent transesophageal echocardiography exhibited global hypokinesia. Further assessment through angiography unveiled diffuse CAS with three vessels involved (Figure 1).

---

## **FINAL DIAGNOSIS**

---

Cardiogenic shock due to diffuse CAS with three vessels involved.

---

## **TREATMENT**

---


A total of 1600 mg intracoronary isosorbide dinitrate and 360 mg adenosine were administered. After transient relief, refractory spasm was noted at the right coronary artery (Figure 2); thus, a bare metal stent was placed (Figure 3). An intra-arterial balloon pump was placed due to poor contraction of the left ventricle, and then the patient was transferred to the intensive care unit.

---

## **OUTCOME AND FOLLOW-UP**


---

The patient regained consciousness on the following day. A week later, echocardiography revealed improved left ventricular systolic function; thus, the intra-arterial balloon pump and ECMO were removed. He was then transferred to a ward for a rehabilitation program and discharged. However, he had pneumonia and progressed into sepsis 5 mo later and expired due to multiorgan failure.



DOI: 10.12998/wjcc.v11.i36.8589 Copyright ©The Author(s) 2023.

Figure 2 Refractory spasm of the right coronary artery after intracoronary isosorbide dinitrate and adenosine administration (arrow).



DOI: 10.12998/wjcc.v11.i36.8589 Copyright ©The Author(s) 2023.

Figure 3 A bare metal stent was placed at the right coronary artery (arrow).

## DISCUSSION

CAS is a rare condition, and its diverse manifestations can sometimes be critical, especially perioperatively. Some risk factors for CAS have been identified, such as age, sex, smoking, and physical and mental stress, and the usage of sympathomimetic and parasympathomimetic agents can be precipitating factors. The pathophysiology of CAS can be multifactorial, including endothelial dysfunction, autonomic nervous system disorder, and oxidative stress. One study reviewed 115 cases with perioperative CAS, and most cases of CAS occurred during abdominal or thoracic surgery. The authors considered inadequate depth of general anesthesia, use of vasopressors, and vagus nerve stimulation as possible contributing factors. Most patients had normal preoperative electrocardiograms. However, almost every patient (97%) presented ST segment changes when CAS occurred, and approximately 20% were associated with ventricular fibrillation or cardiac arrest[2].

MG is an autoimmune neuromuscular disease, and antibodies to acetylcholine receptors at neuromuscular junctions cause muscle weakness. Usually, antibodies bind only to the skeletal system. However, in patients with MG combined

with thymoma, specific striational antibodies bind to heart muscle, which may be related to the myocarditis and myositis that occur in MG patients[3]. Several case reports have presented the occurrence of CAS in patients with MG after cholinesterase inhibitor or intravenous immunoglobulin treatment[4]. Acetylcholine, as a parasympathetic neurotransmitter of the endothelium, is usually related to coronary dilation; however, it can induce vasospasm through vascular smooth muscle constriction when the endothelium is damaged[5,6]. While the precise mechanism is not fully understood, MG and its treatments can influence myocardial and coronary function through different pathways, causing patients to be at risk of cardiovascular events.

To the best of our knowledge, this is the first reported case of intraoperative coronary spasm in a patient with MG who underwent thymectomy. Our case had several risk factors for CAS, including cigarette smoking and atherosclerotic coronary artery disease. He also suffered from coronary spasm after traumatic cervical spine injury. He was diagnosed with MG and administered anticholinergic and intravenous immunoglobulin treatments. Thus, we cannot simply attribute the cause to a single factor. In our case, high-quality cardiopulmonary cerebral resuscitation and successful ECMO cannulation were crucial, and the patient recovered without complications. A previous study announced that prophylactic coronary vasodilators may bring benefits that reduce the risk of CAS[7]. To avoid noxious stimuli, an adequate anesthesia depth is necessary. The epidural catheter technique is widely used in thymectomy as an effective analgesia method. However, it can also induce coronary spasm, so the risks and benefits need to be determined[2,8]. We suggest that sugammadex should be used for the reversal of neuromuscular function if postoperative extubation is indicated. In summary, the potential risk of cardiovascular events should be taken into consideration for patients with MG undergoing surgery.

---

## CONCLUSION

In past literature reviews, an association between MG and CAS has been reported. Patients with these conditions often present with chest pain. However, when MG is combined with thymoma, surgical intervention is frequently required. Diagnosing CAS under general anesthesia can be very challenging. Our case report presented a particularly devastating CAS that necessitated the use of ECMO and an intra-aortic balloon pump. We identified severe heart failure using transesophageal echocardiography and promptly closed the surgical incision to transfer the patient to the catheterization room for further treatment. Reflecting on this case, we propose an appropriate anesthesia plan, including proactive pain management, prophylactic coronary vasodilators, and always keeping the possibility of such complications as a differential diagnosis. Further research is needed to explore this complex relationship in the future.

---

## FOOTNOTES

**Author contributions:** Hus CW and Chang CC interpreted the patient data and wrote the draft of the manuscript; Lin CS wrote the manuscript as the corresponding author; all authors read and approved the final manuscript.

**Informed consent statement:** Written informed consent for publication of the clinical details and images was obtained from the relative of the patient.

**Conflict-of-interest statement:** The authors declare that they have no conflict of interest to disclose.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016) and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** Taiwan

**ORCID number:** Chao-Shun Lin [0000-0001-6964-3862](https://orcid.org/0000-0001-6964-3862).

**Corresponding Author's Membership in Professional Societies:** Taiwan Society of Cardiothoracic and Vascular Anesthesia, 0022.

**S-Editor:** Lin C

**L-Editor:** Wang TQ

**P-Editor:** Lin C

---

## REFERENCES

- 1 Pasupathy S, Tavella R, Beltrame JF. Myocardial Infarction With Nonobstructive Coronary Arteries (MINOCA): The Past, Present, and

- Future Management. *Circulation* 2017; **135**: 1490-1493 [PMID: 28416521 DOI: 10.1161/CIRCULATIONAHA.117.027666]
- 2 **Koshiba K**, Hoka S. Clinical characteristics of perioperative coronary spasm: reviews of 115 case reports in Japan. *J Anesth* 2001; **15**: 93-99 [PMID: 14566530 DOI: 10.1007/s005400170034]
  - 3 **Suzuki S**, Utsugisawa K, Yoshikawa H, Motomura M, Matsubara S, Yokoyama K, Nagane Y, Maruta T, Satoh T, Sato H, Kuwana M, Suzuki N. Autoimmune targets of heart and skeletal muscles in myasthenia gravis. *Arch Neurol* 2009; **66**: 1334-1338 [PMID: 19752287 DOI: 10.1001/archneurol.2009.229]
  - 4 **Yanagihashi M**, Okamoto R, Morioka H, Sawada M, Matsumoto S, Ikeda T, Kano O. Coronary spastic angina after the administration of intravenous immunoglobulin in myasthenia gravis: a case report. *BMC Neurol* 2020; **20**: 319 [PMID: 32859168 DOI: 10.1186/s12883-020-01901-2]
  - 5 **Sato K**, Kaikita K, Nakayama N, Horio E, Yoshimura H, Ono T, Ohba K, Tsujita K, Kojima S, Tayama S, Hokimoto S, Matsui K, Sugiyama S, Yamabe H, Ogawa H. Coronary vasomotor response to intracoronary acetylcholine injection, clinical features, and long-term prognosis in 873 consecutive patients with coronary spasm: analysis of a single-center study over 20 years. *J Am Heart Assoc* 2013; **2**: e000227 [PMID: 23858100 DOI: 10.1161/JAHA.113.000227]
  - 6 **Ishii M**, Kaikita K, Sato K, Tanaka T, Sugamura K, Sakamoto K, Izumiya Y, Yamamoto E, Tsujita K, Yamamuro M, Kojima S, Soejima H, Hokimoto S, Matsui K, Ogawa H. Acetylcholine-Provoked Coronary Spasm at Site of Significant Organic Stenosis Predicts Poor Prognosis in Patients With Coronary Vasospastic Angina. *J Am Coll Cardiol* 2015; **66**: 1105-1115 [PMID: 26337988 DOI: 10.1016/j.jacc.2015.06.1324]
  - 7 **Kurabayashi M**, Okishige K, Asano M, Suzuki H, Shimura T, Iwai S, Kato N, Ihara K, Aoyagi H, Isobe M. Cardiopulmonary arrest caused by coronary spasm after coronary vasodilator withdrawal during the peri-operative period of gastrectomy. *Intern Med* 2013; **52**: 81-84 [PMID: 23291678 DOI: 10.2169/internalmedicine.52.8918]
  - 8 **Easley RB**, Rosen RE, Lindeman KS. Coronary artery spasm during initiation of epidural anesthesia. *Anesthesiology* 2003; **99**: 1015-1017 [PMID: 14508337 DOI: 10.1097/0000542-200310000-00037]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

