

World Journal of *Clinical Cases*

World J Clin Cases 2023 December 16; 11(35): 8242-8433



Contents

Thrice Monthly Volume 11 Number 35 December 16, 2023

EDITORIAL

- 8242 Antibiotic treatment in cirrhotic patients
Fiore M, Leone S

MINIREVIEWS

- 8247 Research progress on preparation of lateral femoral tunnel and graft fixation in anterior cruciate ligament reconstruction
Dai Y, Gao WJ, Li WC, Xiang XX, Wang WM
- 8256 Accessory navicular in children
Xiang F, Liu ZQ, Zhang XP, Li YJ, Wen J
- 8263 Non-pharmacological pain palliation methods in chronic pancreatitis
Tez M, Şahingöz E, Marlı HF

ORIGINAL ARTICLE

Retrospective Study

- 8270 Ratio of hemoglobin to mean corpuscular volume: A new index for discriminating between iron deficiency anemia and thalassemia trait
Yao QC, Zhai HL, Wang HC
- 8276 Influence of standardized nursing intervention combined with mindfulness stress reduction training on the curative effect in patients with acute pancreatitis
Li S, Yin D, Guo XC
- 8284 Clinical analysis of 114 cases of bronchiolitis in infants
Shi C, Wu MH, Zuo A, Yang MM, Jiang RR
- 8291 Endovenous laser treatment *vs* conventional surgery for great saphenous vein varicosities: A propensity score matching analysis
Li Q, Zhang C, Yuan Z, Shao ZQ, Wang J
- 8300 Efficacy of prednisone combined with mycophenolate mofetil for immunoglobulin A nephropathy with moderate-to-severe renal dysfunction
Meng MJ, Hu L, Fan Y, Gao H, Chen HZ, Chen CM, Qi Z, Liu B
- 8310 Efficacy of surgical resection and ultra-reduced tension suture combined with superficial radiation in keloid treatment
Hu XY, Yang Q, Guan XY, Li JY, Wang LL, Li K, Zhang XT

Observational Study

- 8320** Prior abdominal surgery as a potential risk factor for colonic diverticulosis or diverticulitis
Ariam E, Richter V, Bermont A, Sandler Y, Cohen DL, Shirin H

META-ANALYSIS

- 8330** Vericiguat treatment of heart failure: A systematic review and meta-analysis
Yang H, Luo C, Lan WQ, Tang YH

CASE REPORT

- 8343** Rare synchronous colorectal carcinoma with three pathological subtypes: A case report and review of the literature
Li F, Zhao B, Zhang L, Chen GQ, Zhu L, Feng XL, Yao H, Tang XF, Yang H, Liu YQ
- 8350** Twin pregnancy with sudden heart failure and pulmonary hypertension after atrial septal defect repair: A case report
Tong CX, Meng T
- 8357** Diffuse arterial atherosclerosis presenting with acute ischemic gastritis: A case report
Wei RY, Zhu JH, Li X, Wu JY, Liu JW
- 8364** Balloon venoplasty for disdialysis syndrome due to pacemaker-related superior vena cava syndrome with chylothorax post-bacteraemia: A case report
Yamamoto S, Kamezaki M, Ooka J, Mazaki T, Shimoda Y, Nishihara T, Adachi Y
- 8372** Malignant pleural mesothelioma mimics thoracic empyema: A case report
Yao YH, Kuo YS
- 8379** Multifocal papillary thyroid cancer in Graves' disease: A case report
Alzaman N
- 8385** Anlotinib in combination with pembrolizumab for low-grade myofibroblastic sarcoma of the pancreas: A case report
Wu RT, Zhang JC, Fang CN, Qi XY, Qiao JF, Li P, Su L
- 8392** Ankle and toe weakness caused by calcified ligamentum flavum cyst: A case report
Jung HY, Kim GU, Joh YW, Lee JS
- 8399** Atypical case of bow hunter's syndrome linked to aberrantly coursing vertebral artery: A case report
Ahn JH, Jun HS, Kim IK, Kim CH, Lee SJ
- 8404** Phleboscrosis: An overlooked complication of varicose veins that affects clinical outcome: A case report
Ren SY, Qian SY, Gao RD
- 8411** Inflammatory cutaneous metastases originating from gastric cancer: A case report
Tian L, Ye ZB, Du YL, Li QF, He LY, Zhang HZ

- 8416** Metastatic pancreatic solitary fibrous tumor: A case report

Yi K, Lee J, Kim DU

- 8425** Abemaciclib-induced lung damage leading to discontinuation in brain metastases from breast cancer: A case report

Yamashiro H, Morii N

LETTER TO THE EDITOR

- 8431** Letter to the editor: Aggressive variant prostate cancer: An exemplary case study and comprehensive literature survey

Ke HW, Zhang WY, Xu KX

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Md Moshir Rahman, MBBS, Assistant Professor, Department of Neurosurgery, Holy Family Red Crescent Medical College Hospital, Dhaka 1000, Bangladesh. dr.tutul@yahoo.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Hua-Ge Yin; Production Department Director: Xiang Li; Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

December 16, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Clinical analysis of 114 cases of bronchiolitis in infants

Cheng Shi, Meng-Hua Wu, An Zuo, Mi-Mi Yang, Rong-Rong Jiang

Specialty type: Pediatrics

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Alsamhi SH, Ireland

Received: September 22, 2023

Peer-review started: September 22, 2023

First decision: October 9, 2023

Revised: October 12, 2023

Accepted: December 4, 2023

Article in press: December 4, 2023

Published online: December 16, 2023



Cheng Shi, Meng-Hua Wu, An Zuo, Mi-Mi Yang, Rong-Rong Jiang, Department of Pediatrics, Tongling Maternal and Child Health Hospital, Tongling 244000, Anhui Province, China

Corresponding author: Meng-Hua Wu, Doctor, Associate Chief Physician, Department of Pediatrics, Tongling Maternal and Child Health Hospital, No. 98 Yangjiashan Road, Tongguan District, Tongling 244000, Anhui Province, China. menghuawu12@163.com

Abstract

BACKGROUND

Bronchiolitis is a common lower respiratory tract infection in infants and young children. Severe cases may be accompanied by obvious dyspnea and oxygen saturation decline.

AIM

To summarize the clinical features, standard diagnosis, and treatment of bronchiolitis.

METHODS

This is a retrospective analysis of 114 pediatric patients (74 males, 40 females) who were first diagnosed as having bronchioles at the Department of Pediatrics of Tongling Maternal and Child Health Hospital from January 2019 to December 2019. The clinical features, imaging features, treatment, and other clinical data were recorded and analyzed.

RESULTS

The age of onset of the disease was mainly from 1 mo to 6 mo (75.4%), and the time to hospital visit was mostly from the 2nd day to the 4th day of the course of the disease (75.4%). Lung imaging examination showed increase in lung texture, fuzzy (93.8%). The main treatment was atomization therapy: Budesonide combined with terbutaline (45.6%) and budesonide combined with salbutamol (38.5%). The average hospitalization time was 7.1 ± 2.4 d, and the overall cure rate was 94.7%. In patients without bacterial infection, the use of antibiotics significantly prolonged the length of hospital stay (7.8 ± 2.5 d vs 5.7 ± 1.8 d) and improved the cure rate (98.3% vs 87.9%, $P < 0.05$).

CONCLUSION

Infants with bronchiolitis are mainly male and tend to have a good prognosis. However, the unneeded use of antibiotics may prolong the length of hospital stay significantly, which imposes the burden both on the patients and hospital system.

Key Words: Bronchiolitis; Diagnosis; Treatment; Infants; Antibiotics; Retrospective analysis

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: A retrospective analysis of the data of infants and young children with bronchiolitis showed that the disease had a male gender preference, the age of onset was relatively young, and the prognosis was good. Through comparative analysis of the use of antibiotics in infants and young children with bronchiolitis without bacterial infection, it was found that there was controversy over the abuse of antibiotics in the treatment of the disease. Increasing the use of antibiotics can improve the cure rate of patients, but it will prolong the length of hospital stay, and there is controversy over unreasonable use of medical resources.

Citation: Shi C, Wu MH, Zuo A, Yang MM, Jiang RR. Clinical analysis of 114 cases of bronchiolitis in infants. *World J Clin Cases* 2023; 11(35): 8284-8290

URL: <https://www.wjgnet.com/2307-8960/full/v11/i35/8284.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i35.8284>

INTRODUCTION

In infants, bronchiolitis is one of the more common lower respiratory tract infection, its clinical manifestations mainly include cough, shortness of breath, and difficulty breathing, and lung auscultation may reveal wheezing sound and wet rales. Patients with severe bronchiolitis may have obvious dyspnea and oxygen saturation decline[1]. The peak age of its onset is 2-6 mo, and it is the main cause of hospitalization in children under 24 mo old[2]. At present, it is known that bronchiolitis is mainly caused by viral infection, the most common of which is respiratory syncytial virus[3], followed by adenovirus and influenza virus[4]. The main treatment measures were symptomatic and supportive treatment such as relieving cough and asthma and resolving phlegm[5]. The prognosis of children with this disease is mostly good, and a small number of children will develop into childhood asthma. Therefore, bronchiolitis should be actively prevented and treated to reduce the occurrence of asthma.

A total of 114 hospitalized children diagnosed with bronchiolitis at the Department of Pediatrics of Tongling Maternal and Child Health Care Hospital in the past year were retrospectively analyzed, and according to the treatment experience, the hospital's diagnosis and treatment standards for the disease were formulated to save medical costs. The main flowchart of the study is shown in Figure 1. This study was approved by the Medical Ethics Committee of Tongling Maternal and Child Health Hospital, and informed consent was obtained from the children's guardians.

MATERIALS AND METHODS

General data

The clinical data of 114 children hospitalized at Tongling Maternal and Child Health Hospital from January to December 2019 with the first diagnosis of bronchiolitis were collected. All the children met the diagnostic criteria for bronchiolitis. The diagnostic criteria were based on the 8th edition of "Zhufang Practical Pediatrics". Cases with missing data or incomplete primary data were excluded, and as a result, 18 patients were excluded from the study. The criteria for discharge included patients who were either cured or showed significant improvement, had stable vital signs, and exhibited clinical symptoms and signs that had returned to normal for more than 24 hours. Additionally, a review conducted before leaving the hospital confirmed that all infection indicators were normal.

Auxiliary examination

All children underwent routine blood tests, examinations for hypersensitive C-reactive protein (CRP) and liver and kidney function, respiratory virus detection, and sputum culture checks. Children with the following conditions required further examination: (1) Serum electrolytes were measured when there were signs of diarrhea or dehydration; and (2) Blood culture was done when the body temperature was > 38.5 °C or there were symptoms of infection and poisoning. All children underwent pulmonary X-ray examination during their illness.

Treatment

The following treatments were administered: (1) All hospitalized children were randomly treated with nebulized bronchodilator and/or β_2 receptor agonist according to their condition; (2) Glucocorticoid atomization treatment was routinely used, and for severe cases, intravenous corticosteroids or intravenous immunoglobulin can be used; (3) Antibiotic treatment was given or not according to the results of the first blood routine test and CRP within 24 h before and after admission; and (4) Antiviral drugs were not used routinely.




Figure 1 Study flowchart. Bronchiolitis is a prevalent lower respiratory tract infection in infants, often characterized by pronounced dyspnea, decreased oxygen saturation, and symptoms such as cough, asthma, and phlegm. In this study, a total of 114 pediatric patients were analyzed. Routine blood tests, hypersensitive C-reactive protein assessments, and liver and kidney function tests were conducted. Notably, standard respiratory virus detection and sputum culture checks were performed. Serum electrolytes were measured in cases with diarrhea or dehydration. Blood cultures were taken when body temperatures exceeded 38.5 °C or when signs of infection or poisoning were evident. Comprehensive data, including clinical and imaging features as well as treatment details, were meticulously recorded and subsequently analyzed.

Statistical analysis

SPSS19.0 software package was applied for statistical analyses. Measurement data with a normal distribution are expressed as the mean \pm SD, and the two independent samples *t* test was used for comparison. Count data are expressed as percentages (%) or rates, and the chi-square test was used for comparison. $P < 0.05$ was considered statistically significant.

RESULTS

As shown in Table 1, among the 114 children, 74 were male and 40 were female with a male to female ratio of about 1:0.54; 71 (62.3%) were born *via* natural delivery and 43 (37.7%) *via* cesarean section. The age of onset was mainly from 1 mo to 6 mo (93.9%), the time to visit the hospital was mostly 2-4 d (79.8%), the average length of hospital stay was (7.1 ± 2.4) d, and 23 cases (20.2%) were confirmed to have bacterial infection. In the study group, there were no deaths or serious complications. Six people were transferred, and the cure rate was 94.7%.

All children underwent routine chest X-ray examination. As shown in Table 2, of 107 cases (93.8%) showed thickened and blurred lung texture, and 7 (6.1%) were combined with patchy shadows. Wheezing was found in 47 cases (41.2%), pulmonary rales in 40 (35.1%), wet rales in 22 (19.3%), fever in 12 (10.5%), and perioral cyanosis in 3 (2.6%).

In terms of treatment measures, as shown in Table 3, of 81 children (41.2%) were treated with methylprednisolone when the disease progressed rapidly and wheezing was obvious. Fifty-two cases (45.6%) were treated with budesonide combined with terbutaline, 44 (38.5%) with budesonide combined with salbutamol, 81 (71.1%) with antibiotics, and 81 (71.1%) with interferon.

According to the results of the first routine blood test combined with CRP within 24 h before and after admission, the children were divided into children with clear bacterial infection ($n = 23$) and those with unknown infection status ($n = 91$). As shown in Table 4, of 23 children with clear bacterial infection were treated with antibiotics, while 58 children with unknown bacterial infection status were treated with antibiotics (63.7%); there was a significant difference between them ($P < 0.05$). Between the two groups, there were no significant differences in other drug treatments, length of hospital stay,

Table 1 Basic information of children with bronchiolitis

Basic information		
Gender, <i>n</i> (%)		
Male		74 (64.9)
Female		40 (35.1)
Onset age, <i>n</i> (%)		
1-6 mo		107 (93.9)
6-12 mo		7 (6.1)
Delivery, <i>n</i> (%)		
Natural labour		71 (62.3)
Cesarean section		43 (37.7)
Time to hospital visit (d), <i>n</i> (%)		
2-4		91 (79.8)
5-10		23 (20.2)
Length of stay (d)		7.1 ± 2.4
Co-infection with bacteria, <i>n</i> (%)		23 (20.2)
Cure rate, <i>n</i> (%)		108 (94.7)

Table 2 Clinical symptoms and imaging characteristics of children with bronchiolitis, *n* (%)

Bronchiolitis		
Wheezing		47 (41.2)
Fever		12 (10.5)
Rhonchi		40 (35.1)
Crackles		22 (19.3)
Cyanosis		3 (2.6)
Increased lung texture		107 (93.8)
Patchy shadow		7 (6.1)

Table 3 Treatments for children with bronchiolitis, *n* (%)

Treatment		
Antibiotic treatment		81 (71.1)
Hormone treatment		47 (41.2)
Antiviral treatment		81 (71.1)
Atomized medication		
Budesonide + terbutaline		52 (45.6)
Budesonide + salbutamol		44 (38.5)

or lung imaging examination results ($P > 0.05$).

According to whether antibiotics were used or not, the children with bronchiolitis without bacterial infection were divided into a antibiotic treatment group ($n = 58$) and a non-antibiotic treatment group ($n = 33$). As shown in Table 5, the average length of hospital stay in the antibiotic treatment group was 7.8 ± 2.5 d, which was significantly longer than that of the non-antibiotic treatment group (5.7 ± 1.8 d; $P < 0.05$). The cure rate was 98.3% in the antibiotic treatment group and 87.9% in the non-antibiotic treatment group ($P < 0.05$).

Table 4 Comparison of bronchiolitis children with bacterial infection and those with unknown bacterial infection

	Bacterial infection (n = 23)	Unknown bacterial infection (n = 91)	t/χ^2	P value
Antibiotic treatment, n (%)	23 (100)	58 (63.7)	11.636	0.000 ¹
Hormone treatment, n (%)	10 (43.5)	35 (38.5)	0.193	0.660
Antiviral treatment, n (%)	17 (73.9)	64 (70.3)	0.115	0.735
Atomized medication, n (%)			1.424	0.491
Budesonide + terbutaline	8 (34.8)	44 (48.4)		
Budesonide + salbutamol	11 (47.8)	33 (36.3)		
Length of stay (d)	7.3 ± 2.9	6.8 ± 2.3	0.959	0.735
Increased lung texture, n (%)	21 (91.3)	86 (94.5)	0.326	0.568
Wheezing, n (%)	11 (47.8)	36 (39.6)	0.518	0.472

¹Significant difference (chi-square test).**Table 5 Effect of antibiotics in children with bronchiolitis without bacterial infection**

	Antibiotic treatment (n = 58)	No antibiotic treatment (n = 33)	t/χ^2	P value
Length of stay (d)	7.8 ± 2.5	5.7 ± 1.8	4.291	0.001 ¹
Cure rate, n (%)	57 (98.3)	29 (87.9)	4.331	0.037 ²

¹Significant difference (t-test).²Significant difference (chi-square test).

DISCUSSION

Acute bronchiolitis is a common cause of infant emergency department attendance and hospitalization[6]. In developed countries, infants under 1 year old with bronchiolitis account for 18% of all pediatric inpatients[7]. In this study, bronchiolitis infants under 1 year old accounted for about 13.5% (132/976) of all pediatric inpatients, which is slightly lower than previous reports. Bronchiolitis is a viral infectious disease, and the virus detection rate in severe bronchiolitis is 100%. Respiratory syncytial virus is the most important pathogenic factor for bronchiolitis[3,4]. More convincingly, routine respiratory virus examinations were not conducted for all children, an aspect that requires improvement. Furthermore, the study observed a higher incidence in male children compared to female, with a male-to-female incidence ratio of approximately 1:0.54. The age of onset was primarily between January and 6 months of age. These findings are not entirely consistent with related reports, which may be attributed to the short duration of the statistical period and the small sample size. Further verification is needed.

Most children hospitalized for acute bronchiolitis have a stable course[8]; however, approximately 2%-6% require admission to a pediatric intensive care unit[9]. The diagnosis of the disease mainly depends on the patient's clinical manifestations, signs, and imaging examination. Clinical manifestations mainly include common symptoms of viral infections, such as sneezing, runny nose, and fever, the gradual development of paroxysmal cough, asthma, *etc.* The chest manifestations were nonspecific, and most of them were accompanied by wheezing and wet rales in the lungs. In this study, in addition to cough, the children mainly presented with wheezing, increased and blurred lung texture on pulmonary imaging, and dry rales and wet rales on clinical auscultation, with a small number of children having fever and perioral cyanosis.

At present, the treatment of this disease is controversial, and there is no clear recommendation to use any drug for the treatment of bronchiolitis[10]. Common medications such as bronchodilators, corticosteroids, and hypertonic saline only provide symptom relief[11]. There is still insufficient evidence for other treatments, such as aerosol inhalation of hypertonic saline, and their role should be further clarified in the future. In this study, 71.7% of the patients were treated with antibiotics, about 41.2% were treated with hormone drugs, and 71.1% were treated with antiviral therapy. The commonly used nebulised drugs were budesonide plus terbutaline (45.6%) and budesonide plus salbutamol (38.5%).

Children with bronchiolitis mostly have a good prognosis. However, some children have persistent problems (*e.g.*, wheezing and cough) after acute infection (> 14 d), increasing the risk of later persistent lung disease and costs to health systems. Active anti-infection treatment could improve the status of more children. In a small randomized trial, azithromycin treatment appeared to reduce the risk of subsequent long-term respiratory disease in 40 hospitalized children with respiratory syncytial virus bronchiolitis during an acute exacerbation[12], which is similar to the findings in this study. In this study, we found that the probability of bacterial infection in patients with capillary bronchitis was low, yet the use of antibiotics was common. Further research revealed that in patients without a clear bacterial infection,

antibiotic use significantly improved the cure rate ($P < 0.05$). However, it also prolonged the duration of hospitalization ($P < 0.05$), highlighting the need for standardized antibiotic use. Moreover, bronchiolitis may lead to lung injury, which can result in recurring wheezing episodes and the development of asthma[13]. It is suggested that severe bronchiolitis during the first episode could be a critical factor or event, and should be considered in designing secondary prevention strategies for asthma. Post-discharge follow-up for up to one year revealed that two cases still experienced symptoms like cough, wheezing, shortness of breath, and other respiratory issues, which were particularly pronounced during respiratory infections or mild exercise.

CONCLUSION

Bronchiolitis is a common acute respiratory infectious disease in infants and young children. It mainly affects male children and the age onset is between 1 to 6 mo. The standard use of antibiotics should be emphasized in view of the prolonged average length of hospital stay between the antibiotic treatment group and the non-antibiotic treatment group. When the course of disease is more than 7 d or the treatment effect is poor, active anti-infective treatment is needed to improve the long-term prognosis. Very few children have recurrent cough and wheezing symptoms within 1 year, which may be related to the risk of later asthma attack.

ARTICLE HIGHLIGHTS

Research background

Bronchiolitis is one of the common lower respiratory tract infections in infants and young children. Severe cases are accompanied by dyspnea and decreased oxygen saturation.

Research motivation

To investigate the clinical characteristics, standardized diagnosis, and treatment of bronchiolitis.

Research objectives

Comparative analysis of the use of antibiotics in bronchiolitis infants with and without bacterial infection has revealed that the use of antibiotics in the treatment of bronchiolitis is controversial. Increasing the use of antibiotics can improve the cure rate of patients, but it will prolong the length of hospital stay. It is necessary to optimize the treatment methods to reduce the length of hospital stay and financial burden of patients.

Research methods

The clinical data of 114 children with newly diagnosed bronchiolitis at the Department of Pediatrics, Tongling Maternal and Child Health Hospital from January 2019 to December 2019 were retrospectively analyzed. The clinical features, imaging features, treatment, and other data were recorded and analyzed.

Research results

The age of onset was mostly from 1 mo to 6 mo, and the time to hospital visit was mostly from the second day to the fourth day of the course of the disease. Lung imaging studies showed increased and blurred lung texture. The main treatment was aerosol inhalation. The average hospitalization time was 7.1 ± 2.4 d, and the total cure rate was 94.7%. In patients without bacterial infection, the use of antibiotics significantly prolonged the length of hospital stay (7.8 ± 2.5 d *vs* 5.7 ± 1.8 d) and improved the cure rate (98.3% *vs* 87.9%, $P < 0.05$).

Research conclusions

Most of the affected infants are male and have a good prognosis. Unnecessary use of antibiotics may significantly prolong the length of hospital stay, thereby putting pressure on both patients and the hospital system.

Research perspectives

Measures should be taken to optimize the treatment methods, and shorten the treatment time and the length of hospital stay, so as to reduce the economic pressure on both patients and on the hospital system.

FOOTNOTES

Author contributions: Shi C and Wu MH contributed to study conception and design, and provision of study materials or patients; Shi C contributed to administrative support; Zuo A, Yang MM, and Jiang RR contributed to data collection and assembly; Shi C contributed to data analysis and interpretation, and manuscript writing; all authors contributed to the final approval of the manuscript.

Institutional review board statement: The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). The study was approved by Institutional Review Board of Tongling Maternal and Child Health Hospital.

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: The authors declare no conflict of interest for this article.

Data sharing statement: All data are provided in this study, and raw data can be obtained from the corresponding author.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Meng-Hua Wu 0009-0008-5196-9491.

S-Editor: Yan JP

L-Editor: Wang TQ

P-Editor: Yan JP

REFERENCES

- 1 Dalziel SR, Haskell L, O'Brien S, Borland ML, Plint AC, Babl FE, Oakley E. Bronchiolitis. *Lancet* 2022; **400**: 392-406 [PMID: 35785792 DOI: 10.1016/S0140-6736(22)01016-9]
- 2 Meissner HC. Viral Bronchiolitis in Children. *N Engl J Med* 2016; **374**: 62-72 [PMID: 26735994 DOI: 10.1056/NEJMra1413456]
- 3 Manti S, Staiano A, Orfeo L, Midulla F, Marseglia GL, Ghizzi C, Zampogna S, Carnielli VP, Favilli S, Ruggieri M, Perri D, Di Mauro G, Gattinara GC, D'Avino A, Becherucci P, Prete A, Zampino G, Lanari M, Biban P, Manzoni P, Esposito S, Corsello G, Baraldi E. UPDATE - 2022 Italian guidelines on the management of bronchiolitis in infants. *Ital J Pediatr* 2023; **49**: 19 [PMID: 36765418 DOI: 10.1186/s13052-022-01392-6]
- 4 Wainwright C. Acute viral bronchiolitis in children- a very common condition with few therapeutic options. *Paediatr Respir Rev* 2010; **11**: 39-45; quiz 45 [PMID: 20113991 DOI: 10.1016/j.prrv.2009.10.001]
- 5 Milési C, Baudin F, Durand P, Emeriaud G, Essouri S, Pouyau R, Baleine J, Beldjilali S, Bordessoule A, Breinig S, Demaret P, Desprez P, Gaillard-Leroux B, Guichoux J, Guilbert AS, Guillot C, Jean S, Levy M, Noizet-Yverneau O, Rambaud J, Recher M, Reynaud S, Valla F, Radoui K, Faure MA, Ferraro G, Mortamet G; French Speaking Group for Pediatric Intensive and Emergency Care. Clinical practice guidelines: management of severe bronchiolitis in infants under 12 months old admitted to a pediatric critical care unit. *Intensive Care Med* 2023; **49**: 5-25 [PMID: 36592200 DOI: 10.1007/s00134-022-06918-4]
- 6 Ghazaly M, Nadel S. Characteristics of children admitted to intensive care with acute bronchiolitis. *Eur J Pediatr* 2018; **177**: 913-920 [PMID: 29654399 DOI: 10.1007/s00431-018-3138-6]
- 7 Hasegawa K, Tsugawa Y, Brown DF, Mansbach JM, Camargo CA Jr. Trends in bronchiolitis hospitalizations in the United States, 2000-2009. *Pediatrics* 2013; **132**: 28-36 [PMID: 23733801 DOI: 10.1542/peds.2012-3877]
- 8 Birkhaug IM, Inchley CS, Aamodt G, Ånestad G, Nystad W, Nakstad B. Infectious burden of respiratory syncytial virus in relation to time of birth modifies the risk of lower respiratory tract infection in infancy: the Norwegian Mother and Child Cohort. *Pediatr Infect Dis J* 2013; **32**: e235-e241 [PMID: 23385949 DOI: 10.1097/INF.0b013e31828ab9ff]
- 9 Ghazaly MMH, Abu Faddan NH, Raafat DM, Mohammed NA, Nadel S. Acute viral bronchiolitis as a cause of pediatric acute respiratory distress syndrome. *Eur J Pediatr* 2021; **180**: 1229-1234 [PMID: 33161501 DOI: 10.1007/s00431-020-03852-9]
- 10 Ralston SL, Lieberthal AS, Meissner HC, Alverson BK, Baley JE, Gadomski AM, Johnson DW, Light MJ, Maraqa NF, Mendonca EA, Phelan KJ, Zorc JJ, Stanko-Lopp D, Brown MA, Nathanson I, Rosenblum E, Sayles S 3rd, Hernandez-Cancio S; American Academy of Pediatrics. Clinical practice guideline: the diagnosis, management, and prevention of bronchiolitis. *Pediatrics* 2014; **134**: e1474-e1502 [PMID: 25349312 DOI: 10.1542/peds.2014-2742]
- 11 Florin TA, Plint AC, Zorc JJ. Viral bronchiolitis. *Lancet* 2017; **389**: 211-224 [PMID: 27549684 DOI: 10.1016/S0140-6736(16)30951-5]
- 12 Stokholm J, Chawes BL, Vissing NH, Bjarnadóttir E, Pedersen TM, Vinding RK, Schoos AM, Wolsk HM, Thorsteinsdóttir S, Hallas HW, Arianto L, Schjørring S, Krogfelt KA, Fischer TK, Pipper CB, Bønnelykke K, Bisgaard H. Azithromycin for episodes with asthma-like symptoms in young children aged 1-3 years: a randomised, double-blind, placebo-controlled trial. *Lancet Respir Med* 2016; **4**: 19-26 [PMID: 26704020 DOI: 10.1016/S2213-2600(15)00500-7]
- 13 Jartti T, Smits HH, Bønnelykke K, Bircan O, Elenius V, Konradsen JR, Maggina P, Makrinioti H, Stokholm J, Hedlin G, Papadopoulos N, Ruszczyński M, Ryczaj K, Schaub B, Schwarze J, Skevaki C, Stenberg-Hammar K, Feleszko W; EAACI Task Force on Clinical Practice Recommendations on Preschool Wheeze. Bronchiolitis needs a revisit: Distinguishing between virus entities and their treatments. *Allergy* 2019; **74**: 40-52 [PMID: 30276826 DOI: 10.1111/all.13624]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

