

90946\_Auto\_Edited.docx

## Early prediction and prevention of infected pancreatic necrosis

Lv C *et al.* Early prediction and prevention of IPN

5 Approximately 20%-30% of patients with acute necrotizing pancreatitis develop infected pancreatic necrosis (IPN), a highly morbid and potentially lethal complication. Early identification of patients at high risk of IPN may facilitate appropriate preventive measures to improve clinical outcomes. In the past two decades, several markers and predictive tools were proposed and evaluated for this purpose. Conventional biomarkers like C-reactive protein, procalcitonin, lymphocyte count, interleukin 6 and 8, and newly developed biomarkers like angiopoietin-2 all showed significant association with IPN. On the other hand, scoring systems like the Acute Physiology and Chronic Health Evaluation II score and Pancreatitis Activity Scoring System score were tested, and they may provide better accuracy. For early prevention of IPN, several new therapies were tested, including early enteral nutrition, antibiotics, probiotics, immune enhancement, *etc.*, and the results varied. Taken together, several evidence-supported predictive markers and scoring systems are readily available for predicting IPN. However, effective treatments to reduce the incidence of IPN are still lacking apart from early enteral nutrition. In this editorial, we summarized evidence concerning early prediction and prevention of IPN, providing insights into future practice and study design. A more homogenous patient population with reliable risk-stratification tools may help find effective treatments to reduce the risk of IPN, thereby achieving individualized treatment.

**Key Words:** Acute pancreatitis; Infected pancreatic necrosis; Biomarker; Scoring system; nutrition therapy; Selective digestive decontamination; Probiotics; Antibiotics; Immune enhancement therapy

Lv C, Zhang ZX, Ke L. Early prediction and prevention of infected pancreatic necrosis. *World J Gastroenterol* 2024; In press

**Core Tip:** Several evidence-supported predictive markers and scoring systems are readily available for predicting infected pancreatic necrosis (IPN). However, effective treatments to reduce the incidence of IPN are still lacking apart from early enteral nutrition. In future research and practice, a more homogenous patient population should be targeted with reliable risk-stratification tools since such a strategy may help find the effective treatment to reduce the risk of IPN, thereby achieving individualized treatment.

2

## **INTRODUCTION**

Acute pancreatitis (AP) is one of the most common gastrointestinal illnesses worldwide<sup>[1]</sup>. The majority of AP cases are mild, self-limited, and are discharged without complications. However, approximately 20% of AP patients develop a complex, prolonged clinical course characterized by pancreatic necrosis, especially when infected pancreatic necrosis (IPN) occurs<sup>[2,3]</sup>. Therefore, it is of clinical value to identify patients at high risk of IPN in the early phase of AP and provide appropriate preventive measures to improve their clinical outcomes.

In the past two decades, several new predictors and predictive tools have been proposed and evaluated, and several new therapies have been tested in trials to prevent IPN. In this editorial, we summarized evidence concerning the early prediction and prevention of IPN (Figure 1), providing insights into future practice and study design.

## **EARLY PREDICTION OF IPN**

### ***Biomarkers***

Many classical biomarkers indicating the inflammation and severity of AP, including C-reactive protein (CRP)<sup>[3,4]</sup>, procalcitonin (PCT)<sup>[5]</sup>, interleukin 6 and 8 (IL-6 and IL-8)<sup>[6,7]</sup>, had showed significant association with IPN in individual studies and meta-analysis<sup>[8]</sup>.

Moreover, in a substudy of the PROPATRIA Trial, Buddingh *et al*<sup>[9]</sup> discovered that early plasma angiopoietin-2 (Ang-2), which plays an important role in the autocrine regulation of vascular stability and permeability, was a better biomarker than conventional predictors such as CRP, PCT, and the imrie score with a cut-off value at 4.51 mg/L.

The relationship between immunosuppression and the development of IPN has also been recognized in the past<sup>[10]</sup>, and biomarkers such as interferon- $\gamma$ <sup>[11]</sup> and monocyte surface expression of HLA-DR antigens<sup>[12,13]</sup> have been tested to reflect the severity of immunosuppression. However, most of these markers are not readily available in hospitals. Through a post-hoc analysis of the TRACE Trial<sup>[14]</sup>, Cai *et al*<sup>[15]</sup> found that absolute lymphocyte count (ALC), a more readily available clinical measure, can predict the occurrence of IPN. Since ALC is a routine laboratory measurement, it might be of wider clinical use in practice.

### Scoring systems

Several clinical scoring systems have been shown to predict IPN with adequate accuracy. The first is the <sup>1</sup> Acute Physiology and Chronic Health Evaluation (APACHE) II score. An APACHE II score of more than 8 at admission was found to be a risk factor for IPN in patients with severe AP (SAP)<sup>[8,16]</sup>. Systemic Inflammatory Response Syndrome (SIRS) score was another option since persistent inflammation is involved in the development of IPN<sup>[16,17]</sup>. An observational study showed that longer SIRS duration was significantly associated with a higher incidence of IPN<sup>[18]</sup>.

The Pancreatitis Activity Scoring System (PASS), which is an AP-specific score to reflect the disease severity, was tested in the prediction of IPN. In a retrospective study conducted by Ke *et al*<sup>[19]</sup>, the predictive accuracy of the PASS score at admission was better than the APACHE II score in predicting IPN. However, considering the dominating weight assigned to opioid usage in the PASS, Paragomi *et al*<sup>[20]</sup> modified the original PASS score by removing or partly reducing the weight of opioid usage (mPASS <sup>4</sup> 1-4). The mPASS could predict SAP with reasonable accuracy and differentiate between

patients with different early trajectories in patients with different severities. For the prediction of IPN, Mao *et al*<sup>[21]</sup> found that the mPASS-4 model outperformed the conventional indices in predicting IPN, thereby increasing the likelihood of clinical usage.

## **EARLY PREVENTION OF IPN**

In the past few decades, multiple attempts have been made to reduce the incidence of IPN, including nutrition therapy, selective digestive decontamination (SDD), antibiotic therapy, and immune enhancement. Unfortunately, most of the studies have not come to a positive conclusion.

### ***Nutrition therapy***

Two different RCTs<sup>[22,23]</sup> conducted in patients with SAP demonstrated that early total enteral nutrition, compared with total parenteral nutrition, could reduce the incidence of IPN, thereby reducing organ failure and mortality. On the one hand, the lack of enteral feeding results in atrophy of the gastrointestinal mucosa, bacterial overgrowth, and increased intestinal permeability<sup>[24]</sup>. On the other hand, parenteral nutrition (enteric starvation) was associated with rapid and severe atrophy of lymphoid tissue associated with the gut<sup>[25-27]</sup>. As a result, early enteral nutrition may alleviate the translocation of bacteria or bacterial products into the circulation<sup>[28-31]</sup>.

### ***SDD***

In the 1980s, investigations into the source of infection in SAP patients found that Gram-negative aerobic bacteria, originating from the digestive tract are predominantly isolated from IPN samples<sup>[32,33]</sup>. Accordingly, SDD gained broad interest among the research community. The results of a more recent randomized trial<sup>[34]</sup> with a relatively small sample size of 102 SAP patients showed that SDD could significantly reduce the incidence of IPN, which was associated with improved morbidity and mortality.

However, these results have not been confirmed by a large, multicenter trial, and therefore, SDD has not become a standard of care in current guidelines<sup>[35]</sup>.

### **Probiotics**

In experimental and small clinical studies, <sup>1</sup> certain strains of probiotic bacteria might prevent infectious complications by reducing small-bowel bacterial overgrowth, restoring gastrointestinal barrier function, and modulating the immune system<sup>[36,37]</sup>. To confirm the clinical significance of probiotics in SAP patients, Besselink *et al*<sup>[38]</sup> conducted a large, randomized, double-blind, controlled trial testing the effect of probiotic therapy on the incidence of infectious complications. Unfortunately, the results <sup>1</sup> showed no beneficial effect of probiotic prophylaxis on multiple infectious complications. On the contrary, mortality in the probiotics group was about twice as high as in the placebo group, which might be attributed to increased incidence of bowel ischemia. The administration of probiotic bacteria daily as an adjunct to <sup>8</sup> enteral nutrition might increase local oxygen demand, with a combined deleterious effect on the <sup>1</sup> already compromised blood flow. Another possible explanation could be that the presence of probiotics caused local inflammation at the mucosal level. Experimental studies have shown that gut epithelial cells under metabolic stress react to commensal bacteria with an inflammatory response<sup>[39]</sup>. Recently, in addition to gram-negative bacteria, gram-positive bacteria as well as yeasts related infections were observed with increasing incidence<sup>[40-42]</sup>. Therefore, research interests in the source of bacteria in IPN patients have been raised again, and the corresponding preventive measures need to be further studied.

### **Antibiotics**

Systemic antibiotic prophylaxis has long been considered effective in preventing secondary infection in AP<sup>[43]</sup>. The results from a randomized controlled trial testing prophylactic meropenem suggest that although early antibiotic treatment might reduce the occurrence of septic complications and improve the prognosis of AP, it does not



prevent the occurrence of IPN<sup>[44]</sup>. However, another randomized, double-blind trial conducted in patients with sterile necrotizing pancreatitis demonstrated similar rates of infection, operation, and death between the treatment groups receiving meropenem or placebo<sup>[45]</sup>. In addition, imipenem-cilastatin was also tested in patients with ANP. However, it did not reduce the incidence of IPN and increased the risk of fungal infections<sup>[46]</sup>.

In summary, current evidence does not support the use of prophylactic antibiotics in patients with necrotizing pancreatitis since it is ineffective in reducing IPN and may be associated with potential risks.

### *Immune enhancement therapy*

Given that there is evidence of immunosuppression in the early phase of SAP and its association with infectious complications<sup>[10,47-49]</sup>. The Chinese Acute Pancreatitis Clinical Trials Group (CAPCTG) conducted serial trials to test the effects of immune enhancement by subcutaneous injection of thymosin alpha 1 (Tα1) on the incidence of IPN<sup>[14]</sup>. In the pilot trial, it was found that the 28-d positive blood culture rate was almost half in the thymosin α1 group than the control group (16.6% vs 41.7%,  $P = 0.012$ ), and the rate of IPN decreased from 29.4% to 8.3% ( $P = 0.036$ ) after the treatment of thymosin α1. However, the phase-III confirmatory trial found that the immune-enhancing Tα1 treatment did not significantly reduce the incidence of IPN compared with placebo in patients with predicted severe ANP. This was followed by a post-hoc analysis of the trial<sup>[50]</sup>, which found that patients with predicted severe ANP and no lymphopenia (baseline ALC  $\geq 0.8 \times 10^9/L$ ) may benefit from Tα1. However, due to the post-hoc design, new trials are needed to confirm the findings before any formal recommendation can be made.

### **CONCLUSION**

In conclusion, several evidence-supported predictive markers and scoring systems are readily available for predicting IPN. However, effective treatments to reduce the

incidence of IPN are still lacking apart from early enteral nutrition. In future research, a more homogenous group of patients should be selected with reliable risk-stratification tools since such a strategy may help find the effective treatment to reduce the risk of IPN, thereby achieving individualized treatment.

## REFERENCES

- 1 **Xiao AY**, Tan ML, Wu LM, Asrani VM, Windsor JA, Yadav D, Petrov MS. Global incidence and mortality of pancreatic diseases: a systematic review, meta-analysis, and meta-regression of population-based cohort studies. *Lancet Gastroenterol Hepatol* 2016; **1**: 45-55 [PMID: 28404111 DOI: 10.1016/S2468-1253(16)30004-8]
- 2 **Forsmark CE**, Vege SS, Wilcox CM. Acute Pancreatitis. *N Engl J Med* 2016; **375**: 1972-1981 [PMID: 27959604 DOI: 10.1056/NEJMra1505202]
- 3 **Banks PA**, Bollen TL, Dervenis C, Gooszen HG, Johnson CD, Sarr MG, Tsiotos GG, Vege SS; Acute Pancreatitis Classification Working Group. Classification of acute pancreatitis--2012: revision of the Atlanta classification and definitions by international consensus. *Gut* 2013; **62**: 102-111 [PMID: 23100216 DOI: 10.1136/gutjnl-2012-302779]
- 4 **Gomatos IP**, Xiaodong X, Ghaneh P, Halloran C, Raraty M, Lane B, Sutton R, Neoptolemos JP. Prognostic markers in acute pancreatitis. *Expert Rev Mol Diagn* 2014; **14**: 333-346 [PMID: 24649820 DOI: 10.1586/14737159.2014.897608]
- 5 **Rau BM**, Kemppainen EA, Gumbs AA, Büchler MW, Wegscheider K, Bassi C, Puolakkainen PA, Beger HG. Early assessment of pancreatic infections and overall prognosis in severe acute pancreatitis by procalcitonin (PCT): a prospective international multicenter study. *Ann Surg* 2007; **245**: 745-754 [PMID: 17457167 DOI: 10.1097/01.sla.0000252443.22360.46]
- 6 **Rau B**, Steinbach G, Gansauge F, Mayer JM, Grünert A, Beger HG. The potential role of procalcitonin and interleukin 8 in the prediction of infected necrosis in acute pancreatitis. *Gut* 1997; **41**: 832-840 [PMID: 9462219 DOI: 10.1136/gut.41.6.832]



- 7 **Ji L**, Lv JC, Song ZF, Jiang MT, Li L, Sun B. Risk factors of infected pancreatic necrosis secondary to severe acute pancreatitis. *Hepatobiliary Pancreat Dis Int* 2016; **15**: 428-433 [PMID: 27498584 DOI: 10.1016/s1499-3872(15)60043-1]
- 8 **Li W**, Ou L, Fu Y, Chen Y, Yin Q, Song H. Risk factors for concomitant infectious pancreatic necrosis in patients with severe acute pancreatitis: A systematic review and meta-analysis. *Clin Res Hepatol Gastroenterol* 2022; **46**: 101901 [PMID: 35304319 DOI: 10.1016/j.clinre.2022.101901]
- 9 **Buddingh KT**, Koudstaal LG, van Santvoort HC, Besselink MG, Timmer R, Rosman C, van Goor H, Nijmeijer RM, Gooszen H, Leuvenink HG, Ploeg RJ, Nieuwenhuijs VB. Early angiopoietin-2 levels after onset predict the advent of severe pancreatitis, multiple organ failure, and infectious complications in patients with acute pancreatitis. *J Am Coll Surg* 2014; **218**: 26-32 [PMID: 24355874 DOI: 10.1016/j.jamcollsurg.2013.09.021]
- 10 **Ueda T**, Takeyama Y, Yasuda T, Shinzeki M, Sawa H, Nakajima T, Ajiki T, Fujino Y, Suzuki Y, Kuroda Y. Immunosuppression in patients with severe acute pancreatitis. *J Gastroenterol* 2006; **41**: 779-784 [PMID: 16988767 DOI: 10.1007/s00535-006-1852-8]
- 11 **Gardiner BJ**, Lee SJ, Cristiano Y, Levvey BJ, Sullivan LC, Snell GI, Peleg AY, Westall GP. Evaluation of Quantiferon®-Monitor as a biomarker of immunosuppression and predictor of infection in lung transplant recipients. *Transpl Infect Dis* 2021; **23**: e13550 [PMID: 33351991 DOI: 10.1111/tid.13550]
- 12 **Minkov G**, Dimitrov E, Yovtchev Y, Enchev E, Lokova R, Halacheva K. Prognostic value of peripheral blood CD14+HLA-DR+ monocytes in patients with acute pancreatitis. *J Immunoassay Immunochem* 2021; **42**: 478-492 [PMID: 33818295 DOI: 10.1080/15321819.2021.1903491]
- 13 **Chéron A**, Monneret G, Landelle C, Floccard B, Allaouchiche B. [Low monocytic HLA-DR expression and risk of secondary infection]. *Ann Fr Anesth Reanim* 2010; **29**: 368-376 [PMID: 20356708 DOI: 10.1016/j.annfar.2010.02.015]
- 14 **Ke L**, Zhou J, Mao W, Chen T, Zhu Y, Pan X, Mei H, Singh V, Buxbaum J, Doig G, He C, Gu W, Lu W, Tu S, Ni H, Zhang G, Zhao X, Sun J, Chen W, Song J, Shao M, Tu J, Xia L, He W, Zhu Q, Li K, Yao H, Wu J, Fu L, Jiang W, Zhang H, Lin J, Li B, Tong Z,

- Windsor J, Liu Y, Li W; Chinese Acute Pancreatitis Clinical Trials Group (CAPCTG). Immune enhancement in patients with predicted severe acute necrotising pancreatitis: a multicentre double-blind randomised controlled trial. *Intensive Care Med* 2022; **48**: 899-909 [PMID: 35713670 DOI: 10.1007/s00134-022-06745-7]
- 15 **Cai T**, Mao W, Liu M, Zhou J, Wang X, Zhou J, Liu Y, Lv G, Ke L, Zhang Y. Early mean absolute lymphocyte count in acute necrotizing pancreatitis is associated with infected pancreatic necrosis. *Int Immunopharmacol* 2023; **117**: 109883 [PMID: 36827921 DOI: 10.1016/j.intimp.2023.109883]
- 16 **Pando E**, Alberti P, Hidalgo J, Vidal L, Dopazo C, Caralt M, Blanco L, Gómez-Gavara C, Bilbao I, Balsells J, Charco R. The role of extra-pancreatic infections in the prediction of severity and local complications in acute pancreatitis. *Pancreatology* 2018; **18**: 486-493 [PMID: 29802078 DOI: 10.1016/j.pan.2018.05.481]
- 17 **Zerem E**. Treatment of severe acute pancreatitis and its complications. *World J Gastroenterol* 2014; **20**: 13879-13892 [PMID: 25320523 DOI: 10.3748/wjg.v20.i38.13879]
- 18 **Tan C**, Yang L, Shi F, Hu J, Zhang X, Wang Y, Deng Z, Li J, Yuan H, Shi T, Li C, Xiao Y, Peng Y, Xu W, Huang Y. Early Systemic Inflammatory Response Syndrome Duration Predicts Infected Pancreatic Necrosis. *J Gastrointest Surg* 2020; **24**: 590-597 [PMID: 30891659 DOI: 10.1007/s11605-019-04149-5]
- 19 **Ke L**, Mao W, Li X, Zhou J, Li G, Ye B, Tong Z, Li W. The Pancreatitis Activity Scoring System in Predicting Infection of Pancreatic Necrosis. *Am J Gastroenterol* 2018; **113**: 1393-1394 [PMID: 29880970 DOI: 10.1038/s41395-018-0112-x]
- 20 **Paragomi P**, Hinton A, Pothoulakis I, Talukdar R, Kochhar R, Goenka MK, Gulla A, Gonzalez JA, Singh VK, Bogado MF, Stevens T, Barbu ST, Nawaz H, Gutierrez SC, Zarnescu N, Archibugi L, Easler JJ, Triantafyllou K, Peláez-Luna M, Thakkar S, Ocampo C, Enrique de-Madaria, Cote GA, Lee PJ, Krishna S, Lara LF, Han S, Wu BU, Papachristou GI. The Modified Pancreatitis Activity Scoring System Shows Distinct Trajectories in Acute Pancreatitis: An International Study. *Clin Gastroenterol Hepatol* 2022; **20**: 1334-1342.e4 [PMID: 34543736 DOI: 10.1016/j.cgh.2021.09.014]

- 21 **Mao W**, Li K, Zhou J, Chen M, Ye B, Li G, Singh V, Buxbaum J, Fu X, Tong Z, Liu Y, Windsor J, Li W, Ke L; Chinese Acute Pancreatitis Clinical Trials Group (CAPCTG). Prediction of infected pancreatic necrosis in acute necrotizing pancreatitis by the modified pancreatitis activity scoring system. *United European Gastroenterol J* 2023; **11**: 69-78 [PMID: 36579414 DOI: 10.1002/ueg2.12353]
- 22 **Petrov MS**, Kukosh MV, Emelyanov NV. A randomized controlled trial of enteral versus parenteral feeding in patients with predicted severe acute pancreatitis shows a significant reduction in mortality and in infected pancreatic complications with total enteral nutrition. *Dig Surg* 2006; **23**: 336-44; discussion 344-5 [PMID: 17164546 DOI: 10.1159/000097949]
- 23 **Wu XM**, Ji KQ, Wang HY, Li GF, Zang B, Chen WM. Total enteral nutrition in prevention of pancreatic necrotic infection in severe acute pancreatitis. *Pancreas* 2010; **39**: 248-251 [PMID: 19910834 DOI: 10.1097/MPA.0b013e3181bd6370]
- 24 **Li P**, Jian JN, Chen RL. Effect of Early Enteral Nutrition on Serum Inflammatory Factors and Intestinal Mucosal Permeability in Patients with Severe Acute Pancreatitis. *Turk J Gastroenterol* 2021; **32**: 907-912 [PMID: 34787096 DOI: 10.5152/tjg.2021.201033]
- 25 **King BK**, Li J, Kudsk KA. A temporal study of TPN-induced changes in gut-associated lymphoid tissue and mucosal immunity. *Arch Surg* 1997; **132**: 1303-1309 [PMID: 9403534 DOI: 10.1001/archsurg.1997.01430360049009]
- 26 **Janu P**, Li J, Renegar KB, Kudsk KA. Recovery of gut-associated lymphoid tissue and upper respiratory tract immunity after parenteral nutrition. *Ann Surg* 1997; **225**: 707-15; discussion 715-7 [PMID: 9230811 DOI: 10.1097/00000658-199706000-00008]
- 27 **Kudsk KA**, Li J, Renegar KB. Loss of upper respiratory tract immunity with parenteral feeding. *Ann Surg* 1996; **223**: 629-35; discussion 635-8 [PMID: 8645036 DOI: 10.1097/00000658-199606000-00001]
- 28 **Hadfield RJ**, Sinclair DG, Houldsworth PE, Evans TW. Effects of enteral and parenteral nutrition on gut mucosal permeability in the critically ill. *Am J Respir Crit Care Med* 1995; **152**: 1545-1548 [PMID: 7582291 DOI: 10.1164/ajrccm.152.5.7582291]

- 29 **Nakasaki H**, Mitomi T, Tajima T, Ohnishi N, Fujii K. Gut bacterial translocation during total parenteral nutrition in experimental rats and its countermeasure. *Am J Surg* 1998; **175**: 38-43 [PMID: 9445237 DOI: 10.1016/S0002-9610(97)00231-6]
- 30 **Shou J**, Lappin J, Minnard EA, Daly JM. Total parenteral nutrition, bacterial translocation, and host immune function. *Am J Surg* 1994; **167**: 145-150 [PMID: 8311126 DOI: 10.1016/0002-9610(94)90065-5]
- 31 **Qiu JG**, Delany HM, Teh EL, Freundlich L, Gliedman ML, Steinberg JJ, Chang CJ, Levenson SM. Contrasting effects of identical nutrients given parenterally or enterally after 70% hepatectomy: bacterial translocation. *Nutrition* 1997; **13**: 431-437 [PMID: 9225335 DOI: 10.1016/S0899-9007(97)91281-8]
- 32 **Beger HG**, Bittner R, Block S, Büchler M. Bacterial contamination of pancreatic necrosis. A prospective clinical study. *Gastroenterology* 1986; **91**: 433-438 [PMID: 3522342 DOI: 10.1016/0016-5085(86)90579-2]
- 33 **Runkel NS**, Moody FG, Smith GS, Rodriguez LF, LaRocco MT, Miller TA. The role of the gut in the development of sepsis in acute pancreatitis. *J Surg Res* 1991; **51**: 18-23 [PMID: 2067354 DOI: 10.1016/0022-4804(91)90064-S]
- 34 **Luiten EJ**, Hop WC, Lange JF, Bruining HA. Controlled clinical trial of selective decontamination for the treatment of severe acute pancreatitis. *Ann Surg* 1995; **222**: 57-65 [PMID: 7618970 DOI: 10.1097/00000658-199507000-00010]
- 35 **Baron TH**, DiMaio CJ, Wang AY, Morgan KA. American Gastroenterological Association Clinical Practice Update: Management of Pancreatic Necrosis. *Gastroenterology* 2020; **158**: 67-75.e1 [PMID: 31479658 DOI: 10.1053/j.gastro.2019.07.064]
- 36 **Bengmark S**. Ecological control of the gastrointestinal tract. The role of probiotic flora. *Gut* 1998; **42**: 2-7 [PMID: 9505873 DOI: 10.1136/gut.42.1.2]
- 37 **Guarner F**, Malagelada JR. Gut flora in health and disease. *Lancet* 2003; **361**: 512-519 [PMID: 12583961 DOI: 10.1016/S0140-6736(03)12489-0]
- 38 **Besselink MG**, van Santvoort HC, Buskens E, Boermeester MA, van Goor H, Timmerman HM, Nieuwenhuijs VB, Bollen TL, van Ramshorst B, Witterman BJ, Rosman C, Ploeg RJ, Brink MA, Schaapherder AF, Dejong CH, Wahab PJ, van Laarhoven CJ, van

der Harst E, van Eijck CH, Cuesta MA, Akkermans LM, Gooszen HG; Dutch Acute Pancreatitis Study Group. Probiotic prophylaxis in predicted severe acute pancreatitis: a randomised, double-blind, placebo-controlled trial. *Lancet* 2008; **371**: 651-659 [PMID: 18279948 DOI: 10.1016/S0140-6736(08)60207-X]

39 **Okumura R**, Takeda K. Roles of intestinal epithelial cells in the maintenance of gut homeostasis. *Exp Mol Med* 2017; **49**: e338 [PMID: 28546564 DOI: 10.1038/emm.2017.20]

40 **Iseemann R**, Schwarz M, Rau B, Trautmann M, Schober W, Beger HG. Characteristics of infection with *Candida* species in patients with necrotizing pancreatitis. *World J Surg* 2002; **26**: 372-376 [PMID: 11865377 DOI: 10.1007/s00268-001-0146-9]

41 **Grewe M**, Tsiotos GG, Luque de-Leon E, Sarr MG. Fungal infection in acute necrotizing pancreatitis. *J Am Coll Surg* 1999; **188**: 408-414 [PMID: 10195725 DOI: 10.1016/S1072-7515(98)00334-2]

42 **Hoerauf A**, Hammer S, Müller-Myhsok B, Rupprecht H. Intra-abdominal *Candida* infection during acute necrotizing pancreatitis has a high prevalence and is associated with increased mortality. *Crit Care Med* 1998; **26**: 2010-2015 [PMID: 9875913 DOI: 10.1097/00003246-199812000-00031]

43 **Working Party of the British Society of Gastroenterology**; Association of Surgeons of Great Britain and Ireland; Pancreatic Society of Great Britain and Ireland; Association of Upper GI Surgeons of Great Britain and Ireland. UK guidelines for the management of acute pancreatitis. *Gut* 2005; **54** Suppl 3: iii1-iii9 [PMID: 15831893 DOI: 10.1136/gut.2004.057026]

44 **Manes G**, Uomo I, Menchise A, Rabitti PG, Ferrara EC, Uomo G. Timing of antibiotic prophylaxis in acute pancreatitis: a controlled randomized study with meropenem. *Am J Gastroenterol* 2006; **101**: 1348-1353 [PMID: 16771960 DOI: 10.1111/j.1572-0241.2006.00567.x]

45 **Dellinger EP**, Tellado JM, Soto NE, Ashley SW, Barie PS, Dugernier T, Imrie CW, Johnson CD, Knaebel HP, Laterre PF, Maravi-Poma E, Kissler JJ, Sanchez-Garcia M, Utzolino S. Early antibiotic treatment for severe acute necrotizing pancreatitis: a

randomized, double-blind, placebo-controlled study. *Ann Surg* 2007; **245**: 674-683 [PMID: 17457158 DOI: 10.1097/01.sla.0000250414.09255.84]

46 **Maraví-Poma E**, Gener J, Alvarez-Lerma F, Olaechea P, Blanco A, Domínguez-Muñoz JE; Spanish Group for the Study of Septic Complications in Severe Acute Pancreatitis. Early antibiotic treatment (prophylaxis) of septic complications in severe acute necrotizing pancreatitis: a prospective, randomized, multicenter study comparing two regimens with imipenem-cilastatin. *Intensive Care Med* 2003; **29**: 1974-1980 [PMID: 14551680 DOI: 10.1007/s00134-003-1956-z]

47 **Yu WK**, Li WQ, Li N, Li JS. Mononuclear histocompatibility leukocyte antigen-DR expression in the early phase of acute pancreatitis. *Pancreatology* 2004; **4**: 233-243 [PMID: 15166475 DOI: 10.1159/000078748]

48 **Pan T**, Zhou T, Li L, Liu Z, Chen Y, Mao E, Li M, Qu H, Liu J. Monocyte programmed death ligand-1 expression is an early marker for predicting infectious complications in acute pancreatitis. *Crit Care* 2017; **21**: 186 [PMID: 28705256 DOI: 10.1186/s13054-017-1781-3]

49 **Li J**, Yang WJ, Huang LM, Tang CW. Immunomodulatory therapies for acute pancreatitis. *World J Gastroenterol* 2014; **20**: 16935-16947 [PMID: 25493006 DOI: 10.3748/wjg.v20.i45.16935]

50 **Ke L**, Mao W, Shao F, Zhou J, Xu M, Chen T, Liu Y, Tong Z, Windsor J, Ma P, Li W; Chinese Acute Pancreatitis Clinical Trials Group (CAPCTG). Association between pretreatment lymphocyte count and efficacy of immune-enhancing therapy in acute necrotising pancreatitis: a post-hoc analysis of the multicentre, randomised, placebo-controlled TRACE trial. *EClinicalMedicine* 2023; **58**: 101915 [PMID: 37007743 DOI: 10.1016/j.eclinm.2023.101915]



**Figure 1 Summary of evidence concerning the early prediction and prevention of infected pancreatic necrosis.** CRP: C-reactive protein; PCT: Procalcitonin; IL-6 and 8: Interleukin 6 and 8; Ang-2: Angiopoietin-2; ALC: Absolute lymphocyte count; APACHE II: Acute Physiology and Chronic Health Evaluation II; SIRS: systemic inflammatory response syndrome; PASS: Pancreatitis Activity Scoring System; mPASS: Modified Pancreatitis Activity Scoring System; EN: Enteral nutrition; PN: Parenteral nutrition; Tα1: Thymosin alpha 1.

18%

SIMILARITY INDEX

---

PRIMARY SOURCES

---

- |                                                                                                                                                                            |                                                                                                                                                                                                                                                                                                             |               |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| <div style="background-color: red; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">1</div>     | <a href="http://www.natap.org" style="color: red;">www.natap.org</a><br><small>Internet</small>                                                                                                                                                                                                             | 95 words — 5% |
| <hr/>                                                                                                                                                                      |                                                                                                                                                                                                                                                                                                             |               |
| <div style="background-color: magenta; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">2</div> | <a href="#">Na Yang, Lu Ke, Zhihui Tong, Weiqin Li. "The effect of thymosin α1 for prevention of infection in patients with severe acute pancreatitis", Expert Opinion on Biological Therapy, 2018</a><br><small>Crossref</small>                                                                           | 59 words — 3% |
| <hr/>                                                                                                                                                                      |                                                                                                                                                                                                                                                                                                             |               |
| <div style="background-color: purple; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">3</div>  | <a href="http://www.pubmedcentral.nih.gov" style="color: purple;">www.pubmedcentral.nih.gov</a><br><small>Internet</small>                                                                                                                                                                                  | 47 words — 3% |
| <hr/>                                                                                                                                                                      |                                                                                                                                                                                                                                                                                                             |               |
| <div style="background-color: teal; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">4</div>    | <a href="#">Wenjian Mao, Kang Li, Jing Zhou, Miao Chen et al. "Prediction of infected pancreatic necrosis in acute necrotizing pancreatitis by the modified pancreatitis activity scoring system", United European Gastroenterology Journal, 2022</a><br><small>Crossref</small>                            | 33 words — 2% |
| <hr/>                                                                                                                                                                      |                                                                                                                                                                                                                                                                                                             |               |
| <div style="background-color: green; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">5</div>   | <a href="http://rcastoragev2.blob.core.windows.net" style="color: green;">rcastoragev2.blob.core.windows.net</a><br><small>Internet</small>                                                                                                                                                                 | 33 words — 2% |
| <hr/>                                                                                                                                                                      |                                                                                                                                                                                                                                                                                                             |               |
| <div style="background-color: brown; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">6</div>   | <a href="#">Lu Ke, Wenjian Mao, Fang Shao, Jing Zhou et al. "Association between pretreatment lymphocyte count and efficacy of immune-enhancing therapy in acute necrotising pancreatitis: a post-hoc analysis of the multicentre, randomised, placebo-controlled TRACE trial", eClinicalMedicine, 2023</a> | 22 words — 1% |

---

7	link.springer.com	16 words — 1%
	Internet	

---

8	www.hindawi.com	15 words — 1%
	Internet	

---

EXCLUDE QUOTES	ON	EXCLUDE SOURCES	< 15 WORDS
EXCLUDE BIBLIOGRAPHY	ON	EXCLUDE MATCHES	< 10 WORDS