

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 26243

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Genetic polymorphism in CD14 gene, a co-receptor of TLR4 is associated with nonalcoholic fatty liver disease

Kapil S *et al.* CD14 polymorphism in NAFLD

Shweta Kapil, Ajay Duseja, BalKrishan Sharma, Bhupesh Singla, Anuradha Chakraborti, Ashim Das, Pallab Ray, Radha K Dhiman, Yogesh Chawla

Shweta Kapil, Ajay Duseja, BalKrishan Sharma, Bhupesh Singla, Radha K Dhiman, Yogesh Chawla, Departments of Hepatology, Postgraduate Institute of Medical Education and Research, Chandigarh, India

Match Overview

1	Internet 122 words crawled on 01-Jul-2015 www.wjgnet.com	3%
2	Internet 117 words crawled on 31-Jul-2016 onlinelibrary.wiley.com	2%
3	Internet 74 words crawled on 07-Nov-2008 www.clinchem.org	2%
4	CrossCheck 46 words Naik, Adviti, Rok Košir, and Damjana Rozman. "Genomic aspects of NAFLD pathogenesis", <i>Genomics</i> , 2013.	1%
5	Internet 44 words crawled on 31-Jul-2016 ejnmires.springeropen.com	1%
6	CrossCheck 20 words "Liver", <i>The American Journal of Gastroenterology</i> , 2015.	<1%

找到约 6,780 条结果 (用时 1.16 秒)

Google 学术: Genetic polymorphism in CD14 gene, a co-receptor of TLR4 is associated with nonalcoholic fatty liver disease

The genetics of nonalcoholic fatty liver disease - Osterreicher - 被引用次数: 22

Toll-like receptor 4 signaling in liver injury and hepatic ... - Guo - 被引用次数: 106

Toll-like receptor signaling in the liver - Schwabe - 被引用次数: 274

Genetic Predisposition in NAFLD and NASH: Impact on Severity of ...

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: P Dongiovanni - 2013 - 被引用次数: 64 - [相关文章](#)

Association studies confirm that the I148M polymorphism is also a strong modifier of Heritable components of non-alcoholic fatty liver disease (NAFLD) and liver indices ... These were localized in or near the genes neurocan (NCAN, SNP affect the strength of interactions with either agonists and/or co-receptors [191].

Toll-like receptor 4 signaling in liver injury and hepatic fibrogenesis

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: J Guo - 2010 - 被引用次数: 106 - [相关文章](#)

2010年10月21日 - In humans, single nucleotide polymorphisms of the TLR4 gene have an effect on ... including viral hepatitis, alcoholic and non-alcoholic liver diseases, autoimmune ... TLR4 and the two co-receptors CD14 and myeloid differentiation protein TLR4 is involved in the generation of steatosis, congestion and ...

Genetic modifiers of non-alcoholic fatty liver disease progression

www.sciencedirect.com/science/article/pii/S0925443911001700 ▾ [翻译此页](#)

作者: QM Anstee - 2011 - 被引用次数: 40 - [相关文章](#)

Non-alcoholic fatty liver disease (NAFLD) is now recognised as the most ... GWAS and candidate gene studies have identified several genetic modifiers. found no association with polymorphisms in either TLR4 or the NOD2 receptor for ... Carriage of the CD14 minor allele is associated with increased expression of both ...

The genetics of nonalcoholic fatty liver disease (PDF Download ...

全部 图片 新闻 视频 购物 更多 ▾ 搜索工具

找到约 7,890 条结果 (用时 1.00 秒)

Google 学术 : Genetic polymorphism in CD14 gene, a co-receptor of TLR4 is associated with nonalcoholic fatty liver disease

The genetics of nonalcoholic fatty liver disease - Osterreicher - 被引用次数 : 22

Toll-like receptor 4 signaling in liver injury and hepatic ... - Guo - 被引用次数 : 112

Toll-like receptor signaling in the liver - Schwabe - 被引用次数 : 279

CD14 CD14 molecule [(human)] - NCBI

www.ncbi.nlm.nih.gov/gene/929 ▾ 翻译此页

2016年9月4日 - Association between CD14 gene promoter polymorphisms with serum total-IgE ... Genetic variants of NOS3, CD14, MMP3 and IL4R are implicated in the ... liver inflammation grade in patients with nonalcoholic fatty liver disease. does not require the CD14 co-receptor to elicit TLR4-mediated responses ...

Toll-like receptor 4 signaling in liver injury and hepatic fibrogenesis

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者 : J Guo - 2010 - 被引用次数 : 112 - 相关文章

2010年10月21日 - In humans, single nucleotide polymorphisms of the TLR4 gene have an effect on ... including viral hepatitis, alcoholic and non-alcoholic liver diseases, autoimmune ... TLR4 and the two co-receptors CD14 and myeloid differentiation protein TLR4 is involved in the generation of steatosis, congestion and ...

Genetic modifiers of non-alcoholic fatty liver disease progression

www.sciencedirect.com/science/article/pii/S0925443911001700 ▾ 翻译此页

作者 : QM Anstee - 2011 - 被引用次数 : 39 - 相关文章

Non-alcoholic fatty liver disease (NAFLD) is now recognised as the most ... GWAS and candidate gene studies have identified several genetic modifiers, found no association with polymorphisms in either TLR4 or the NOD2 receptor for ... Carriage of the CD14 minor allele is associated with increased expression of both ...

The genetics of nonalcoholic fatty liver disease (PDF Download ...