

World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2020 April 27; 12(4): 129-196





REVIEW

- 129 Management of complicated acute appendicitis in children: Still an existing controversy
Zavras N, Vaos G

ORIGINAL ARTICLE

Retrospective Cohort Study

- 138 Newly developed self-expandable Niti-S MD colonic metal stent for malignant colonic obstruction
Miyasako Y, Kuwai T, Ishaq S, Tao K, Konishi H, Miura R, Sumida Y, Kuroki K, Tamaru Y, Kusunoki R, Yamaguchi A, Kouno H, Kohno H

- 149 Mammalian target of rapamycin inhibitors after post-transplant hepatocellular carcinoma recurrence: Is it too late?
Au KP, Chok KSH

Retrospective Study

- 159 Contemporary indications for and outcomes of hepatic resection for neuroendocrine liver metastases
Scoville SD, Xourafas D, Ejaz AM, Tsung A, Pawlik T, Cloyd JM

- 171 Clinical course of suspected small gastrointestinal stromal tumors in the stomach
Ye LS, Li Y, Liu W, Yao MH, Khan N, Hu B

Observational Study

- 178 Short term outcomes of minimally invasive selective lateral pelvic lymph node dissection for low rectal cancer
Wong KY, Tan AM

- 190 Feasibility of robotic assisted bladder sparing pelvic exenteration for locally advanced rectal cancer: A single institution case series
Heah NH, Wong KY

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Krasimir Dimitrov Ivanov, DSc, MD, PhD, Professor, Department of General and Operative Surgery, University Hospital "St. Marina" 1, Varna 9000, Bulgaria

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (WJGS, *World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagoplasty, esophagostomy, fundoplication, gastrectomy, gastroenterostomy, gastropexy, hepatectomy, jejunioileal bypass, liver transplantation, pancreas transplantation, pancreatectomy, pancreaticoduodenectomy, and pancreaticojejunostomy, etc.

INDEXING/ABSTRACTING

The WJGS is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (CSTJ), and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: Yan-Xia Xing

Proofing Production Department Director: Xiang Li

Responsible Editorial Office Director: Ruo-Yu Ma

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Varut Lohsiriwat, Shu-You Peng

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

April 27, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/gerinfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/gerinfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/gerinfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Contemporary indications for and outcomes of hepatic resection for neuroendocrine liver metastases

Steven D Scoville, Dimitrios Xourafas, Aslam M Ejaz, Allan Tsung, Timothy Pawlik, Jordan M Cloyd

ORCID number: Steven D Scoville (0000-0003-1597-1267); Dimitrios Xourafas (0000-0002-7811-867X); Aslam M Ejaz (0000-0002-1387-2604); Allan Tsung (0000-0002-3916-8965); Timothy Pawlik (0000-0002-7994-9870); Jordan M Cloyd (0000-0002-2373-8433).

Author contributions: All authors helped to perform the research; Scoville SD, Xourafas D and Cloyd JM performed drafting conception and design, data analysis, manuscript writing; Ejaz AM, Tsung A and Pawlik T contributed to writing the manuscript.

Institutional review board statement: Not applicable as data was de-identified and publicly available.

Informed consent statement: Not applicable as data was de-identified and publicly available.

Conflict-of-interest statement: We have no financial relationships to disclose.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works

Steven D Scoville, Dimitrios Xourafas, Aslam M Ejaz, Allan Tsung, Timothy Pawlik, Jordan M Cloyd, Department of Surgery, Division of Surgical Oncology at The Ohio State University, James Cancer Center, Columbus, OH 43210, United States

Steven D Scoville, The Arthur G James Comprehensive Cancer Center and Solove Research Institute, The Ohio State University, Columbus, OH 43210, United States

Dimitrios Xourafas, Department of Surgery, Harvard Medical School, Department of Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA 02215, United States

Aslam M Ejaz, Timothy Pawlik, Jordan M Cloyd, Department of Surgery, The Ohio State University, Columbus, OH 43210, United States

Allan Tsung, Department of Surgery, The Ohio State University Wexner Medical Center, Columbus, OH 43210, United States

Jordan M Cloyd, Department of Surgery, Division of Surgical Oncology, The Ohio State University Wexner Medical Center, Columbus, OH 43210, United States

Corresponding author: Jordan M Cloyd, MD, Assistant Professor, Department of Surgery, Division of Surgical Oncology, The Ohio State University Wexner Medical Center, 410 W 10th Ave, N-907 Doan Hall, Columbus, OH 43210, United States. jordan.cloyd@osumc.edu

Abstract

BACKGROUND

Although surgical resection is associated with the best long-term outcomes for neuroendocrine liver metastases (NELM), the current indications for and outcomes of surgery for NELM from a population perspective are not well understood.

AIM

To determine the current indications for and outcomes of liver resection (LR) for NELM using a population-based cohort.

METHODS

A retrospective review of the 2014-2017 American College of Surgeons National Surgical Quality Improvement Program and targeted hepatectomy databases was performed to identify patients who underwent LR for NELM. Perioperative characteristics and 30-d morbidity and mortality were analyzed.

RESULTS

on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Received: November 21, 2019

Peer-review started: November 21, 2019

First decision: December 13, 2019

Revised: February 21, 2020

Accepted: March 5, 2020

Article in press: March 5, 2020

Published online: April 27, 2020

P-Reviewer: Campana D, Nacak M, Vagholkar K, Mrzljak A

S-Editor: Zhou JJ

L-Editor: A

E-Editor: Wu YXJ



Among 669 patients who underwent LR for NELM, the median age was 60 (interquartile range: 51-67) and 51% were male. While the number of metastases resected ranged from 1 to 9, the most common (45%) number of tumors resected was one. The majority (68%) of patients had a largest tumor size of < 5 cm. Most patients underwent partial hepatectomy (71%) while fewer underwent a right or left hepatectomy or trisectionectomy. The majority of operations were open (82%) versus laparoscopic (17%) or robotic (1%). In addition, 30% of patients underwent intraoperative ablation while 45% had another concomitant operation including cholecystectomy (28.8%), bowel resection (20.2%), or partial pancreatectomy (3.4%). Overall 30-d morbidity and mortality was 29% and 1.3%, respectively. On multivariate analysis, American Society of Anesthesiologists class ≥ 3 [odds ratios (OR), OR = 2.089, 95% confidence intervals (CI): 1.197-3.645], open approach (OR = 1.867, 95% CI: 1.148-3.036), right hepatectomy (OR = 1.618, 95% CI: 1.014-2.582), and prolonged operative time of > 230 min (OR = 1.731, 95% CI: 1.168-2.565) were associated with higher 30-d morbidity while intraoperative ablation and concomitant procedures were not.

CONCLUSION

LR for NELM was performed with relatively low postoperative morbidity and mortality. Concomitant procedures performed at the time of LR did not increase morbidity.

Key words: Carcinoid; Neuroendocrine tumor; Primary tumor resection; Intraoperative ablation; Cholecystectomy; Small bowel resection

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Surgical resection of neuroendocrine liver metastases is associated with the best long-term outcomes, however the current indications for and outcomes of surgery are not well understood. In this study, we performed a retrospective review of the 2014-2017 American College of Surgeons National Surgical Quality Improvement Program to identify 669 patients who underwent liver resection to define characteristics associated with increased 30-d postoperative morbidity and mortality. Overall morbidity and mortality were relatively low at 29% and 1.3% respectively. Factors associated with increased 30-d morbidity included open and prolonged cases (> 230 min), right hepatectomy, and American Society of Anesthesiologists class ≥ 3 while concomitant procedures including intraoperative ablation did not influence morbidity.

Citation: Scoville SD, Xourafas D, Ejaz AM, Tsung A, Pawlik T, Cloyd JM. Contemporary indications for and outcomes of hepatic resection for neuroendocrine liver metastases. *World J Gastrointest Surg* 2020; 12(4): 159-170

URL: <https://www.wjgnet.com/1948-9366/full/v12/i4/159.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v12.i4.159>

INTRODUCTION

Neuroendocrine tumors (NET) are a heterogeneous group of neoplasms that can occur anywhere in the body but commonly arise from the gastrointestinal tract. While relatively rare, the incidence and prevalence of NETs are steadily increasing, at least in part due to improved imaging and diagnostic techniques^[1,2]. Despite their low grade nature, a substantial proportion (60%-80%) of well-differentiated NETs are diagnosed with or will develop neuroendocrine liver metastases (NELM), which is one of the strongest prognostic factors among patients with NETs^[3]. For example, the 5-year overall survival of patients with untreated NELM range from 13% to 54%, compared with 61% to 79% among individuals who undergo treatment^[4-8].

Multiple treatments exist for patients with NELM including surgical resection, ablative techniques, transarterial therapies, somatostatin analogs, cytotoxic chemotherapy, targeted therapies, and peptide receptor radionuclide therapy^[4]. Other novel systemic and targeted therapies are rapidly emerging^[9,10]. Despite the absence of level I evidence, surgical resection is associated with the best long-term outcomes

based on retrospective cohort studies and meta-analyses^[6,11,12]. Indeed, even cytoreductive surgery (*i.e.*, surgical debulking) of NELM has been associated with improved overall survival if residual disease less than 10%-30% can be achieved^[5,13-15]. Based on these data, surgical resection of NELM has been recommended as the preferred initial approach, when feasible, by the European Neuroendocrine Tumors Society and North American Neuroendocrine Tumors Society^[16,17].

Previous studies evaluating the short-term outcomes of surgery for NELM have frequently been limited by their retrospective, single-institution nature^[18-22]. Other more recent multi-institutional studies have been limited to high-volume institutions and conducted over long study periods^[6,14,23]. Thus, there is a need for an evaluation of contemporary practice patterns and outcomes from a population-based perspective. Such information would inform patient selection and facilitate patient education and the informed consent process. Therefore, we utilized the American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP) targeted hepatectomy database to analyze contemporary characteristics of patients with NELM who underwent liver resection (LR) in the United States and evaluate factors associated with postoperative morbidity and mortality.

MATERIALS AND METHODS

Data acquisition and study population

The ACS-NSQIP data set is a national validated and risk adjusted outcomes based database that includes demographic, clinical, perioperative, and 30-d postoperative details of patients undergoing surgery from 600 eligible hospitals across the United States. The ACS-NSQIP uses a systemic sampling process to ensure that each case has an equal chance of selection and it is frequently monitored to minimize sample bias. A retrospective review of the 2014-2017 ACS-NSQIP and targeted hepatectomy databases were performed and patients were matched based on case ID numbers. All adult patients undergoing LR were identified using Current Procedural Terminology codes 47120, 47122, 47125, and 47130.

Study variables and outcomes

Independent variables included demographic, preoperative health status, relevant comorbidities, operative, and postoperative outcomes. Demographics included age and gender. Preoperative health included American Society of Anesthesiologists (ASA) classification, body mass index, weight loss within 6 mo from surgery, smoking, chronic steroid use, preoperative sepsis (systemic inflammatory response syndrome or septic shock) and preoperative transfusion. Comorbidities included diabetes mellitus, chronic obstructive pulmonary disease, hypertension requiring medications, bleeding disorders, congestive heart failure. Targeted hepatectomy variables included neoadjuvant therapy within 90 d of surgery, ascites, viral hepatitis, and pre-operative biliary stent placement. Operative variables consisted of operative approach (open, laparoscopic, robotic), type of resection (trisegmentectomy, total right or left hemihepatectomy, or partial hepatectomy), concomitant procedures (cholecystectomy, intestinal resection, partial pancreatectomy), size of metastatic lesion (< 2, 2-5, or > 5 cm) number of metastatic lesions (1-2, 3-5, or > 5), number of concurrent partial hepatectomies (0, 1-5, 6-9, or > 10), liver texture (normal, congested, cirrhotic or fatty), operative time, pringle maneuver during resection, concomitant intraoperative ablation (IA), and biliary reconstruction.

Overall morbidity included each of the following within 30 d from the date of surgery: Superficial, deep and organ/space surgical site infection, sepsis, respiratory complications including pneumonia or reintubation, pulmonary embolism, deep vein thrombosis, myocardial infarct, cardiac arrest, stroke, renal complications such as renal insufficiency or urinary tract infection, hemorrhage requiring at least 4 U of packed red blood cells, bile leak, liver intervention post hepatectomy and postoperative liver failure. Perioperative mortality was also measured and defined as death within 30 d after LR. Length of stay, discharge disposition, 30-d readmission and reoperation were also among the postoperative measures assessed.

Statistical analysis

Descriptive statistics were reported as percentages of the total number of patients in the study. Univariate and multivariate analyses were used to identify factors associated with the development of overall morbidity and mortality within 30 d of surgery. Logistic regression analysis was used for univariate analysis. Stepwise logistic regression analysis was used for multivariable analysis and included all non-collinear variables. Results are reported as odds ratios (OR) and 95% confidence

intervals (CI). Analyses were performed using SAS 9.2 (SAS Institute, Inc., Cary, NC, United States). *P* values < 0.05 were considered statistically significant. All statistics were performed by an experienced biostatistician.

RESULTS

Patient characteristics

Between 2014-2017, 669 patients were identified within the ACS-NSQIP database who underwent LR for NELM. Complete demographic and clinicopathologic criteria of the cohort are listed in [Table 1](#). The median age was 60 [interquartile range (IQR): 51-67], 51% were male and the mean body mass index was 27.4 (IQR: 24-32). Most patients underwent partial hepatectomy (71%) while fewer underwent hemihepatectomy (23%) or trisegmentectomy (6%). The majority of operations were open (82%); fewer were laparoscopic (17%) or robotic (1%). The mean number of lesions resected was 3.8 (range 1-9) and the most common size of the largest tumor was 2-5 cm (45%) ([Figure 1A and B](#)). Among all patients, 30.6% underwent concomitant IA in conjunction with a LR. In addition, 45.4% of patients underwent a combined resection in addition to the LR. The most common concomitant procedure was a cholecystectomy (28.8%), followed by intestinal resection (20.2%) and pancreatectomy (3.4%) ([Figure 1C](#)).

Perioperative outcomes

[Table 2](#) reports the postoperative outcomes of patients undergoing resection of NELM. Median operative time was 232 min (IQR: 179-297) and the mean length of stay was 6 (IQR: 4-8) d. The overall 30-d complication rate was 29% with the most common perioperative complications being perioperative transfusion (15.5%), intraabdominal infection (7.3%), bile leakage (5.9%), sepsis (5.6%), surgical site infection (3.5%), reoperation (3.1%), pneumonia (2.9%), liver failure (2.6%), and pulmonary embolism (2%). Of note, serious adverse events such as stroke, cardiac arrest, and myocardial infarction occurred in less than 1% of all patients undergoing resection for NELM. Postoperative mortality occurred in 1.3%. The vast majority (95%) of patients were able to be discharged home and readmission was required in 11.2% of the patients.

Predictors of postoperative morbidity

Factors associated with 30-d morbidity on univariate analysis are reported in [Table 3](#). ASA class of ≥ 3 (OR = 2.418, 95%CI: 1.422-4.113, *P* = 0.0011), open approach (OR = 1.943, 95%CI: 1.218-3.102, *P* = 0.0053), right, left or trisection hepatectomy (OR = 1.660, 95%CI: 1.169-2.355, *P* = 0.0046), and operative time > 230 min, OR = 2.403, 95%CI: 1.407-2.968, *P* = 0.0002) were all associated with increased morbidity while IA was associated with a decrease in perioperative morbidity (OR = 0.686, 95%CI: 0.476-0.988, *P* = 0.0431). Interestingly, the use of concomitant procedures (including bowel resections, cholecystectomy, or pancreatectomy), as well as the size or number of tumors were not associated with postoperative morbidity.

On multivariable logistic regression, ASA class of ≥ 3 (OR = 2.089, 95%CI: 1.197-3.645, *P* = 0.0095), open approach (OR = 1.867, 95%CI: 1.148-3.036, *P* = 0.0118), right, left or trisegmental hepatectomy (OR = 1.618, 95%CI: 1.014-2.582, *P* = 0.0437), and operative time > 230 min (OR = 1.731, 95%CI: 1.168-2.565, *P* = 0.0062) were independently associated with increased morbidity while normal liver texture was protective of overall morbidity (OR = 0.641, 95%CI: 0.433-0.950, *P* = 0.0266) ([Table 4](#)).

DISCUSSION

The incidence of NETs is increasing worldwide and a majority of patients will present with metastatic disease in their liver^[1]. NELM is a strong negative prognostic factor for survival and is associated with significant reductions in patient quality of life^[24]. While several systemic and liver-directed therapies are available, surgical resection is typically recommended when feasible^[25]. In this paper, we used a contemporary, population-based, prospective database to define the characteristics and outcomes of patients undergoing surgery for NELM in the United States. These results highlight several important findings. First, the majority of operations are being performed for small tumors in the setting of multifocal disease and typically are minor resections. Second, a significant proportion of cases are being performed concomitant with another operation, either liver IA, cholecystectomy, or (presumably) primary tumor resection. Finally, modern surgery for NELM can be performed with relatively minimal postoperative morbidity (29%) and mortality (1.3%). These results are critical

Table 1 Demographic, clinical, and operative characteristics of patients with neuroendocrine liver metastases undergoing resection

	NELM (<i>n</i> = 669)
Median age in years, <i>n</i> (IQR)	60 (51-67)
Male gender, <i>n</i> (%)	341 (51)
ASA classification, <i>n</i> (%)	
I	2 (0.3)
II	143 (21)
III	459 (69)
IV	61 (9)
Median BMI (kg/m ²), <i>n</i> (IQR)	27.4 (24-32)
Comorbidities/preoperative	
> 10% loss body weight in last 6 mo, <i>n</i> (%)	30 (4.4)
Diabetes mellitus with oral agents or insulin, <i>n</i> (%)	115 (17)
Current smoker within one yr, <i>n</i> (%)	79 (12)
Severe chronic obstructive pulmonary disease, <i>n</i> (%)	9 (0.3)
Congestive heart failure in 30 d before surgery, <i>n</i> (%)	5 (0.7)
Hypertension requiring medications, <i>n</i> (%)	326 (49)
Viral hepatitis, <i>n</i> (%)	13 (1.9)
Preoperative biliary stent, <i>n</i> (%)	13 (1.9)
Ascites within 30 d, <i>n</i> (%)	3 (0.4)
Preoperative sepsis, <i>n</i> (%)	2 (0.3)
Steroid use for a chronic condition, <i>n</i> (%)	20 (3)
Bleeding disorders, <i>n</i> (%)	18 (2.6)
Preoperative transfusion, <i>n</i> (%)	3 (0.4)
Neoadjuvant therapy, <i>n</i> (%)	119 (17.7)
Patients with concomitant procedure (%)	304 (45.4)
Total number of cholecystectomy	193 (28.8)
Total number of small/large bowel resection	135 (20.2)
Total number of partial pancreatectomy	23 (3.4)
Operative approach, <i>n</i> (%)	
Open	546 (82)
Laparoscopic	113 (17)
Robotic	10 (1)
Liver resection type, <i>n</i> (%)	
Trisegmentectomy	42 (6)
Right hepatectomy	99 (15)
Left hepatectomy	52 (8)
Partial lobectomy	476 (71)
Size of metastatic lesion, <i>n</i> (%)	
< 2 cm	152 (23)
2-5 cm	303 (45)
> 5 cm	181 (27)
Unknown	33 (5)
Number of metastatic lesions, <i>n</i> (%)	
< 2	298 (45)
3-5	168 (26)
> 5	166 (27)
Unknown	37 (2)
Concurrent partial liver resections, <i>n</i> (%)	
0	205 (30.6)
1-5	385 (57.5)
6-9	44 (6.6)
> 10	5 (0.7)
Unknown	30 (4.5)

Liver texture, <i>n</i> (%)	
Normal	190 (28)
Congested	8 (1)
Cirrhotic	12 (2)
Fatty	57 (9)
Unknown	402 (60)
Median optime in minutes, <i>n</i> (IQR)	232 (179-297)
Pringle maneuver during resection, <i>n</i> (%)	161 (24)
Biliary reconstruction, <i>n</i> (%)	14 (2)
Intraoperative ablation, <i>n</i> (%)	205 (30)
Drain placement	253 (38)

NELM: Neuroendocrine liver metastases; IQR: Interquartile range; BMI: Body mass index; ASA: American Society of Anesthesiology.

for informed preoperative discussions with patients as well as future comparative effectiveness research with other liver-directed treatments.

Recent advances in the perioperative management of patients undergoing LR have improved the safety of hepatectomy and expanded criteria for selecting patients for surgery. Indeed, a recent study by Cloyd *et al*^[26] evaluated nearly 4000 patients who underwent LR over two decades and noted steady improvements in postoperative morbidity despite increases in case complexity. Improvements in the outcomes of LR are likely multifactorial but improved patient selection, evaluation and optimization, are paramount. Accurate liver volumetry and future liver remnant augmentation have been important strategies for minimizing post hepatectomy and postoperative liver failure^[27,28]. Improved perioperative and anesthetic care, including reduced intravenous fluid administration and less blood loss, have similarly been critical advances in contemporary hepatic surgery. The introduction of enhanced recovery after surgery processes have further contributed to reduced morbidity following surgery and are now commonly routinized at major medical centers^[29]. Finally, implementation of laparoscopic and robotic approaches for surgery have led to reduced postoperative pain, blood loss, and length of hospital stay with similar outcomes compared with open approaches^[30-33].

Prior studies evaluating the role of IA for NELM have demonstrated that this therapeutic approach is generally well tolerated and is indicated for patients whose tumors are not amenable to resection^[6,34] though the safety of IA during surgery for NELM has not been thoroughly evaluated^[34-36]. In this study, we noted that IA was associated with decreased 30-d morbidity in univariate analysis though this association did not persist on multivariate analysis. These findings are consistent with a recent study evaluating IA during resection of colorectal liver metastasis which found lower overall morbidity, hospital length of stay, and readmission rates in patients who underwent LR and IA compared to patients who underwent LR alone^[37]. Based on these results and others, IA appears to be a safe and effective strategy to expand the surgical options for patients with multifocal NELM.

While the role of primary tumor resection in the setting of unresectable NELM remains controversial, resection of the primary NET is indicated when liver metastases are resectable^[19,38-41]. The current study suggests that resection of the primary (*e.g.*, pancreatectomy, intestinal resection) can be performed safely and is not associated with increased postoperative morbidity. These findings are consistent with the large body of literature which suggests that most LRs for colorectal liver metastasis can be performed safely in a combined fashion with standard colorectal resections^[42,43].

While the ACS-NSQIP targeted hepatectomy database has the advantage of containing hepatectomy-specific perioperative variables, a limitation of the current database was the lack of cancer- and patient-specific information. For example, the database lacked relevant information such as the symptomatic status of patients, functional status of tumors, tumor grade, or presence of extra-hepatic disease. In addition, it lacked Furthermore, as the study was limited to the 30-d postoperative period, we were unable to describe the long-term efficacy of LR for NELM. However, the purpose of the current study was to evaluate the indications for and short-term outcomes of surgery for NELM. Multiple prior studies have found that LR for NELM is associated with good long-term survival^[4,6,44]. Similarly, the ACS-NSQIP database did not have information on carcinoid crisis, however, previous studies have shown this to be a relatively rare event^[45,46]. This study had several other limitations,




Figure 1 Indications for liver resection for neuroendocrine liver metastases. Overall size (A) and number (B) of neuroendocrine liver metastases and frequency of concomitant procedures including intraoperative ablation (C) among patients undergoing liver resection for neuroendocrine liver metastases.

primarily related to its retrospective nature and the fact that data were limited to the 30-d postoperative period which may be insufficient to capture all complications.

In conclusion, in this contemporary population-based analysis, we demonstrated that LR can be performed for NELM with relatively low postoperative morbidity and mortality. Concomitant operations such as cholecystectomy, bowel resection, pancreatectomy, and IA can safely be performed and do not contribute to increased morbidity. Careful patient selection, minimizing operative time, and utilizing minimally invasive surgical approaches may help reduce postoperative morbidity. While multiple therapeutic options exist for NELM, given the excellent long-term outcomes observed in the literature and the satisfactory short-term outcomes demonstrated herein, surgical resection should remain the standard of care when feasible.

Table 2 Postoperative outcomes of patients with neuroendocrine liver metastases undergoing liver resection

	NELM (<i>n</i> = 669)
Post-hepatectomy	
Bile leakage, <i>n</i> (%)	40 (5.9)
Post hepatectomy invasive intervention, <i>n</i> (%)	65 (9.7)
Post hepatectomy liver failure, <i>n</i> (%)	18 (2.6)
Specific complications	
Superficial surgical site infection, <i>n</i> (%)	24 (3.5)
Deep incisional surgical site infection, <i>n</i> (%)	4 (0.6)
Organ/space surgical site infection, <i>n</i> (%)	49 (7.3)
Bleeding requiring transfusion, <i>n</i> (%)	104 (15.5)
Unplanned re-intubation, <i>n</i> (%)	9 (1.3)
Pneumonia, <i>n</i> (%)	20 (2.9)
Pulmonary embolism, <i>n</i> (%)	14 (2)
Progressive renal insufficiency, <i>n</i> (%)	7 (1.1)
Urinary tract infection, <i>n</i> (%)	14 (2)
Stroke, <i>n</i> (%)	1 (0.1)
Cardiac arrest, <i>n</i> (%)	2 (0.3)
Myocardial infarction, <i>n</i> (%)	3 (0.4)
Deep venous thrombosis/thrombophlebitis, <i>n</i> (%)	9 (1.4)
Sepsis, <i>n</i> (%)	38 (5.6)
Overall	
Median length of hospital stay in days, <i>n</i> (IQR)	6 (4-8)
Discharge destination to home, <i>n</i> (%)	637 (95.2)
30-d readmission, <i>n</i> (%)	75 (11.2)
Reoperation, <i>n</i> (%)	21 (3.1)
30-d overall morbidity, <i>n</i> (%)	194 (29)
Mortality, <i>n</i> (%)	9 (1.3)

NELM: Neuroendocrine liver metastases; IQR: Interquartile range.

Table 3 Significant predictors of 30-d overall morbidity among patients undergoing hepatectomy for neuroendocrine liver metastases based on univariate logistic regression analysis

	OR	95%CI	P value
Age > 60	0.931	0.658-1.316	0.6835
Male gender	1.232	0.890-1.707	0.2092
ASA class ≥ 3	2.418	1.422-4.113	0.0011
BMI > 27	1.030	0.743-1.428	0.8596
Preop biliary stent	2.562	0.850-7.719	0.0946
Viral hepatitis	2.538	0.841-7.660	0.0983
Concomitant bowel resection	1.278	0.885-1.844	0.1906
Concomitant cholecystectomy	1.094	0.749-1.598	0.6431
Concomitant pancreatectomy	1.579	0.626-3.984	0.3335
Open approach	1.943	1.218-3.102	0.0053
Size < 2 cm (ref)			
Size 2-5 cm	0.989	0.645-1.516	0.9591
Size > 5 cm	1.397	0.882-2.215	0.1546
Number of tumors > 1	0.984	0.681-1.422	0.9317
Right/left/triseg hepatectomy	1.660	1.169-2.355	0.0046
Abnormal liver texture	1.340	0.818-2.193	0.2447
Intraoperative ablation	0.686	0.476-0.988	0.0431
Biliary reconstruction	3.979	1.317-12.023	0.0144
Operative time > 230 min	2.043	1.407-2.968	0.0002
Pringle	1.429	0.986-2.070	0.0593

OR: Odds ratio; CI: Confidence intervals; ASA: American Society of Anesthesiologists; BMI: Body mass index.

Table 4 Significant predictors of 30-d overall morbidity among patients undergoing hepatectomy for neuroendocrine liver metastases based on multivariate stepwise logistic regression analysis

	OR	95%CI	P value
ASA class ≥ 3	2.089	1.197-3.645	0.0095
Normal liver texture	0.641	0.433-0.950	0.0266
Open approach	1.867	1.148-3.036	0.0118
Right hepatectomy	1.618	1.014-2.582	0.0437
Intraoperative ablation	0.697	0.473-1.029	0.0697
Biliary reconstruction	2.802	0.870-9.021	0.0842
Operative time > 230 min	1.731	1.168-2.565	0.0062

OR: Odds ratio; CI: Confidence intervals; ASA: American Society of Anesthesiologists.

ARTICLE HIGHLIGHTS

Research background

Multiple liver-directed therapies, including hepatic resection, exist for patients with neuroendocrine liver metastases (NELM). While surgical resection is associated with the best long-term outcomes, the current indications for and outcomes of surgery for NELM from a population perspective are not well understood.

Research motivation

A better understanding of the frequency and predictors of postoperative complications will improve shared-decision making for patients with NELM, especially given the expanding number of liver-directed and systemic therapies available.

Research objectives

The purpose of the current study was to define the current indications for surgery for NELM, characterize the short-term outcomes of patients undergoing surgery, and evaluate predictors of complications using a population-based approach.

Research methods

A retrospective review of the 2014-2017 American College of Surgeons National Surgical Quality Improvement Program targeted hepatectomy database was performed to identify patients who underwent hepatic resection for NELM. Perioperative characteristics and 30-d morbidity and mortality were analyzed.

Research results

Among 669 patients who underwent liver resection for NELM, the number of metastases resected ranged from 1 to 9 though the most common (45%) number of tumors resected was one. The majority (68%) of patients had a largest tumor size of < 5 cm and most patients underwent partial hepatectomy (71%). The majority of operations were open (82%) versus laparoscopic (17%) or robotic (1%). In addition, 30% of patients underwent intraoperative ablation while 45% had another concomitant operation including cholecystectomy (28.8%), bowel resection (20.2%), or partial pancreatectomy (3.4%). Overall 30-d morbidity and mortality was 29% and 1.3%, respectively. On multivariate analysis, American Society of Anesthesiologists class ≥ 3 , open approach, formal hemi-hepatectomy or trisectionectomy, and prolonged operative time were associated with higher 30-d morbidity. Concomitant procedures including intraoperative ablation, small bowel resection, or pancreatectomy were not independently associated with higher morbidity.

Research conclusions

In this contemporary population-based analysis, we demonstrated that hepatic resection can be performed with relatively low postoperative morbidity and mortality for patients with NELM. Concomitant operations such as cholecystectomy, bowel resection, pancreatectomy, and liver ablation can safely be performed and do not contribute to increased morbidity. Careful patient selection, minimizing operative time, and utilizing minimally invasive approaches may help reduce postoperative morbidity. While multiple therapeutic options exist for NELM, given the excellent long-term outcomes observed in the literature and the satisfactory short-term outcomes demonstrated in the current study, surgical resection should remain the standard of care when feasible.

Research perspectives

This study highlights the current population-based indications for liver resection for patients with neuroendocrine liver metastases and confirms satisfactory short-term outcomes. In light of these findings, future research should focus on expanding the indications for hepatic resection particularly given the increasing number of liver-directed and systemic therapy options available. Future prospective studies should evaluate the optimal sequencing of liver-directed therapies including neoadjuvant and adjuvant strategies to improve long-term outcomes.

REFERENCES

1. Dasari A, Shen C, Halperin D, Zhao B, Zhou S, Xu Y, Shih T, Yao JC. Trends in the Incidence, Prevalence, and Survival Outcomes in Patients With Neuroendocrine Tumors in the United States. *JAMA Oncol* 2017; **3**: 1335-1342 [PMID: 28448665 DOI: 10.1001/jamaoncol.2017.0589]
2. Lee MR, Harris C, Baeg KJ, Aronson A, Wisnivesky JP, Kim MK. Incidence Trends of Gastroenteropancreatic Neuroendocrine Tumors in the United States. *Clin Gastroenterol Hepatol* 2019; **17**: 2212-2217.e1 [PMID: 30580091 DOI: 10.1016/j.cgh.2018.12.017]
3. Spolverato G, Bagante F, Wagner D, Buettner S, Gupta R, Kim Y, Maqsood H, Pawlik TM. Quality of life after treatment of neuroendocrine liver metastasis. *J Surg Res* 2015; **198**: 155-164 [PMID: 26095419 DOI: 10.1016/j.jss.2015.05.048]
4. Farley HA, Pommier RF. Treatment of Neuroendocrine Liver Metastases. *Surg Oncol Clin N Am* 2016; **25**: 217-225 [PMID: 26610783 DOI: 10.1016/j.soc.2015.08.010]
5. Sarmiento JM, Heywood G, Rubin J, Ilstrup DM, Nagorney DM, Que FG. Surgical treatment of neuroendocrine metastases to the liver: a plea for resection to increase survival. *J Am Coll Surg* 2003; **197**: 29-37 [PMID: 12831921 DOI: 10.1016/S1072-7515(03)00230-8]
6. Mayo SC, de Jong MC, Pulitano C, Clary BM, Reddy SK, Gamblin TC, Celinksi SA, Kooby DA, Staley CA, Stokes JB, Chu CK, Ferrero A, Schulick RD, Choti MA, Mentha G, Strub J, Bauer TW, Adams RB, Aldrighetti L, Capussotti L, Pawlik TM. Surgical management of hepatic neuroendocrine tumor metastasis: results from an international multi-institutional analysis. *Ann Surg Oncol* 2010; **17**: 3129-3136 [PMID: 20585879 DOI: 10.1245/s10434-010-1154-5]
7. Chamberlain RS, Canes D, Brown KT, Saltz L, Jarnagin W, Fong Y, Blumgart LH. Hepatic neuroendocrine metastases: does intervention alter outcomes? *J Am Coll Surg* 2000; **190**: 432-445 [PMID: 10757381 DOI: 10.1016/S1072-7515(00)00222-2]
8. Bacchetti S, Bertozzi S, Londero AP, Uzzau A, Pasqual EM. Surgical treatment and survival in patients with liver metastases from neuroendocrine tumors: a meta-analysis of observational studies. *Int J Hepatol* 2013; **2013**: 235040 [PMID: 23509630 DOI: 10.1155/2013/235040]
9. Cloyd JM, Konda B, Shah MH, Pawlik TM. The emerging role of targeted therapies for advanced well-differentiated gastroenteropancreatic neuroendocrine tumors. *Expert Rev Clin Pharmacol* 2019; **12**: 101-108 [PMID: 30582383 DOI: 10.1080/17512433.2019.1561273]
10. Scoville SD, Cloyd JM, Pawlik TM. New and emerging systemic therapy options for well-differentiated gastroenteropancreatic neuroendocrine tumors. *Expert Opin Pharmacother* 2020; **21**: 183-191 [PMID: 31760823 DOI: 10.1080/14656566.2019.1694003]
11. Fairweather M, Swanson R, Wang J, Brais LK, Dutton T, Kulke MH, Clancy TE. Management of Neuroendocrine Tumor Liver Metastases: Long-Term Outcomes and Prognostic Factors from a Large Prospective Database. *Ann Surg Oncol* 2017; **24**: 2319-2325 [PMID: 28303430 DOI: 10.1245/s10434-017-5839-x]

- 12 **Woltering EA**, Voros BA, Beyer DT, Wang YZ, Thiagarajan R, Ryan P, Wright A, Ramirez RA, Ricks MJ, Boudreaux JP. Aggressive Surgical Approach to the Management of Neuroendocrine Tumors: A Report of 1,000 Surgical Cyto reductions by a Single Institution. *J Am Coll Surg* 2017; **224**: 434-447 [PMID: 28088602 DOI: 10.1016/j.jamcollsurg.2016.12.032]
- 13 **Morgan RE**, Pommier SJ, Pommier RF. Expanded criteria for debulking of liver metastasis also apply to pancreatic neuroendocrine tumors. *Surgery* 2018; **163**: 218-225 [PMID: 29103583 DOI: 10.1016/j.surg.2017.05.030]
- 14 **Ejaz A**, Reames BN, Maithel S, Poultides GA, Bauer TW, Fields RC, Weiss MJ, Marques HP, Aldrighetti L, Pawlik TM. Cyto reductive debulking surgery among patients with neuroendocrine liver metastasis: a multi-institutional analysis. *HPB (Oxford)* 2018; **20**: 277-284 [PMID: 28964630 DOI: 10.1016/j.hpb.2017.08.039]
- 15 **Scott AT**, Breheny PJ, Keck KJ, Bellizzi AM, Dillon JS, O'Dorisio TM, Howe JR. Effective cyto reduction can be achieved in patients with numerous neuroendocrine tumor liver metastases (NETLMs). *Surgery* 2019; **165**: 166-175 [PMID: 30343949 DOI: 10.1016/j.surg.2018.04.070]
- 16 **O'Toole D**, Kianmanesh R, Caplin M. ENETS 2016 Consensus Guidelines for the Management of Patients with Digestive Neuroendocrine Tumors: An Update. *Neuroendocrinology* 2016; **103**: 117-118 [PMID: 26731186 DOI: 10.1159/000443169]
- 17 **Howe JR**, Cardona K, Fraker DL, Kebebew E, Untch BR, Wang YZ, Law CH, Liu EH, Kim MK, Menda Y, Morse BG, Bergsland EK, Strosberg JR, Nakakura EK, Pommier RF. The Surgical Management of Small Bowel Neuroendocrine Tumors: Consensus Guidelines of the North American Neuroendocrine Tumor Society. *Pancreas* 2017; **46**: 715-731 [PMID: 28609357 DOI: 10.1097/MPA.0000000000000846]
- 18 **Jensen EH**, Kvols L, McLoughlin JM, Lewis JM, Alvarado MD, Yeatman T, Malafa M, Shibata D. Biomarkers predict outcomes following cyto reductive surgery for hepatic metastases from functional carcinoid tumors. *Ann Surg Oncol* 2007; **14**: 780-785 [PMID: 17146740 DOI: 10.1245/s10434-006-9148-z]
- 19 **Boudreaux JP**, Wang YZ, Diebold AE, Frey DJ, Anthony L, Uhlhorn AP, Ryan P, Woltering EA. A single institution's experience with surgical cyto reduction of stage IV, well-differentiated, small bowel neuroendocrine tumors. *J Am Coll Surg* 2014; **218**: 837-844 [PMID: 24655881 DOI: 10.1016/j.jamcoll-surg.2013.12.035]
- 20 **de Mestier L**, Neuzillet C, Hentic O, Kianmanesh R, Hammel P, Ruszniewski P. Prolonged survival in a patient with neuroendocrine tumor of the cecum and diffuse peritoneal carcinomatosis. *Case Rep Gastroenterol* 2012; **6**: 205-210 [PMID: 22649333 DOI: 10.1159/000338740]
- 21 **Chambers AJ**, Pasiacka JL, Dixon E, Rorstad O. The palliative benefit of aggressive surgical intervention for both hepatic and mesenteric metastases from neuroendocrine tumors. *Surgery* 2008; **144**: 645-51; discussion 651-653 [PMID: 18847650 DOI: 10.1016/j.surg.2008.06.008]
- 22 **Chung MH**, Pisegna J, Spirt M, Giuliano AE, Ye W, Ramming KP, Bilchik AJ. Hepatic cyto reduction followed by a novel long-acting somatostatin analog: a paradigm for intractable neuroendocrine tumors metastatic to the liver. *Surgery* 2001; **130**: 954-962 [PMID: 11742323 DOI: 10.1067/msy.2001.118388]
- 23 **Sham JG**, Ejaz A, Gage MM, Bagante F, Reames BN, Maithel S, Poultides GA, Bauer TW, Fields RC, Weiss MJ, Marques HP, Aldrighetti L, Pawlik TM, He J. The Impact of Extent of Liver Resection Among Patients with Neuroendocrine Liver Metastasis: an International Multi-institutional Study. *J Gastrointest Surg* 2019; **23**: 484-491 [PMID: 29980977 DOI: 10.1007/s11605-018-3862-2]
- 24 **Frilling A**, Sotiropoulos GC, Li J, Kornasiewicz O, Plöckinger U. Multimodal management of neuroendocrine liver metastases. *HPB (Oxford)* 2010; **12**: 361-379 [PMID: 20662787 DOI: 10.1111/j.1477-2574.2010.00175.x]
- 25 **Frilling A**, Modlin IM, Kidd M, Russell C, Breitenstein S, Salem R, Kwekkeboom D, Lau WY, Klersy C, Vilgrain V, Davidson B, Siegler M, Caplin M, Solcia E, Schilsky R; Working Group on Neuroendocrine Liver Metastases. Recommendations for management of patients with neuroendocrine liver metastases. *Lancet Oncol* 2014; **15**: e8-21 [PMID: 24384494 DOI: 10.1016/S1470-2045(13)70362-0]
- 26 **Cloyd JM**, Mizuno T, Kawaguchi Y, Lillemoen HA, Karagkounis G, Omichi K, Chun YS, Conrad C, Tzeng CD, Odisio BC, Huang SY, Hicks M, Wei SH, Aloia TA, Vauthey JN. Comprehensive Complication Index Validates Improved Outcomes Over Time Despite Increased Complexity in 3707 Consecutive Hepatectomies. *Ann Surg* 2020; **271**(4): 724-731 [PMID: 30339628 DOI: 10.1097/SLA.0000000000003043]
- 27 **Rassam F**, Olthof PB, Bennink RJ, van Gulik TM. Current Modalities for the Assessment of Future Remnant Liver Function. *Visc Med* 2017; **33**: 442-448 [PMID: 29344518 DOI: 10.1159/000480385]
- 28 **Rahnama-Azar AA**, Cloyd JM, Weber SM, Dillhoff M, Schmidt C, Winslow ER, Pawlik TM. Update on Liver Failure Following Hepatic Resection: Strategies for Prediction and Avoidance of Post-operative Liver Insufficiency. *J Clin Transl Hepatol* 2018; **6**: 97-104 [PMID: 29577036 DOI: 10.14218/JCTH.2017.00060]
- 29 **Wang C**, Zheng G, Zhang W, Zhang F, Lv S, Wang A, Fang Z. Enhanced Recovery after Surgery Programs for Liver Resection: a Meta-analysis. *J Gastrointest Surg* 2017; **21**: 472-486 [PMID: 28101720 DOI: 10.1007/s11605-017-3360-y]
- 30 **Jackson NR**, Hauch A, Hu T, Buell JF, Slakey DP, Kandil E. The safety and efficacy of approaches to liver resection: a meta-analysis. *JSLs* 2015; **19**: e2014.00186 [PMID: 25848191 DOI: 10.4293/JSLs.2014.00186]
- 31 **Nguyen KT**, Marsh JW, Tsung A, Steel JJ, Gamblin TC, Geller DA. Comparative benefits of laparoscopic vs open hepatic resection: a critical appraisal. *Arch Surg* 2011; **146**: 348-356 [PMID: 21079109 DOI: 10.1001/archsurg.2010.248]
- 32 **Tsilimigras DI**, Moris D, Vagios S, Merath K, Pawlik TM. Safety and oncologic outcomes of robotic liver resections: A systematic review. *J Surg Oncol* 2018; **117**: 1517-1530 [PMID: 29473968 DOI: 10.1002/jso.25018]
- 33 **Gani F**, Ejaz A, Dillhoff M, He J, Weiss M, Wolfgang CL, Cloyd J, Tsung A, Johnston FM, Pawlik TM. A national assessment of the utilization, quality and cost of laparoscopic liver resection. *HPB (Oxford)* 2019; **21**: 1327-1335 [PMID: 30850188 DOI: 10.1016/j.hpb.2019.02.005]
- 34 **Mazzaglia PJ**, Berber E, Milas M, Siperstein AE. Laparoscopic radiofrequency ablation of neuroendocrine liver metastases: a 10-year experience evaluating predictors of survival. *Surgery* 2007; **142**: 10-19 [PMID: 17629995 DOI: 10.1016/j.surg.2007.01.036]
- 35 **Orgera G**, Krokidis M, Cappucci M, Gourtsoyianni S, Tipaldi MA, Hatzidakis A, Rebonato A, Rossi M. Current status of Interventional Radiology in the management of Gastro-Entero-Pancreatic Neuroendocrine Tumours (GEP-NETs). *Cardiovasc Intervent Radiol* 2015; **38**: 13-24 [PMID: 25366087]

- DOI: [10.1007/s00270-014-1005-z](https://doi.org/10.1007/s00270-014-1005-z)
- 36 **Kose E**, Kahramangil B, Aydin H, Donmez M, Takahashi H, Aucejo F, Siperstein A, Berber E. Outcomes of laparoscopic tumor ablation for neuroendocrine liver metastases: a 20-year experience. *Surg Endosc* 2020; **34**: 249-256 [PMID: [30945061](https://pubmed.ncbi.nlm.nih.gov/30945061/) DOI: [10.1007/s00464-019-06759-1](https://doi.org/10.1007/s00464-019-06759-1)]
 - 37 **Xourafas D**, Pawlik TM, Cloyd JM. Early Morbidity and Mortality after Minimally Invasive Liver Resection for Hepatocellular Carcinoma: a Propensity-Score Matched Comparison with Open Resection. *J Gastrointest Surg* 2019; **23**: 1435-1442 [PMID: [30377911](https://pubmed.ncbi.nlm.nih.gov/30377911/) DOI: [10.1007/s11605-018-4016-2](https://doi.org/10.1007/s11605-018-4016-2)]
 - 38 **Hellman P**, Lundström T, Ohrvall U, Eriksson B, Skogseid B, Oberg K, Tiensuu Janson E, Akerström G. Effect of surgery on the outcome of midgut carcinoid disease with lymph node and liver metastases. *World J Surg* 2002; **26**: 991-997 [PMID: [12016480](https://pubmed.ncbi.nlm.nih.gov/12016480/) DOI: [10.1007/s00268-002-6630-z](https://doi.org/10.1007/s00268-002-6630-z)]
 - 39 **Givi B**, Pommier SJ, Thompson AK, Diggs BS, Pommier RF. Operative resection of primary carcinoid neoplasms in patients with liver metastases yields significantly better survival. *Surgery* 2006; **140**: 891-897; discussion 897-898 [PMID: [17188135](https://pubmed.ncbi.nlm.nih.gov/17188135/) DOI: [10.1016/j.surg.2006.07.033](https://doi.org/10.1016/j.surg.2006.07.033)]
 - 40 **Tsilimigras DI**, Ntanasis-Stathopoulos I, Kostakis ID, Moris D, Schizas D, Cloyd JM, Pawlik TM. Is Resection of Primary Midgut Neuroendocrine Tumors in Patients with Unresectable Metastatic Liver Disease Justified? A Systematic Review and Meta-Analysis. *J Gastrointest Surg* 2019; **23**: 1044-1054 [PMID: [30671800](https://pubmed.ncbi.nlm.nih.gov/30671800/) DOI: [10.1007/s11605-018-04094-9](https://doi.org/10.1007/s11605-018-04094-9)]
 - 41 **Xiang JX**, Zhang XF, Beal EW, Weiss M, Aldrighetti L, Poultides GA, Bauer TW, Fields RC, Maithel SK, Marques HP, Pawlik TM. Hepatic Resection for Non-functional Neuroendocrine Liver Metastasis: Does the Presence of Unresected Primary Tumor or Extrahepatic Metastatic Disease Matter? *Ann Surg Oncol* 2018; **25**: 3928-3935 [PMID: [30218247](https://pubmed.ncbi.nlm.nih.gov/30218247/) DOI: [10.1245/s10434-018-6751-8](https://doi.org/10.1245/s10434-018-6751-8)]
 - 42 **Reddy SK**, Pawlik TM, Zorzi D, Gleisner AL, Ribero D, Assumpcao L, Barbas AS, Abdalla EK, Choti MA, Vauthey JN, Ludwig KA, Mantyh CR, Morse MA, Clary BM. Simultaneous resections of colorectal cancer and synchronous liver metastases: a multi-institutional analysis. *Ann Surg Oncol* 2007; **14**: 3481-3491 [PMID: [17805933](https://pubmed.ncbi.nlm.nih.gov/17805933/) DOI: [10.1245/s10434-007-9522-5](https://doi.org/10.1245/s10434-007-9522-5)]
 - 43 **Idrees JJ**, Bagante F, Gani F, Rosinski BF, Chen Q, Merath K, Dillhoff M, Cloyd J, Pawlik TM. Population level outcomes and costs of single stage colon and liver resection versus conventional two-stage approach for the resection of metastatic colorectal cancer. *HPB (Oxford)* 2019; **21**: 456-464 [PMID: [30266492](https://pubmed.ncbi.nlm.nih.gov/30266492/) DOI: [10.1016/j.hpb.2018.08.007](https://doi.org/10.1016/j.hpb.2018.08.007)]
 - 44 **Glazer ES**, Tseng JF, Al-Refaie W, Solorzano CC, Liu P, Willborn KA, Abdalla EK, Vauthey JN, Curley SA. Long-term survival after surgical management of neuroendocrine hepatic metastases. *HPB (Oxford)* 2010; **12**: 427-433 [PMID: [20662794](https://pubmed.ncbi.nlm.nih.gov/20662794/) DOI: [10.1111/j.1477-2574.2010.00198.x](https://doi.org/10.1111/j.1477-2574.2010.00198.x)]
 - 45 **Kimbrough CW**, Beal EW, Dillhoff ME, Schmidt CR, Pawlik TM, Lopez-Aguilar AG, Poultides G, Makris E, Rocha FG, Crown A, Abbott DE, Fisher AV, Fields RC, Krasnick BA, Idrees K, Marincola-Smith P, Cho CS, Beems M, Maithel SK, Cloyd JM. Influence of carcinoid syndrome on the clinical characteristics and outcomes of patients with gastroenteropancreatic neuroendocrine tumors undergoing operative resection. *Surgery* 2019; **165**: 657-663 [PMID: [30377003](https://pubmed.ncbi.nlm.nih.gov/30377003/) DOI: [10.1016/j.surg.2018.09.008](https://doi.org/10.1016/j.surg.2018.09.008)]
 - 46 **Byrne RM**, Pommier RF. Small Bowel and Colorectal Carcinoids. *Clin Colon Rectal Surg* 2018; **31**: 301-308 [PMID: [30186052](https://pubmed.ncbi.nlm.nih.gov/30186052/) DOI: [10.1055/s-0038-1642054](https://doi.org/10.1055/s-0038-1642054)]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

