

World Journal of *Clinical Cases*

World J Clin Cases 2023 September 16; 11(26): 6031-6317



MINIREVIEWS

- 6031 Diabetes among Muslims during Ramadan: A narrative review
Ochani RK, Shaikh A, Batra S, Pikale G, Surani S

ORIGINAL ARTICLE**Retrospective Cohort Study**

- 6040 Clinical evaluation of ventilation mode on acute exacerbation of chronic obstructive pulmonary disease with respiratory failure
Wang JJ, Zhou Z, Zhang LY

Retrospective Study

- 6051 Predictive value of preoperative albumin-bilirubin score and other risk factors for short-term outcomes after open pancreatoduodenectomy
Zavrtanik H, Cosola D, Badovinac D, Hadžialjević B, Horvat G, Plevel D, Bogoni S, Tarchi P, de Manzini N, Tomažič A

- 6066 Lyophilized recombinant human brain natriuretic peptide for chronic heart failure: Effects on cardiac function and inflammation
Li F, Li H, Luo R, Pei JB, Yu XY

- 6073 Continuous renal replacement therapy with oXiris® in patients with hematologically malignant septic shock: A retrospective study
Wang J, Wei SR, Ding T, Zhang LP, Weng ZH, Cheng M, Zhou Y, Zhang M, Liu FJ, Yan BB, Wang DF, Sun MW, Cheng WX

- 6083 Serum basic fibroblast growth factor and interleukin-1 β predict the effect of first-line chemotherapy in patients with advanced gastric cancer
Zheng L, Gan LH, Yao L, Li B, Huang YQ, Zhang FB, Kuang MQ, Fang N

- 6091 Multinucleated giant cells of bladder mucosa are modified telocytes: Diagnostic and immunohistochemistry algorithm and relation to PD-L1 expression score
Gulinac M, Velikova T, Dikov D

Clinical Trials Study

- 6105 Comparing the efficacy of regen-cov, remdesivir, and favipiravir in reducing invasive mechanical ventilation need in hospitalized COVID-19 patients
Hegazy SK, Tharwat S, Hassan AH

META-ANALYSIS

- 6122 Risk factors for stroke recurrence in young patients with first-ever ischemic stroke: A meta-analysis
Xia Y, Liu H, Zhu R

SCIENTOMETRICS

- 6132** Unveiling the hidden world of gut health: Exploring cutting-edge research through visualizing randomized controlled trials on the gut microbiota
Zyoud SH, Shakhshir M, Abushanab AS, Koni A, Shahwan M, Jairoun AA, Abu Taha A, Al-Jabi SW

CASE REPORT

- 6147** Rivaroxaban for the treatment of heparin-induced thrombocytopenia with thrombosis in a patient undergoing artificial hip arthroplasty: A case report
Ly FF, Li MY, Qu W, Jiang ZS
- 6154** Mepolizumab induced palmoplantar psoriasis: A case report
Artosi F, Diluvio L, Vultaggio M, Campione E, Bianchi L
- 6159** Early diagnosis of renal pelvis villous adenoma: A case report
Li LL, Song PX, Xing DF, Liu K
- 6165** Identification of the dominant loop of a dual-loop macro-reentry left atrial flutter without prior intervention using high-density mapping technology: A case report
Yu SD, Chu YP
- 6170** Surgery for fibrous dysplasia associated with aneurysmal-bone-cyst-like changes in right proximal femur: A case report
Xie LL, Yuan X, Zhu HX, Pu D
- 6176** Efficacy of abatacept treatment in a patient with enteropathy carrying a variant of unsignificance in *CTLA4* gene: A case report
Musabak U, Erdoğan T, Ceylaner S, Özbek E, Suna N, Özdemir BH
- 6183** Postpartum hemophagocytic lymphohistiocytosis: A case report
An JH, Ahn JH
- 6189** Non-arteritic anterior ischemic optic neuropathy combined with branch retinal vein obstruction: A case report
Gong HX, Xie SY
- 6194** Large colonic lipoma with a laterally spreading tumor treated by endoscopic submucosal dissection: A case report
Bae JY, Kim HK, Kim YJ, Kim SW, Lee Y, Ryu CB, Lee MS
- 6200** T/myeloid mixed-phenotype acute leukemia treated with venetoclax and decitabine: A case report
Park S, Jeong EJ, Kang JH, Lee GW, Go SI, Lee DH, Koh EH
- 6206** Severe inflammatory disorder in trisomy 8 without myelodysplastic syndrome and response to methylprednisolone: A case report
Pan FY, Fan HZ, Zhuang SH, Pan LF, Ye XH, Tong HJ

- 6213** Aggressive variant prostate cancer: A case report and literature review
Weng XT, Lin WL, Pan QM, Chen TF, Li SY, Gu CM
- 6223** Typical Zollinger-Ellison syndrome-atypical location of gastrinoma and absence of hypergastrinemia: A case report and review of literature
Zhang JM, Zheng CW, Li XW, Fang ZY, Yu MX, Shen HY, Ji X
- 6231** Left epigastric isolated tumor fed by the inferior phrenic artery diagnosed as ectopic hepatocellular carcinoma: A case report
Liu HB, Zhao LH, Zhang YJ, Li ZF, Li L, Huang QP
- 6240** Squamous cell carcinoma associated with endometriosis in the uterus and ovaries: A case report
Cai Z, Yang GL, Li Q, Zeng L, Li LX, Song YP, Liu FR
- 6246** Intestinal obstruction due to giant liver cyst: A case report
Küçük A, Mohamed SS, Abdi AM, Ali AY
- 6252** Difficulties in diagnosing angiomatoid fibrous histiocytoma of the head and neck region: A case report
Michcik A, Bieñ M, Wojciechowska B, Polcyn A, Garbacewicz Ł, Kowalski J, Drogoszewska B
- 6262** Efficacy of tolvaptan in an infant with syndrome of inappropriate antidiuretic hormone secretion associated with holoprosencephaly: A case report
Mori M, Takeshita S, Nakamura N, Mizuno Y, Tomita A, Aoyama M, Kakita H, Yamada Y
- 6268** Recurrent hemoptysis in pediatric bronchial Dieulafoy's disease with inferior phrenic artery supply: A case report
Wang F, Tang J, Peng M, Huang PJ, Zhao LJ, Zhang YY, Wang T
- 6274** Variant of Guillain-Barré syndrome with anti-sulfatide antibody positivity and spinal cord involvement: A case report
Liu H, Lv HG, Zhang R
- 6280** Secondary pulmonary infection by *Fusarium solani* and *Aspergillus niger* during systemic steroid treatment for COVID-19: A case report
Usuda D, Kato M, Sugawara Y, Shimizu R, Inami T, Tsuge S, Sakurai R, Kawai K, Matsubara S, Tanaka R, Suzuki M, Shimozawa S, Hoichi Y, Osugi I, Katou R, Ito S, Mishima K, Kondo A, Mizuno K, Takami H, Komatsu T, Oba J, Nomura T, Sugita M
- 6289** Collision tumor of primary malignant lymphoma and adenocarcinoma in the colon diagnosed by molecular pathology: A case report and literature review
Jiang M, Yuan XP
- 6298** Successful resolution of gastric perforation caused by a severe complication of pancreatic walled-off necrosis: A case report
Noh BG, Yoon M, Park YM, Seo HI, Kim S, Hong SB, Park JK, Lee MW
- 6304** Bilateral dislocation of the long head of biceps tendon with intact rotator cuff tendon: A case report
Sohn HJ, Cho CH, Kim DH

6311 Delayed diagnosis of abdominal Henoch-Schonlein purpura in children: A case report

Guo H, Wang ZL, Tao Z

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Vikram K Mahajan, MD, Professor, Dermatology, Venereology and Leprosy, Dr. Radhakrishnan Government Medical College, Kangra 177001, Himachal Pradesh, India. vkm1@rediffmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJCC* as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Hua-Ge Yu*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jin-Lei Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

September 16, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Bilateral dislocation of the long head of biceps tendon with intact rotator cuff tendon: A case report

Hyuk-Joon Sohn, Chul-Hyun Cho, Du-Han Kim

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Koumantakis GA, Greece; Oommen AT, India

Received: July 5, 2023

Peer-review started: July 5, 2023

First decision: August 10, 2023

Revised: August 16, 2023

Accepted: August 21, 2023

Article in press: August 21, 2023

Published online: September 16, 2023



Hyuk-Joon Sohn, Chul-Hyun Cho, Du-Han Kim, Department of Orthopedic Surgery, Keimyung University Dongsan Hospital, Keimyung University School of Medicine, Daegu 42601, South Korea

Corresponding author: Du-Han Kim, MD, PhD, Doctor, Department of Orthopedic Surgery, Keimyung University Dongsan Hospital, Keimyung University School of Medicine, 1035 Dalgubul-ro, Dalseo-gu, Daegu 42601, South Korea. osmdkdh@gmail.com

Abstract

BACKGROUND

Dislocation of the long head of biceps tendon (LHBT) usually involves rotator cuff injury, and isolated dislocation with an intact rotator cuff is rare. Some cases of isolated dislocation have been reported. However, to the best of our knowledge, there has been no report of bilateral dislocation of the LHBT without rotator cuff pathology.

CASE SUMMARY

A 23-year-old male presented to our outpatient clinic with left side dominant pain in both shoulders. The patient had no history of trauma or overuse. The patient underwent intra-articular injection and physical therapy, but his symptoms aggravated. Based on preoperative imaging, the diagnosis was bilateral dislocation of the LHBT. Dysplasia of the bicipital groove was detected in both shoulders. Active dislocation of the biceps tendon over an intact subscapularis tendon was identified by diagnostic arthroscopy. Staged biceps tenodesis was performed and continuous passive motion therapy was administered immediately after surgery. The patient's pain was resolved, and full functional recovery was achieved, and he was satisfied with the condition of his shoulders.

CONCLUSION

This study describes a rare case of bilateral dislocations of the LHBT without rotator cuff injury due to dysplasia of the bicipital groove.

Key Words: Shoulder; Biceps; Dislocation; Rotator cuff; Tenodesis; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The long head of biceps tendon (LHBT), which has been regarded as a significant cause of anterior shoulder pain, often occurs with other shoulder pathologies. Trauma and repetitive mechanical wear can lead to development of biceps subluxation and dislocations. Association of sports activity and injury with LHBT pathology has also been reported. The subscapularis fiber function as the medial and proximal portion of the soft tissue sling and subscapularis lesions are almost always accompanied by dislocation. This report describes a rare case of bilateral dislocation of the LHBT without rotator cuff pathology.

Citation: Sohn HJ, Cho CH, Kim DH. Bilateral dislocation of the long head of biceps tendon with intact rotator cuff tendon: A case report. *World J Clin Cases* 2023; 11(26): 6304-6310

URL: <https://www.wjgnet.com/2307-8960/full/v11/i26/6304.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i26.6304>

INTRODUCTION

The long head of biceps tendon (LHBT), which has been regarded as a significant cause of anterior shoulder pain, often occurs with other shoulder pathologies[1,2]. LHBT, which originates from the supraglenoid tubercle, runs within the intracapsular space, making a 30-degree turn into the bicipital groove. The bicipital groove is located between the greater and lesser tuberosities; this osseous component of the bicipital groove provides stability to the LHBT. The soft tissue system, including the coracohumeral ligament (CHL), superior glenohumeral ligament (SGHL), and interwoven fibers of the subscapularis and supraspinatus tendon, are the most important stabilizers[3-5]. Trauma and repetitive mechanical wear can lead to development of biceps subluxation and dislocations. Association of sports activity and injury with LHBT pathology has also been reported[6]. These cases usually involve pulley lesions. The SGHL/CHL complex and subscapularis fiber function as the medial and proximal portion of the soft tissue sling and subscapularis lesions are almost always accompanied by dislocation[7-11].

This report describes a rare case of bilateral dislocation of the LHBT without rotator cuff pathology. Some authors have reported on cases of isolated dislocation of the LHBT with an intact rotator cuff tendon[7,8,10,12,13]. However, to the best of our knowledge, there have been no reports of bilateral isolated dislocation of the LHBT without any event including trauma or overuse.

CASE PRESENTATION

Chief complaints

A 23-year-old male presented to our clinic with left side dominant pain in both shoulders.

History of present illness

Gradual aggravation of the patient's symptoms occurred over the last year, even after physical therapy and intra-articular injection. The patient reported that raising his arm was difficult. The pain was more severe on the left side and became aggravated when lifting heavy objects.

History of past illness

The patient first visited our hospital four years ago, complaining of left side dominant pain in both shoulders. He was then diagnosed with impingement syndrome in both shoulders and visited another local hospital to undergo physical therapy and receive conservative management. After four years of conservative management, the patient returned to our clinic with aggravated pain in both shoulders.

Personal and family history

The patient was a college student with no history of heavy loading activity or overuse, and there was no history of trauma or sport injury as well. There was no other past illness or family history.

Physical examination

On physical examination, the patient's pain was localized to the bicipital groove area and lesser tuberosity in both shoulders. O'Brien test, Speed's test, and Yegerson test showed a positive result in both shoulders. The patient's range of motion (ROM) was in nearly normal range in both shoulders. The results of manual muscle testing showed no muscle weakness.

Laboratory examinations

The results of preoperative laboratory tests were normal.

Imaging examinations

A simple radiograph showed no signs of bony anomaly. Shoulder ultrasonography showed a shallow bicipital groove with an intact subscapularis tendon in both shoulders (Figure 1). Computed tomography scan showed a dysplastic bicipital groove along with flattening of the medial wall of the groove (Figure 2). Preoperative magnetic resonance imaging (MRI) showed that the LHBT was medially dislocated in both shoulders. LHBT distance was 17.9 mm on left shoulder and 20.9 mm on right shoulder[14]. The subscapularis tendon and the supraspinatus tendon were intact on both sides (Figure 3). The morphology of the bicipital groove was examined using the technique introduced by Yoo *et al*[15]. The opening angle of the bicipital groove was 124.6 degrees on the right side and 137.0 degrees on the left side. The medial wall angle was 25.7 degrees on the right side and 21.4 degrees on the left side. The depth of each bicipital groove was 2.41 mm in the right shoulder, and 1.34 mm in the left shoulder. The bicipital groove in both shoulders was considered shallow, as proposed in previous studies[15,16]. Considering the values measured in MRI, the groove was more shallow on the left side, where the pain was more severe.

FINAL DIAGNOSIS

The final diagnosis for the presented case was bilateral dislocations of the long head of bicep tendon (Walch classification type I, Bennett type 4) with dysplasia of both bicipital grooves.

TREATMENT

Diagnostic shoulder arthroscopy was first performed on the left shoulder because the patient had left side dominant pain. A partial tear and synovitis of the LHBT were observed on the initial inspection. When the LHBT was pulled medially, active dislocation of the LHBT over the intact subscapularis tendon was detected (Figure 4A, Video). An assessment of the intact rotator cuff on the bursa side was performed. Finally, subpectoral biceps tenodesis was performed using a suture anchor (Figure 5A).

Surgical treatment of the right shoulder was administered after functional recovery had been achieved, four months after treatment of the left shoulder. Fraying of the biceps tendon pulley was detected by diagnostic arthroscopy. The LHBT was actively dislocated to the anterior side of the subscapularis tendon with application of medial traction (Figure 4B). As with the left shoulder, pulley release and subpectoral tenodesis were performed using a suture anchor (Figure 5B). The supraspinatus and subscapularis tendons were also intact. The patient started immediate postoperative continuous passive motion therapy with restriction of active elbow flexion. There were no complications in either shoulder.


OUTCOME AND FOLLOW-UP

The patient's symptoms showed significant improvement after subpectoral tenodesis. Before the surgery, the visual analogue scale (VAS) score was 5 and the score for subjective shoulder value (SSV) was 50%. The UCLA score was 21 for both shoulders and the American Shoulder and Elbow Surgeons (ASES) score was 58.33 prior to the surgery. Three months after left biceps tenodesis, the VAS score had improved to 2 with nearly full ROM, the UCLA score improved to 32, SSV improved to 60%, and the ASES score improved to 86.67. At the last follow-up, which was post-operative 22 mo on the left side and 18 mo on the right side, the VAS score was 1 with full ROM, and the UCLA score was 33 for both shoulders. The SSV score was 85%, and the ASES score had improved to 85. The patient returned to his ordinary life within three months after surgery, and he was satisfied with the functional improvement without pain at six months after surgery.

DISCUSSION


Many authors have reported that dislocation of the LHBT often involves rotator cuff pathology[7,9-11]. Desai and Mata [9], who reviewed 141 shoulders with full-thickness rotator cuff tear, asserted that the incidence of LHBT pathology was significantly greater for tears involving subscapularis. In addition, an increase in the size of the rotator cuff tear was statistically related to increased incidence of LHBT pathology. Walch *et al*[10] also emphasized that subluxation of the LHBT is always associated with minor lesions of subscapularis, and dislocation over a completely intact subscapularis is rare.

However, other authors have maintained that isolated dislocation of the LHBT might occur without rotator cuff injury. Ayoubi *et al*[17] reported on a case of bilateral dislocation without detection of any subscapularis tears. The patient's symptoms were remarkably similar to those of our patient. However, our case did not involve the bifid subscapularis tendon, which is a rare congenital anomaly. The LHBT was located inside the substance of the subscapularis tendon. Instability of the biceps tendon with an intact rotator cuff due to injury in other stabilizing factors has also been noted. According to Gambill *et al*[8] and Vopat *et al*[12], isolated dislocation of the LHBT may occur as a result of injury in capsu-



DOI: 10.12998/wjcc.v11.i26.6304 Copyright ©The Author(s) 2023.

Figure 1 Ultrasonography. A: Left shoulder; B: Right shoulder. Intact subscapularis tendon (asterisk) and empty bicipital groove with dysplasia (arrow).




DOI: 10.12998/wjcc.v11.i26.6304 Copyright ©The Author(s) 2023.

Figure 2 Computed tomography scan showing bicipital groove dysplasia. A: Left shoulder; B: Right shoulder.

ligamentous structures of the rotator interval. Posterior instability of the humeral head was reported in both studies. The difference between the two cases is that one patient had posterior instability of the humeral head resulting from a posterior labral tear from a previous injury[8], while the other was the result of a patulous posterior capsule[12]. Despite similar symptoms, their cases differs from ours in that our case did not result from previous trauma and there was no posterior instability.


The stability of the LHBT can also be attributed to the osseous dimensions of the bicipital groove[3]. Yoo *et al*[15], who measured the depth of the bicipital groove, medial angle, and opening angle on shoulder MRI in order to examine the influence of bony morphology of the bicipital groove on the stability of the LHBT, concluded that bony morphology characterized by a shallow groove was associated with increased prevalence of LHBT instability. Levinsohn and Santelli [18] reported comparable results; a dislocated biceps tendon was detected on bicipital groove view on plain radiography in 58% of patients with a flattened bicipital groove and dysplasia of the lesser tuberosity. Because the medial wall of the bicipital groove functions as a trochlea about which the biceps tendon glides, the dysplasia of the medial wall and flattening of the bicipital groove observed in our case could be a reason for bilateral dislocation of the LHBT without rotator cuff tear.

In this study, the patient experienced pain in both shoulders caused by bilateral dislocation of the LHBT. Unlike a typical case, the dislocation was not associated with rotator cuff pathology or a traumatic event. Isolated dislocation of the LHBT was triggered by dysplasia of the bicipital groove in each shoulder. The patient underwent biceps tenodesis on both sides and satisfactory clinical and functional outcomes were finally achieved.



DOI: 10.12998/wjcc.v11.i26.6304 Copyright ©The Author(s) 2023.

Figure 3 Magnetic resonance imaging axial T2w. A: Left shoulder; B: Right shoulder. Dislocation of the long head of biceps tendon (triangle) over the intact subscapularis tendon (asterisk). Flattening of bicipital groove is remarkable (arrow). Long head of biceps tendon distance of left shoulder: 17.9 mm; right shoulder: 20.9 mm. LT: Lesser tuberosity; GT: Greater tuberosity.



DOI: 10.12998/wjcc.v11.i26.6304 Copyright ©The Author(s) 2023.

Figure 4 Arthroscopic photo from the posterior portal. A: Left shoulder; B: Right shoulder. Dislocation of the biceps tendon (triangle), fraying of biceps pulley (arrow), and intact subscapularis tendon (asterisk). HH: Humeral head.

CONCLUSION

We have described a rare case of bilateral dislocation of the LHBT without previous trauma and rotator cuff injury. To the best of our knowledge, this is the first reported case to demonstrate that bilateral dislocation of the biceps tendon may occur as a result of dysplasia of the bicipital groove without rotator cuff pathology. Physical examination and evaluation of the bicipital groove should be emphasized in patients with anterior shoulder pain of unknown cause even in cases where it occurs bilaterally and without trauma or repetitive overhead activity.



DOI: 10.12998/wjcc.v11.i26.6304 Copyright ©The Author(s) 2023.

Figure 5 Postoperative radiographs after subpectoral tenodesis. A: Left shoulder; B: Right shoulder.

FOOTNOTES

Author contributions: Sohn HJ and Cho CH contributed equally to this article as first authors; Cho CH and Kim DH designed the research study; Sohn HJ and Kim DH performed the research; Cho CH contributed new reagents and analytic tools; Cho CH and Sohn HJ analyzed the data and wrote the manuscript; and all authors have read and approved the final manuscript.

Informed consent statement: Informed written consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: South Korea

ORCID number: Hyuk-Joon Sohn 0009-0006-0407-6194; Chul-Hyun Cho 0000-0003-0252-8741; Du-Han Kim 0000-0002-6636-9340.

S-Editor: Wang JJ

L-Editor: A

P-Editor: Wang JJ

REFERENCES

- 1 Dines D, Warren RF, Inglis AE. Surgical treatment of lesions of the long head of the biceps. *Clin Orthop Relat Res* 1982; 165-171 [PMID: 7067280]
- 2 Kim TM, Shin MH, Baek S, Lee DR, Chung SW. Can a Two Simple Stitches Method Provide Secure Fixation Strength in Biceps Tenodesis?: Biomechanical Evaluation of Various Suture Techniques. *Clin Orthop Surg* 2022; 14: 426-433 [PMID: 36061853 DOI: 10.4055/cios21186]
- 3 Varacallo M, Seaman TJ, Mair SD. Biceps Tendon Dislocation and Instability. 2022 Sep 4. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan- [PMID: 30475566]
- 4 Park HB, Gwark JY, Na JB. Risk factors of chronic subscapularis tendon tear. *Clin Shoulder Elb* 2022; 25: 257-264 [PMID: 36330722 DOI: 10.1177/10785465221111111]

- 10.5397/cise.2021.00710]
- 5 **Kim JH**, Min YK, Park MJ, Huh JW, Park JH. Postoperative clinical outcomes and radiological healing according to deep and superficial layer detachment in first facet involving subscapularis tendon tear. *Clin Shoulder Elb* 2022; **25**: 93-100 [PMID: 35295069 DOI: 10.5397/cise.2021.00542]
 - 6 **Oh JH**, Kim JY, Nam KP, Kang HD, Yeo JH. Immediate Changes and Recovery of the Supraspinatus, Long Head Biceps Tendon, and Range of Motion after Pitching in Youth Baseball Players: How Much Rest Is Needed after Pitching? Sonoelastography on the Supraspinatus Muscle-Tendon and Biceps Long Head Tendon. *Clin Orthop Surg* 2021; **13**: 385-394 [PMID: 34484632 DOI: 10.4055/cios20187]
 - 7 **Ditsios K**, Agathangelidis F, Boutsiadis A, Karataglis D, Papadopoulos P. Long head of the biceps pathology combined with rotator cuff tears. *Adv Orthop* 2012; **2012**: 405472 [PMID: 23209915 DOI: 10.1155/2012/405472]
 - 8 **Gambill ML**, Mologne TS, Provencher MT. Dislocation of the long head of the biceps tendon with intact subscapularis and supraspinatus tendons. *J Shoulder Elbow Surg* 2006; **15**: e20-2 [PMID: 17126231 DOI: 10.1016/j.jse.2005.09.008]
 - 9 **Desai SS**, Mata HK. Long Head of Biceps Tendon Pathology and Results of Tenotomy in Full-Thickness Reparable Rotator Cuff Tear. *Arthroscopy* 2017; **33**: 1971-1976 [PMID: 28847573 DOI: 10.1016/j.arthro.2017.06.018]
 - 10 **Walch G**, Nové-Josserand L, Boileau P, Levigne C. Subluxations and dislocations of the tendon of the long head of the biceps. *J Shoulder Elbow Surg* 1998; **7**: 100-108 [PMID: 9593086 DOI: 10.1016/s1058-2746(98)90218-x]
 - 11 **Farshad-Amacker NA**, Jain Palrecha S, Farshad M. The primer for sports medicine professionals on imaging: the shoulder. *Sports Health* 2013; **5**: 50-77 [PMID: 24381700 DOI: 10.1177/1941738112468265]
 - 12 **Vopat ML**, Yang SY, Gregor CM, Kallail KJ, Saunders BM. Medial Dislocation of the Long Head of the Biceps without Concomitant Subscapularis Tear: A Case Report. *J Orthop Case Rep* 2020; **9**: 6-10 [PMID: 32548018 DOI: 10.13107/jocr.2019.v09.i06.1564]
 - 13 **Khazzam M**, George MS, Churchill RS, Kuhn JE. Disorders of the long head of biceps tendon. *J Shoulder Elbow Surg* 2012; **21**: 136-145 [PMID: 22005126 DOI: 10.1016/j.jse.2011.07.016]
 - 14 **Zappia M**, Ascione F, Di Pietto F, Fischetti M, Romano AM, Castagna A, Brunese L. Long head biceps tendon instability: diagnostic performance of known and new MRI diagnostic signs. *Skeletal Radiol* 2021; **50**: 1863-1871 [PMID: 33733694 DOI: 10.1007/s00256-021-03710-1]
 - 15 **Yoo JC**, Iyyampillai G, Park D, Koh KH. The influence of bicipital groove morphology on the stability of the long head of the biceps tendon. *J Orthop Surg (Hong Kong)* 2017; **25**: 2309499017717195 [PMID: 28659056 DOI: 10.1177/2309499017717195]
 - 16 **Abboud JA**, Bartolozzi AR, Widmer BJ, DeMola PM. Bicipital groove morphology on MRI has no correlation to intra-articular biceps tendon pathology. *J Shoulder Elbow Surg* 2010; **19**: 790-794 [PMID: 20713275 DOI: 10.1016/j.jse.2010.04.044]
 - 17 **Ayoubi R**, Darwish M, Nassour N, Aouad D, Maalouly J, Kanj V, El Rassi G. A rare case of an anatomical variation of bilateral long head of the biceps brachii tendons identified within the substance of bifid subscapularis tendons. *Asia Pac J Sports Med Arthrosc Rehabil Technol* 2021; **23**: 18-21 [PMID: 33376677 DOI: 10.1016/j.asmart.2020.09.001]
 - 18 **Levinsohn EM**, Santelli ED. Bicipital groove dysplasia and medial dislocation of the biceps brachii tendon. *Skeletal Radiol* 1991; **20**: 419-423 [PMID: 1925673 DOI: 10.1007/BF00191083]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

