

World Journal of *Clinical Cases*

World J Clin Cases 2023 October 26; 11(30): 7261-7507



Contents

Thrice Monthly Volume 11 Number 30 October 26, 2023

MINIREVIEWS

- 7261 Lower limb amputation rehabilitation status in India: A review
Swarnakar R, Yadav SL, Surendran D
- 7268 Magnetic resonance imaging for acute pancreatitis in type 2 diabetes patients
Ni YH, Song LJ, Xiao B

ORIGINAL ARTICLE

Retrospective Study

- 7277 Efficacy of lidocaine wet compress combined with red-light irradiation for chronic wounds
Bao MZ, Zhou LB, Zhao L, Zhang H, Li Y, Yang L, Tai AT
- 7284 Clinical implications of forkhead box M1, cyclooxygenase-2, and glucose-regulated protein 78 in breast invasive ductal carcinoma
Bai J, Li Y, Cai L
- 7294 Six-year analysis of key monitoring for bacterial strain distribution and antibiotic sensitivity in a hospital
Li ZY, Yang D, Hao CH
- 7302 Clinical pharmacists' involvement in carbapenem antibiotics management at Wenzhou Integrated Hospital
Xu XM, Pan CY, Zeng DL

Observational Study

- 7309 High risk for obstructive sleep apnea and risk of hypertension in military personnel: The CHIEF sleep study
Liu WN, Lin KH, Tsai KZ, Chu CC, Chang YC, Kwon Y, Lin GM

EVIDENCE-BASED MEDICINE

- 7318 Causal relationship association of cheese intake with gestational hypertension and diabetes result from a Mendelian randomization study
Zhong T, Huang YQ, Wang GM

META-ANALYSIS

- 7329 Left lateral decubitus sleeping position is associated with improved gastroesophageal reflux disease symptoms: A systematic review and meta-analysis
Simadibrata DM, Lesmana E, Amangku BR, Wardoyo MP, Simadibrata M
- 7337 Efficacy and safety of anti-vascular endothelial growth factor agents on corneal neovascularization: A meta-analysis
Lai SC, Loh EW, Chiou DI, Hong CT

- 7350** Efficacy and safety of different anti-osteoporotic drugs for the spinal fusion surgery: A network meta-analysis

He XY, Chen HX, Zhao ZR

SCIENTOMETRICS

- 7363** Construction of clinical research nurse training program based on position competence

Sun J, Shan WC, Liu JM, Zhang QQ, Ye Y, Huang ST, Zhong K

CASE REPORT

- 7372** Fatal hemophagocytic lymphohistiocytosis-induced multiorgan dysfunction secondary to *Burkholderia pseudomallei* sepsis: A case report

Sui MZ, Wan KC, Chen YL, Li HL, Wang SS, Chen ZF

- 7380** Interpeduncular cistern intrathecal targeted drug delivery for intractable postherpetic neuralgia: A case report

Fu F, Jiang XF, Wang JJ, Gong L, Yun C, Sun HT, Tang FW

- 7386** Using shape-memory alloy staples to treat comminuted manubrium sterni fractures: A case report

Zhang M, Jiang W, Wang ZX, Zhou ZM

- 7393** Lead helix winding tricuspid chordae tendineae: A case report

Liu TF, Ding CH

- 7398** Fournier gangrene in an infant, complicated with severe sepsis and liver dysfunction: A case report

Bakalli I, Heta S, Kola E, Celaj E

- 7403** Prenatal ultrasound diagnosis of congenital infantile fibrosarcoma and congenital hemangioma: Three case reports

Liang RN, Jiang J, Zhang J, Liu X, Ma MY, Liu QL, Ma L, Zhou L, Wang Y, Wang J, Zhou Q, Yu SS

- 7413** Iatrogenic bladder neck rupture due to traumatic urethral catheterization: A case report

Ekici O, Keskin E, Kocoglu F, Bozkurt AS

- 7418** Near obstructing painful anorectal mass and facial rash in a man with monkeypox: A case report

Akpoigbe K, Yannick J, Culpepper-Morgan J

- 7424** Traditional Chinese medicine for foot pain in a patient with complex regional pain syndrome: A case report

Shin WC, Kim H, Chung WS

- 7432** Diffuse large B-cell lymphoma successfully treated with amplified natural killer therapy alone: A case report

Nagai K, Nagai S, Okubo Y, Teshigawara K

- 7440** Pharmacogenomics-based individualized treatment of hypertension in preterm infants: A case report and review of the literature

Tang LF, Xu A, Liu K

- 7450** Warthin-like papillary renal cell carcinoma: A case report
Li XF, Wang ZJ, Zhang HM, Yang MQ
- 7457** Bladder stone due to late clip migration after prostatic urethral lift procedure: A case report
Bozkurt AS, Ekici O, Keskin E, Kocoglu F
- 7463** Acute-on-chronic liver failure induced by antiviral therapy for chronic hepatitis C: A case report
Zhong JL, Zhao LW, Chen YH, Luo YW
- 7469** Hemodynamic instability following intravenous dexmedetomidine infusion for sedation under brachial plexus block: Two case reports
Kim YS, Lee C, Oh J, Nam S, Doo AR
- 7475** Neonatal methicillin-resistant *Staphylococcus aureus* pneumonia-related recurrent fatal pyopneumothorax: A case report and review of literature
Li XC, Sun L, Li T
- 7485** Infrequent organ involvement in immunoglobulin G4-related prostate disease: A case report
Yu Y, Wang QQ, Jian L, Yang DC
- 7492** Gouty tenosynovitis with compartment syndrome in the hand: A case report
Lee DY, Eo S, Lim S, Yoon JS
- 7497** Acute myocardial infarction after initially diagnosed with unprovoked venous thromboembolism: A case report
Seo J, Lee J, Shin YH, Jang AY, Suh SY
- 7502** Distal clavicle fractures treated by anteroinferior plating with a single screw: Two case reports
Zhao XL, Liu YQ, Wang JG, Liu YC, Zhou JX, Wang BY, Zhang YJ

ABOUT COVER

Editorial Board Member of *World Journal of Clinical Cases*, Ravindra Shukla, MBBS, MD, Additional Professor, Department of Endocrinology and Metabolism, All India Institute of Medical Sciences, Jodhpur 342001, Rajasthan, India. ravindrashukla2@rediffmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (WJCC, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The WJCC is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Current Contents®/Clinical Medicine, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCC as 1.1; IF without journal self cites: 1.1; 5-year IF: 1.3; Journal Citation Indicator: 0.26; Ranking: 133 among 167 journals in medicine, general and internal; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xu Gao, Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Thrice Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng, Salim Surani, Jerzy Tadeusz Chudek, George Kontogeorgos, Maurizio Serati, Ja Hyeon Ku

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

October 26, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Clinical implications of forkhead box M1, cyclooxygenase-2, and glucose-regulated protein 78 in breast invasive ductal carcinoma

Jie Bai, Ying Li, Li Cai

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Kumar S, India; Myers JS, United States

Received: August 15, 2023

Peer-review started: August 15, 2023

First decision: August 31, 2023

Revised: September 11, 2023

Accepted: September 18, 2023

Article in press: September 18, 2023

Published online: October 26, 2023



Jie Bai, Department of Clinical Laboratory, Joint Logistics Support Unit 940 Hospital, Lanzhou 730030, Gansu Province, China

Ying Li, Department of Breast Surgery, The Fourth Hospital of Shijiazhuang, Shijiazhuang 050032, Hebei Province, China

Li Cai, Department of Pathology, Huai'an Maternal and Child Health Care Center, Huai'an 223002, Jiangsu Province, China

Corresponding author: Li Cai, MD, Doctor, Department of Pathology, Huai'an Maternal and Child Health Care Center, No. 104 Renmin South Road, Qingjiangpu District, Huai'an 223002, Jiangsu Province, China. aa14756893@163.com

Abstract

BACKGROUND

Breast infiltrating ductal carcinoma (BIDC) represents the largest heterotypic tumor group, and an in-depth understanding of the pathogenesis of BIDC is key to improving its prognosis.

AIM

To analyze the expression profiles and clinical implications of forkhead box M1 (FOXM1), cyclooxygenase-2 (COX-2), and glucose-regulated protein 78 (GRP78) in BIDC.

METHODS

A total of 65 BIDC patients and 70 healthy controls who presented to our hospital between August 2019 and May 2021 were selected for analysis. The peripheral blood FOXM1, COX-2, and GRP78 levels in both groups were measured and the association between their expression profiles in BIDC was examined. Additionally, we investigated the diagnostic value of FOXM1, COX-2, and GRP78 in patients with BIDC and their correlations with clinicopathological features. Furthermore, BIDC patients were followed for 1 year to identify factors influencing patient prognosis.

RESULTS

The levels of FOXM1, COX-2, and GRP78 were significantly higher in BIDC patients compared to healthy controls ($P < 0.05$), and a positive correlation was observed among them ($P < 0.05$). Receiver operating characteristic analysis

demonstrated that FOXM1, COX-2, and GRP78 had excellent diagnostic value in predicting the occurrence of BIDC ($P < 0.05$). Subsequently, we found significant differences in FOXM1, COX-2, and GRP78 levels among patients with different histological grades and metastasis statuses (with *vs* without) ($P < 0.05$). Cox analysis revealed that FOXM1, COX-2, GRP78, increased histological grade, and the presence of tumor metastasis were independent risk factors for prognostic death in BIDC ($P < 0.001$).

CONCLUSION

FOXM1, COX-2, and GRP78 exhibit abnormally high expression in BIDC, promoting malignant tumor development and closely correlating with prognosis. These findings hold significant research implications for the future diagnosis and treatment of BIDC.

Key Words: Diagnostic value; Forkhead box M1; Cyclooxygenase-2; Glucose-regulated protein 78; Clinical implications

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Forkhead box M1, cyclooxygenase-2, and glucose-regulated protein 78 exhibit elevated expression in breast invasive ductal carcinoma (BIDC) and are associated with a poor prognosis. Their diagnostic value suggests their potential as biomarkers for BIDC detection. Understanding their clinical implications can aid in the diagnosis and treatment of BIDC, contributing to improved patient outcomes.

Citation: Bai J, Li Y, Cai L. Clinical implications of forkhead box M1, cyclooxygenase-2, and glucose-regulated protein 78 in breast invasive ductal carcinoma. *World J Clin Cases* 2023; 11(30): 7284-7293

URL: <https://www.wjgnet.com/2307-8960/full/v11/i30/7284.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i30.7284>

INTRODUCTION

Breast cancer (BC) is the most prevalent malignancy in women, and its incidence has been increasing in recent years[1]. Among the various types of invasive BC, breast infiltrating ductal carcinoma (BIDC) represents the largest heterotypic tumor group, lacking distinct tissue characteristics and therefore being classified as a non-special type cancer[2]. BIDC is also the most common type of invasive BC, accounting for over 80% of cases[3]. While advancements in medical standards have contributed to a decrease in the overall mortality rate of BIDC, the prognosis for advanced BIDC patients remains unsatisfactory[4,5]. Researchers believe that a comprehensive understanding of the pathogenesis of BIDC and the discovery of new diagnostic and treatment methods are key to improving patient prognosis[6].

Forkhead box M1 (FOXM1), a member of the Fox protein family, plays a vital role in modulating the cell cycle[7]. Numerous studies have demonstrated that FOXM1 is overexpressed in various human tumors, promoting oncogenic transformation and participating in tumor occurrence and development[8,9]. Cyclooxygenase-2 (COX-2), a subtype of cyclooxygenase, has been identified as an essential component in the pathogenesis of malignant tumors and is closely associated with the occurrence and progression of BC based on previous data[10,11]. Additionally, glucose-regulated protein 78 (GRP78), a signature stress protein of the endoplasmic reticulum, has been found to induce tumor development, contribute to drug resistance, and facilitate tumor cell survival[12]. Although some studies have shown abnormal expression of FOXM1, COX-2, and GRP78 in BC[13-15], their specific roles in BIDC have yet to be characterized.

To develop new diagnostic and treatment protocols and identify targets for the prevention of BIDC, this study analyzed the expression profiles of FOXM1, COX-2, and GRP78 in BIDC and investigated their clinical implications in the disease. The objective was to establish the correlation between these markers and BIDC, thus laying a foundation for further research.

MATERIALS AND METHODS

Specimen collection

This study obtained approval from the Medical Ethics Committee and enrolled 65 BIDC patients [research group (RG)] and 70 healthy controls [control group (CG)], who presented to our hospital between August 2019 and May 2021. Informed consent was obtained from all participants.

Eligibility criteria

Eligible patients had histopathologically confirmed BIDC[16] and exhibited normal organ function, complete clinical data, cooperation with treatment and follow-up, no contraindications to chemotherapy, and a life expectancy of ≥ 3 mo. Pregnant and lactating women were excluded from the study. Additionally, individuals with immune deficiency, inflam-

matory diseases, severe hematopoietic injury, a history of other malignant tumors, cardio-cerebrovascular diseases, and poor treatment compliance were excluded.

Sampling and testing

Fasting venous blood samples were collected from all participants upon admission. Total RNA was isolated from the blood using Trizol, followed by reverse transcription into cDNA for polymerase chain reaction (PCR) detection. The PCR reaction was conducted under the following conditions: 95 °C/30 s and 40 cycles of 95 °C/5 s, 60 °C/30 s, and 72 °C/30 s. The design and synthesis of primer sequences (Table 1) were performed by Tsingke Biotechnology Co., Ltd. The relative expression of the target genes to β -actin was calculated using the $2^{-\Delta\Delta CT}$.

Follow-up for patient prognosis

BIDC patients were followed for 1 year through regular hospital reexaminations, with reexamination intervals not exceeding 2 mo. Death was considered a termination event, and the patient's prognosis and survival were recorded.

Outcome measures

The expression profiles of FOXM1, COX-2, and GRP78 and their correlations in BIDC were analyzed. The diagnostic value of the three markers in BIDC patients and their correlations with clinicopathological features were also investigated. Finally, the factors influencing patient prognosis were analyzed based on the follow-up results.

Statistical analysis

Statistical analyses were performed using SPSS 22.0 software. Count data, such as previous medical history, are presented as percentages (%), and inter-group differences were assessed using the Chi-square test. Expression levels of FOXM1 and COX-2 and other measurement data are presented as the mean \pm standard deviation, and inter-group and multi-group differences were identified using independent sample *t*-test and analysis of variance with Bonferroni *post-hoc* tests, respectively. Correlation analysis was conducted using the Pearson correlation coefficient, the diagnostic value was determined using receiver operating characteristic (ROC) analysis, and factors related to patient prognosis were identified using COX regression analysis. A significance level of $P < 0.05$ was used to indicate statistical significance.

RESULTS

Comparison of baseline clinical data

Baseline clinical data, including age, family history, and smoking habits, were collected at admission. The analysis revealed no significant inter-group differences ($P > 0.05$, Table 2), confirming the comparability of the study groups.

Comparison of FOXM1, COX-2, and GRP78 expression levels

The mRNA expression level of FOXM1 was (4.63 ± 0.76) in the RG and (3.94 ± 0.73) in the CG, indicating significantly higher FOXM1 levels in BIDC patients ($P < 0.05$, Figure 1A). Similarly, COX-2 mRNA expression in the RG was (3.40 ± 0.77), which was also higher compared to that in the CG ($P < 0.05$, Figure 1B). Finally, the inter-group comparison showed a higher GRP78 mRNA level in the RG compared to the CG ($P < 0.05$, Figure 1C).

Diagnostic value of FOXM1, COX-2, and GRP78 in BIDC

ROC analysis revealed that when the peripheral blood mRNA level of FOXM1 was > 4.05 , it had a sensitivity of 81.54% and specificity of 55.71% for diagnosing BIDC ($P < 0.05$, Figure 2A). Similarly, COX-2 had a sensitivity of 69.23% and specificity of 70.00% for BIDC diagnosis when its mRNA level was > 3.05 ($P < 0.05$, Figure 2B). GRP78 exhibited a sensitivity of 63.08% and specificity of 74.29% for diagnosing BIDC when its mRNA level was > 2.76 ($P < 0.05$, Figure 2C).

Relationship between the expression levels of FOXM1, COX-2, and GRP78

The Pearson correlation analysis demonstrated a positive correlation between FOXM1 and both COX-2 and GRP78 ($P < 0.05$, Figure 3A and B), as well as a positive association between COX-2 and GRP78 ($P < 0.05$, Figure 3C), in the peripheral blood of patients in the RG ($P < 0.05$, Figure 3C).

Correlation of FOXM1, COX-2, and GRP78 with clinicopathological features of BIDC

There were no significant differences in FOXM1, COX-2, and GRP78 levels among patients of different age groups ($P > 0.05$). However, significant differences in their levels were observed in patients with different histological grades and metastasis statuses (with vs without) ($P > 0.05$, Table 3), indicating a close relationship between FOXM1, COX-2, and GRP78 and the above indexes.

Univariate analysis of prognostic mortality in BIDC

During the follow-up period, 12 patients died. Deceased patients exhibited higher age, FOXM1, COX-2, and GRP78 levels compared to the surviving patients, with a higher proportion of histological grade III and tumor metastasis ($P < 0.05$, Table 4). These findings suggest that age, FOXM1, COX-2, GRP78, histological grade, and tumor metastasis were individual factors affecting the prognostic death of BIDC patients.

Table 1 Primer sequences

	Forward (5'-3')	Reverse (5-3')
FOXM1	CAGTCCGATTAGTCAGCTCCT	GTCATTAGCTCCTTGCTG
COX-2	CCGGGTACAATCGCACTTAT	GGCGCTCAGCCATACAG
GRP78	GTTACAATCAAGGTAT	CATTCACATCTATCTCAA
β-actin	CTGATATAGCCGCGCTCG	CACTCGGTGCCGATCATCA

FOXM1: Forkhead box M1; COX-2: Cyclooxygenase-2; GRP78: Glucose-regulated protein 78.

Table 2 Comparison of baseline clinical data

	Control group (n = 70)	Research group (n = 65)	<i>t</i> / χ^2	<i>P</i> value
Age (yr)	60.37 ± 5.25	60.26 ± 7.26	0.101	0.920
Family history of BC			0.457	0.499
Yes	8 (11.43)	10 (15.38)		
No	62 (88.57)	55 (84.62)		
Long-term smoking			0.099	0.753
Yes	20 (28.57)	17 (26.15)		
No	50 (71.43)	48 (73.85)		
Long-term drinking			0.182	0.669
Yes	12 (17.14)	13 (20.00)		
No	58 (82.86)	52 (80.00)		
History of breast disease			0.223	0.637
Yes	20 (28.57)	21 (32.31)		
No	50 (71.43)	44 (67.69)		

BC: Breast cancer.




Figure 1 Comparison of forkhead box M1, cyclooxygenase-2, and glucose-regulated protein 78 expression levels between the two groups. A: Forkhead box M1 expression levels; B: Cyclooxygenase-2 expression levels; C: Glucose-regulated protein 78 expression levels. $^aP < 0.05$. FOXM1: Forkhead box M1; COX-2: Cyclooxygenase-2; GRP78: Glucose-regulated protein 78.

Multivariate analysis of prognostic mortality in BIDC


Univariate indicators of BIDC (Age: ≤ 60 assigned to 0, > 60 assigned to 1; histological grade: Grade I assigned to 0, grade II assigned to 1, grade III assigned to 2; tumor metastasis: No assigned to 0, yes assigned to 1; FOXM1, COX-2, and GRP78 were analyzed using raw data) were input as covariates for COX regression analysis, with patient death used as the dependent variable. Age was not found to be an independent factor for the prognostic death of BIDC patients ($P > 0.05$). However, FOXM1, COX-2, and GRP78, increased histological grade, and tumor metastasis were identified as significant

Table 3 Correlation of forkhead box M1, cyclooxygenase-2, and glucose-regulated protein 78 with clinicopathological features of breast infiltrating ductal carcinoma

	<i>n</i>	FOXM1	COX-2	GRP78
Age (yr)				
≤ 60	33	4.51 ± 0.69	3.22 ± 0.61	3.85 ± 0.71
> 60	32	4.75 ± 0.83	3.60 ± 0.87	3.12 ± 0.69
Family history of BC				
Yes	10	4.66 ± 0.61	3.42 ± 0.65	3.05 ± 0.64
No	55	4.43 ± 1.37	3.31 ± 1.28	2.62 ± 0.95
Long-term smoking				
Yes	17	4.65 ± 0.49	3.59 ± 0.54	2.96 ± 0.49
No	48	4.62 ± 0.84	3.34 ± 0.83	2.99 ± 0.77
Long-term drinking				
Yes	13	4.36 ± 0.80	3.36 ± 0.91	2.81 ± 0.58
No	52	4.69 ± 0.75	3.41 ± 0.74	3.03 ± 0.73
History of breast disease				
Yes	21	4.49 ± 0.63	3.42 ± 0.61	2.77 ± 0.59
No	44	4.69 ± 0.82	3.40 ± 0.84	3.08 ± 0.74
Histological grade				
Grade I	35	4.39 ± 0.76	3.21 ± 0.72	2.75 ± 0.66
Grade II	19	4.57 ± 0.47	3.29 ± 0.61	2.91 ± 0.49
Grade III	11	5.48 ± 0.61 ^{1,2}	4.22 ± 0.70 ^{1,2}	3.83 ± 0.52 ^{1,2}
Tumor metastasis				
Yes	21	5.03 ± 0.75	3.7 ± 0.78	3.41 ± 0.69
No	44	4.43 ± 0.70 ³	3.24 ± 0.71 ³	2.78 ± 0.62 ³

¹Indicates a statistically significant difference from grade I patients.²Indicates a statistically significant difference from grade II patients.³Indicates a statistically significant difference from metastatic patients.

FOXM1: Forkhead box M1; COX-2: Cyclooxygenase-2; GRP78: Glucose-regulated protein 78; BC: Breast cancer.



DOI: 10.12998/wjcc.v11.i30.7284 Copyright ©The Author(s) 2023.

Figure 2 Diagnostic value of forkhead box M1, cyclooxygenase-2, and glucose-regulated protein 78 in breast infiltrating ductal carcinoma. A: Forkhead box M1; B: Cyclooxygenase-2; C: Glucose-regulated protein 78. AUC: Area under the curve; CI: Confidence interval.

Table 4 Univariate analysis of prognostic mortality in breast infiltrating ductal carcinoma

	Surviving patients (n = 53)	Dead patients (n = 12)	<i>t</i> / χ^2	<i>P</i> value
Age (yr)			9.849	0.002 ^b
≤ 60	31 (58.49)	1 (8.33)		
> 60	22 (41.51)	11 (91.67)		
Family disease history of BC			0.562	0.453
Yes	9 (16.98)	1 (8.33)		
No	44 (83.02)	11 (91.67)		
Long-term smoking			0.686	0.408
Yes	15 (28.30)	12 (16.67)		
No	38 (71.70)	10 (83.33)		
Long-term drinking			0.102	0.749
Yes	11 (20.75)	12 (16.67)		
No	42 (79.25)	10 (83.33)		
History of breast disease			1.646	0.200
Yes	19 (35.85)	12 (16.67)		
No	34 (64.15)	10 (83.33)		
Histological grade			37.350	< 0.001 ^b
Grade I	35 (66.04)	0 (0.0)		
Grade II	16 (30.19)	3 (25.00)		
Grade III	2 (3.77)	9 (75.00)		
Tumor metastasis			30.840	< 0.001 ^b
Yes	9 (16.98)	12 (100.0)		
No	44 (83.02)	0 (0.0)		
FOXM1	4.49 ± 0.69	5.22 ± 0.80	3.214	0.002 ^b
COX-2	3.28 ± 0.70	3.96 ± 0.84	2.928	0.005 ^b
GRP78	2.83 ± 0.62	3.66 ± 0.66	4.140	< 0.001 ^b

^b*P* < 0.01 *vs* surviving patients.

FOXM1: Forkhead box M1; COX-2: Cyclooxygenase-2; GRP78: Glucose-regulated protein 78; BC: Breast cancer.

factors (*P* < 0.001, Table 5).

DISCUSSION

The incidence of BC has been increasing in recent years, and it is affecting women at a younger age, posing significant risks to their health and well-being[17]. Early-stage BIDC often goes undetected, leading to a diagnosis at advanced stages with tumor metastasis, contributing to the poor prognosis of BIDC[18]. Therefore, analyzing the expression profiles and clinical implications of FOXM1, COX-2, and GRP78 in BIDC is essential for advancing research in this field.

This study confirmed the high expression levels of FOXM1, COX-2, and GRP78 in BIDC, which is consistent with previous research on BC[19-21]. These findings suggest the involvement of FOXM1, COX-2, and GRP78 in the onset and progression of BIDC. Furthermore, correlation analysis revealed positive associations between the expression levels of FOXM1, COX-2, and GRP78, indicating a synergistic relationship in their abnormal expression. Previous studies have shown that FOXM1 regulates tumor cell activity, including promoting liver cancer cell growth through kinesin superfamily protein 4A and influencing the quiescence and survival state of leukemia stem cells by regulating mixed lineage leukemia[22,23]. Additionally, FOXM1 plays a role in angiogenesis, endothelial cell proliferation, migration, and endothelial cell tube formation[24]. Vascular endothelial growth factor (VEGF), secreted by tumor tissues, induces tumor neovascularization, promoting tumor cell invasion and metastasis[25]. The close correlation between VEGF and tumor-distant metastasis has been well documented[26]. FOXM1 can bind to the VEGF promoter and transcriptionally regulate

Table 5 Multivariate analysis of prognostic mortality in breast infiltrating ductal carcinoma

	β	S.E.	Wald χ^2	P value	OR	95%CI
Age (yr)	-0.426	2.623	3.621	0.527	1.642	0.842-3.513
Histological grade	1.124	0.342	10.624	< 0.001	3.068	1.512-6.084
Tumor metastasis	1.824	0.493	20.184	< 0.001	6.521	2.164-14.813
FOXM1	1.109	0.384	8.147	0.004	3.024	1.541-6.038
COX-2	0.674	0.281	5.264	0.021	1.871	1.064-3.257
GRP78	1.084	0.573	11.806	< 0.001	1.583	0.962-4.823

FOXM1: Forkhead box M1; COX-2: Cyclooxygenase-2; GRP78: Glucose-regulated protein 78; OR: Odds ratio; CI: Confidence interval.




Figure 3 Relationship between expression levels of forkhead box M1, cyclooxygenase-2, and glucose-regulated protein 78 in breast infiltrating ductal carcinoma. A: Correlation between forkhead box M1 (FOXM1) and cyclooxygenase-2 (COX-2) mRNA expression; B: Correlation between FOXM1 and glucose-regulated protein 78 (GRP78) mRNA expression; C: Correlation between COX-2 and GRP78 mRNA expression. FOXM1: Forkhead box M1; COX-2: Cyclooxygenase-2; GRP78: Glucose-regulated protein 78.

its expression[27]. Therefore, it is speculated that this may be the mechanism by which FOXM1 is involved in BIDC. COX-2, a subtype of COX, is minimally expressed under normal physiological conditions but is overexpressed in pathological states. By promoting prostaglandin synthesis, COX-2 increases the risk of tumor growth, proliferation, vascular permeability, and metastasis[28]. Furthermore, COX-2 enhances the anti-apoptotic abilities of tumor cells by upregulating the expression of the proto-oncogene B-cell lymphoma-2 (Bcl-2). It also inhibits lymphokine production, reduces T and B cell proliferation through the production of prostaglandin E2, weakens the immune surveillance function, and makes tumor cells prone to immune escape by inducing the expression of the immunosuppressive enzyme 3-dioxygenase[29]. Increased COX-2 represents an increased risk of cancer cell growth, proliferation, and metastasis, a stronger anti-apoptotic ability, and an increased risk of adverse prognosis in patients undergoing chemotherapy. GRP78, a molecular chaperone, assists in protein folding and transport in the endoplasmic reticulum, cytoplasm, and cell membrane. It also participates in the activation of unfolded protein response signals[30]. Studies have shown that GRP78 promotes cell proliferation and migration, and its mechanism involves inhibiting Bcl-2 and binding to caspase-7 to prevent apoptosis [31,32]. Previous studies on FOXM1, COX-2, and GRP78 have all indicated their association with tumor invasion or migration, thus suggesting that their abnormal expression in BIDC is expected. Moreover, ROC analysis revealed that the three are effective in diagnosing the occurrence of BIDC, which may significantly contribute to promoting early diagnosis in the future. Clinical data analysis demonstrated that FOXM1, COX-2, and GRP78 were closely related to the histological grade and metastasis of BIDC. This reinforced the relationship between the three markers and the pathological progression of BIDC, indicating that abnormally elevated levels of FOXM1, COX-2, and GRP78 can promote the malignant development of BIDC. Finally, the prognosis analysis revealed that FOXM1, COX-2, GRP78, histological grade, and tumor metastasis were independent factors affecting the prognostic outcomes of patients. Histological grade and tumor metastasis, being typical pathological features, naturally influence the malignant progression of BIDC[33]. The analysis results for FOXM1, COX-2, and GRP78 also support the above viewpoint, confirming their role in promoting the malignant development of BIDC. Conversely, these results suggest that targeted silencing of FOXM1, COX-2, and GRP78 may inhibit BIDC progression and facilitate the treatment of BIDC.

However, the exact mechanism by which FOXM1, COX-2, and GRP78 contribute to BIDC still needs to be further confirmed through experiments. Additionally, this study did not analyze tissue samples from BIDC patients, and the short follow-up time limits the evaluation of long-term prognosis. These areas should be addressed and supplemented in future research.

CONCLUSION

The findings of this study indicate that FOXM1, COX-2, and GRP78 exhibit abnormally high expression levels in BIDC, contributing to the malignant development of the tumor and significantly impacting the prognosis of BIDC patients. These results hold significant research implications for the future diagnosis and treatment of BIDC.

ARTICLE HIGHLIGHTS

Research background

Breast infiltrating ductal carcinoma (BIDC) represents the largest heterotypic tumor group, and this study analyzed the clinical significance of forkhead box M1 (FOXM1), cyclooxygenase-2 (COX-2), and glucose-regulated protein 78 (GRP78) in BIDC, which could provide a reliable foundation for subsequent studies.

Research motivation

FOXM1, COX-2, and GRP78 are closely related to breast cancer development and progression, but their roles in BIDC remain unclear. They may play equally important roles and hold promise for future diagnosis and treatment of BIDC.

Research objectives

To analyze the clinical significance of FOXM1, COX-2, and GRP78 in BIDC, and to provide references and new research directions for future diagnosis and treatment of BIDC.

Research methods

In this study, we analyzed the clinical significance of FOXM1, COX-2, and GRP78 in BIDC by detecting the expression levels of FOXM1, COX-2, and GRP78 in the peripheral blood of patients with BIDC and healthy people.

Research results

FOXM1, COX-2, and GRP78 were elevated in BIDC and demonstrated excellent diagnostic and prognostic assessment of BIDC.

Research conclusions

FOXM1, COX-2, and GRP78 exhibit abnormally high expression in BIDC, promoting malignant tumor development and closely correlating with prognosis.

Research perspectives

This study demonstrated the clinical significance of FOXM1, COX-2, and GRP78 in BIDC, and in the future, they can be used in the clinic as reference indexes for the diagnosis, disease evaluation, and prognosis assessment of BIDC. In addition, they can also be used as targets to study new targeted therapeutic options for BIDC in the future.

FOOTNOTES

Author contributions: Bai J and Cai L designed the research study, analyzed the data, and wrote the manuscript; Bai J, Cai L, and Li Y performed the research, and contributed new reagents and analytic tools; all authors have read and approved the final manuscript.

Institutional review board statement: This study was reviewed and approved by the Ethics Committee of the Joint Logistics Support Unit 940 Hospital.

Informed consent statement: All study participants provided written informed consent.

Conflict-of-interest statement: We have no financial relationships to disclose.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Jie Bai 0009-0007-0425-850X; Li Cai 0009-0005-7870-984X.

S-Editor: Qu XL

L-Editor: Wang TQ

P-Editor: Yu HG

REFERENCES

- 1 Fahad Ullah M. Breast Cancer: Current Perspectives on the Disease Status. *Adv Exp Med Biol* 2019; **1152**: 51-64 [PMID: 31456179 DOI: 10.1007/978-3-030-20301-6_4]
- 2 Ullah L, Hameed Y, Ejaz S, Raashid A, Iqbal J, Ullah I, Ejaz SA. Detection of novel infiltrating ductal carcinoma-associated BReast CAncer gene 2 mutations which alter the deoxyribonucleic acid-binding ability of BReast CAncer gene 2 protein. *J Cancer Res Ther* 2020; **16**: 1402-1407 [PMID: 33342804 DOI: 10.4103/jert.JCRT_861_19]
- 3 Zhao H. The prognosis of invasive ductal carcinoma, lobular carcinoma and mixed ductal and lobular carcinoma according to molecular subtypes of the breast. *Breast Cancer* 2021; **28**: 187-195 [PMID: 32812198 DOI: 10.1007/s12282-020-01146-4]
- 4 Dai D, Shi R, Wang Z, Zhong Y, Shin VY, Jin H, Wang X. Competing Risk Analyses of Medullary Carcinoma of Breast in Comparison to Infiltrating Ductal Carcinoma. *Sci Rep* 2020; **10**: 560 [PMID: 31953417 DOI: 10.1038/s41598-019-57168-2]
- 5 Kader T, Hill P, Zethoven M, Goode DL, Elder K, Thio N, Doyle M, Semple T, Sufyan W, Byrne DJ, Pang JB, Murugasu A, Miligy IM, Green AR, Rakha EA, Fox SB, Mann GB, Campbell IG, Gorringer KL. Atypical ductal hyperplasia is a multipotent precursor of breast carcinoma. *J Pathol* 2019; **248**: 326-338 [PMID: 30843206 DOI: 10.1002/path.5262]
- 6 Chen XY, Yeong J, Thike AA, Bay BH, Tan PH. Prognostic role of immune infiltrates in breast ductal carcinoma in situ. *Breast Cancer Res Treat* 2019; **177**: 17-27 [PMID: 31134489 DOI: 10.1007/s10549-019-05272-2]
- 7 Barger CJ, Branick C, Chee L, Karpf AR. Pan-Cancer Analyses Reveal Genomic Features of FOXM1 Overexpression in Cancer. *Cancers (Basel)* 2019; **11** [PMID: 30795624 DOI: 10.3390/cancers11020251]
- 8 Dan VM, Raveendran RS, Baby S. Resistance to Intervention: Paclitaxel in Breast Cancer. *Mini Rev Med Chem* 2021; **21**: 1237-1268 [PMID: 33319669 DOI: 10.2174/1389557520999201214234421]
- 9 Ros S, Wright AJ, D'Santos P, Hu DE, Hesketh RL, Lubling Y, Georgopoulou D, Lerda G, Couturier DL, Razavi P, Pelossof R, Batra AS, Mannion E, Lewis DY, Martin A, Baird RD, Oliveira M, de Boo LW, Linn SC, Scaltriti M, Rueda OM, Bruna A, Caldas C, Brindle KM. Metabolic Imaging Detects Resistance to PI3K α Inhibition Mediated by Persistent FOXM1 Expression in ER(+) Breast Cancer. *Cancer Cell* 2020; **38**: 516-533.e9 [PMID: 32976773 DOI: 10.1016/j.ccell.2020.08.016]
- 10 Peng Y, Wang Y, Tang N, Sun D, Lan Y, Yu Z, Zhao X, Feng L, Zhang B, Jin L, Yu F, Ma X, Lv C. Andrographolide inhibits breast cancer through suppressing COX-2 expression and angiogenesis via inactivation of p300 signaling and VEGF pathway. *J Exp Clin Cancer Res* 2018; **37**: 248 [PMID: 30314513 DOI: 10.1186/s13046-018-0926-9]
- 11 Regulski M, Regulska K, Prukala W, Piotrowska H, Stanis B, Murias M. COX-2 inhibitors: a novel strategy in the management of breast cancer. *Drug Discov Today* 2016; **21**: 598-615 [PMID: 26723915 DOI: 10.1016/j.drudis.2015.12.003]
- 12 Cook KL, Clarke PA, Clarke R. Targeting GRP78 and antiestrogen resistance in breast cancer. *Future Med Chem* 2013; **5**: 1047-1057 [PMID: 23734687 DOI: 10.4155/fmc.13.77]
- 13 Zhang YL, Ma Y, Zeng YQ, Liu Y, He EP, Liu YT, Qiao FL, Yu R, Wang YS, Wu XY, Leng P. A narrative review of research progress on FoxM1 in breast cancer carcinogenesis and therapeutics. *Ann Transl Med* 2021; **9**: 1704 [PMID: 34988213 DOI: 10.21037/atm-21-5271]
- 14 de Souza CP, Alves B, Waisberg J, Fonseca F, Carmo AO, Gehrke F. Detection of COX-2 in liquid biopsy in patients with breast cancer. *J Clin Pathol* 2020; **73**: 826-829 [PMID: 32376713 DOI: 10.1136/jclinpath-2020-206576]
- 15 Zheng Y, Liu P, Wang N, Wang S, Wang B, Li M, Chen J, Situ H, Xie M, Lin Y, Wang Z. Betulinic Acid Suppresses Breast Cancer Metastasis by Targeting GRP78-Mediated Glycolysis and ER Stress Apoptotic Pathway. *Oxid Med Cell Longev* 2019; **2019**: 8781690 [PMID: 31531187 DOI: 10.1155/2019/8781690]
- 16 Chen Y, Zhou JH, Fan HX, Luo Y, Peng YL, Ma BY. Ultrasound Diagnosis of Breast Lymphoma and the Identification of Breast Infiltrating Ductal Carcinoma. *J Ultrasound Med* 2020; **39**: 1203-1211 [PMID: 31891216 DOI: 10.1002/jum.15209]
- 17 Odle TG. Precision Medicine in Breast Cancer. *Radiol Technol* 2017; **88**: 401M-421M [PMID: 28298516]
- 18 Lan T, Lu Y, Luo H, He J, Hu Z, Xu H. Effects of Marital Status on Prognosis in Women with Infiltrating Ductal Carcinoma of the Breast: A Real-World 1: 1 Propensity-Matched Study. *Med Sci Monit* 2020; **26**: e923630 [PMID: 32581209 DOI: 10.12659/MSM.923630]
- 19 Sun HL, Men JR, Liu HY, Liu MY, Zhang HS. FOXM1 facilitates breast cancer cell stemness and migration in YAP1-dependent manner. *Arch Biochem Biophys* 2020; **685**: 108349 [PMID: 32209309 DOI: 10.1016/j.abb.2020.108349]
- 20 Feriancová M, Walter I, Singer CF, Gazdarica J, Pohlodek K. Expression of COX-2, p16, and Ki67 in the range from normal breast tissue to breast cancer. *Neoplasma* 2021; **68**: 342-351 [PMID: 33147051 DOI: 10.4149/neo_2020_200731N798]
- 21 Lager TW, Conner C, Keating CR, Warshaw JN, Panopoulos AD. Cell surface GRP78 and Dermcidin cooperate to regulate breast cancer cell migration through Wnt signaling. *Oncogene* 2021; **40**: 4050-4059 [PMID: 33981001 DOI: 10.1038/s41388-021-01821-6]
- 22 Hu G, Yan Z, Zhang C, Cheng M, Yan Y, Wang Y, Deng L, Lu Q, Luo S. FOXM1 promotes hepatocellular carcinoma progression by regulating KIF4A expression. *J Exp Clin Cancer Res* 2019; **38**: 188 [PMID: 31072351 DOI: 10.1186/s13046-019-1202-3]
- 23 Sheng Y, Yu C, Liu Y, Hu C, Ma R, Lu X, Ji P, Chen J, Mizukawa B, Huang Y, Licht JD, Qian Z. FOXM1 regulates leukemia stem cell quiescence and survival in MLL-rearranged AML. *Nat Commun* 2020; **11**: 928 [PMID: 32066721 DOI: 10.1038/s41467-020-14590-9]
- 24 Lin JZ, Wang WW, Hu TT, Zhu GY, Li LN, Zhang CY, Xu Z, Yu HB, Wu HF, Zhu JG. FOXM1 contributes to docetaxel resistance in castration-resistant prostate cancer by inducing AMPK/mTOR-mediated autophagy. *Cancer Lett* 2020; **469**: 481-489 [PMID: 31738958 DOI: 10.1016/j.canlet.2019.11.014]
- 25 Liu C, Barger CJ, Karpf AR. FOXM1: A Multifunctional Oncoprotein and Emerging Therapeutic Target in Ovarian Cancer. *Cancers (Basel)* 2021; **13** [PMID: 34205406 DOI: 10.3390/cancers13123065]
- 26 Hisada Y, Mackman N. Tissue Factor and Cancer: Regulation, Tumor Growth, and Metastasis. *Semin Thromb Hemost* 2019; **45**: 385-395 [PMID: 31096306 DOI: 10.1055/s-0039-1687894]
- 27 Monteiro LJ, Cubillos S, Sanchez M, Acuña-Gallardo S, Venegas P, Herrera V, Lam EW, Varas-Godoy M, Illanes SE. Reduced FOXM1 Expression Limits Trophoblast Migration and Angiogenesis and Is Associated With Preeclampsia. *Reprod Sci* 2019; **26**: 580-590 [PMID: 31456179 DOI: 10.1007/978-3-030-20301-6_4]

29848205 DOI: [10.1177/1933719118778798](https://doi.org/10.1177/1933719118778798)]

- 28 **Hugo HJ**, Saunders C, Ramsay RG, Thompson EW. New Insights on COX-2 in Chronic Inflammation Driving Breast Cancer Growth and Metastasis. *J Mammary Gland Biol Neoplasia* 2015; **20**: 109-119 [PMID: [26193871](https://pubmed.ncbi.nlm.nih.gov/26193871/) DOI: [10.1007/s10911-015-9333-4](https://doi.org/10.1007/s10911-015-9333-4)]
- 29 **Qin Q**, Ji H, Li D, Zhang H, Zhang Z, Zhang Q. Tumor-associated macrophages increase COX-2 expression promoting endocrine resistance in breast cancer *via* the PI3K/Akt/mTOR pathway. *Neoplasia* 2021; **68**: 938-946 [PMID: [34619972](https://pubmed.ncbi.nlm.nih.gov/34619972/) DOI: [10.4149/neo_2021_201226N1404](https://doi.org/10.4149/neo_2021_201226N1404)]
- 30 **Zheng Y**, Zhang J, Huang W, Zhong LLD, Wang N, Wang S, Yang B, Wang X, Pan B, Situ H, Lin Y, Liu X, Shi Y, Wang Z. Sini San Inhibits Chronic Psychological Stress-Induced Breast Cancer Stemness by Suppressing Cortisol-Mediated GRP78 Activation. *Front Pharmacol* 2021; **12**: 714163 [PMID: [34912211](https://pubmed.ncbi.nlm.nih.gov/34912211/) DOI: [10.3389/fphar.2021.714163](https://doi.org/10.3389/fphar.2021.714163)]
- 31 **Liao M**, Wang C, Yang B, Huang D, Zheng Y, Wang S, Wang X, Zhang J, Tang C, Xu Z, He Y, Huang R, Zhang F, Wang Z, Wang N. Autophagy Blockade by Ai Du Qing Formula Promotes Chemosensitivity of Breast Cancer Stem Cells Via GRP78/ β -Catenin/ABCG2 Axis. *Front Pharmacol* 2021; **12**: 659297 [PMID: [34149413](https://pubmed.ncbi.nlm.nih.gov/34149413/) DOI: [10.3389/fphar.2021.659297](https://doi.org/10.3389/fphar.2021.659297)]
- 32 **Zielinska HA**, Daly CS, Alghamdi A, Bahl A, Sohail M, White P, Dean SR, Holly JMP, Perks CM. Interaction between GRP78 and IGFBP-3 Affects Tumorigenesis and Prognosis in Breast Cancer Patients. *Cancers (Basel)* 2020; **12** [PMID: [33352865](https://pubmed.ncbi.nlm.nih.gov/33352865/) DOI: [10.3390/cancers12123821](https://doi.org/10.3390/cancers12123821)]
- 33 **Zhou X**, Zheng Z, Li Y, Zhao W, Lin Y, Zhang J, Sun Q. The clinical features and prognosis of patients with mucinous breast carcinoma compared with those with infiltrating ductal carcinoma: a population-based study. *BMC Cancer* 2021; **21**: 536 [PMID: [33975551](https://pubmed.ncbi.nlm.nih.gov/33975551/) DOI: [10.1186/s12885-021-08262-0](https://doi.org/10.1186/s12885-021-08262-0)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

