

78614_Auto_Edited.docx

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 78614

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Case series of endobronchial ultrasound-guided transbronchial needle aspiration for the diagnosis of intrathoracic lymphadenopathy in patients with extrathoracic malignancy

EBUS-TBNA in extrathoracic malignancy

Shijie Li, Qi Wu

Abstract

BACKGROUND

Endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA) for the diagnosis of mediastinal and hilar lymph is poorly studied in patients with extrathoracic malignancies.

AIM

To evaluate the value of EBUS-TBNA for the diagnosis of enlarged intrathoracic lymph nodes in patients with extrathoracic malignancies.

METHODS

This was a retrospective study of patients with extrathoracic malignancies who were referred to Peking University Cancer Hospital from January 2013 to December 2018 for EBUS-TBNA due to intrathoracic lymphadenopathy. The specimens were defined as positive for malignancy, negative for non-malignancy (tuberculosis, sarcoidosis, etc), and without a definitive diagnosis. Sensitivity, negative predictive value (NPV) for malignancy, and overall accuracy were calculated. Complications were recorded.

RESULTS

A total of 80 patients underwent EBUS-TBNA and had a final diagnosis, among which 50 (62.5%) were diagnosed with extrathoracic malignancy with intrathoracic lymph nodes metastasis, 14 (17.5%) were diagnosed with primary lung cancer with nodal involvement, and 16 (20.0%) exhibited benign behavior including tuberculosis, sarcoidosis and reactive lymphadenitis or who had benign follow-up. The diagnostic sensitivity, NPV, and accuracy of EBUS-TBNA for intrathoracic lymphadenopathy in patients with extrathoracic malignancy were 93.8% ($n = 60/64$), 80.0% ($n = 16/20$), and 95.0% ($n = 76/80$), respectively. In the multivariate analysis, longer short axis of the lymph node (odds ratio [OR]: 1.200, 95% confidence interval [CI]: 1.024-1.407; $P = 0.024$)

and synchronous lung lesion (OR: 19.449, 95%CI: 1.875-201.753; $P = 0.013$) were independently associated with malignant intrathoracic lymphadenopathy. No characteristics of the lymph nodes and EBUS-TBNA were associated with the location of malignant intrathoracic lymphadenopathy, and no major complication was observed.

CONCLUSION

EBUS-TBNA is a simple and accurate procedure for the diagnosis of intrathoracic lymphadenopathy with extrathoracic malignancy.

Key Words: Endobronchial ultrasound; Intrathoracic lymphadenopathy; Extrathoracic malignancy; Transbronchial needle aspiration; Diagnosis

Li S, Wu Q. Case series of endobronchial ultrasound-guided transbronchial needle aspiration for the diagnosis of intrathoracic lymphadenopathy in patients with extrathoracic malignancy. *World J Clin Cases* 2022; In press

Core Tip: This was a retrospective study of patients referred to Peking University Cancer Hospital from January 2013 to December 2018 for endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA) due to intrathoracic lymphadenopathy. The specimens were defined as positive for malignancy, negative for non-malignancy (tuberculosis, sarcoidosis, etc), and without definite diagnosis. Sensitivity, negative predictive value for malignancy, and overall accuracy were calculated. EBUS-TBNA was found to be a simple and accurate procedure for the diagnosis of intrathoracic lymphadenopathy with extrathoracic malignancy.

INTRODUCTION

Intrathoracic lymphadenopathy is a common incidental finding by computed tomography (CT) or positron emission tomography (PET)-CT in cases with synchronous or metachronous extrathoracic malignancies^[1-3]. In such conditions, the

causes of mediastinal or hilar nodal enlargement may be distal metastasis of the extrathoracic lesion, metastasis from a primary lung cancer synchronous with the extrathoracic malignancy, or even benign lesions including tuberculosis, granulomatous inflammation, and reactive changes^[3-5]. In all of these situations, pathologic confirmation of the enlarged lymph node is crucial for the proper staging and management of patients^[1].

Mediastinoscopy is considered the “gold standard” in nodal evaluation for intrathoracic lymphadenopathy, but it is invasive and requires general anesthesia^[6]. Endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA) is currently the preferred modality to sample both mediastinal and hilar lymph nodes in primary lung cancer because it is not only minimally invasive but also can be performed under moderate conscious sedation or even under intratracheal surface anesthesia only^[7,8]. Tournoy *et al*^[9] showed that EBUS-TBNA is a diagnostic test for lung lesions after failed diagnostic bronchoscopy. A number of studies have shown that EBUS-TBNA is of value for the staging of lung cancer, as reviewed by Sehgal *et al*^[10]. EBUS-TBNA is also of value for the diagnosis of mediastinal lymphoma^[11]. In clinical practice, EBUS-TBNA could be considered the first-line examination for suspicious mediastinal lymph nodes in patients with extrathoracic cancer, preventing surgery in 50% of them^[12]; in contrast, the use of EBUS-TBNA in intrathoracic lymphadenopathy in patients with extrathoracic solid organ malignancy is a relatively less investigated topic^[13,14].

Therefore, the aim of this study was to evaluate the value of EBUS-TBNA for the diagnosis of enlarged intrathoracic lymph nodes in patients with extrathoracic malignancies.

MATERIALS AND METHODS

Patients

This study was designed as a single-center retrospective case series study. Data from patients who were referred to Peking University Cancer Hospital (Beijing, China) from

January 2013 to December 2018 for EBUS-TBNA due to intrathoracic lymphadenopathy were retrieved from the hospital database. The study was conducted according to good clinical practice and the Declaration of Helsinki. The protocol was approved by the ethics committee of Peking University Cancer Hospital (No. 2018YJZ72), which waived the need for individual consent.

The inclusion criteria for patients were synchronous or metachronous extrathoracic solid organ malignancy, and available radiological data including from CT or PET-CT scan. The exclusion criteria were synchronous or metachronous lymphoma or leukemia, or follow-up of < 12 mo when no definite diagnosis could be obtained by EBUS-TBNA with/without other interventional procedures including mediastinoscopy or thoracoscopy.

EBUS-TBNA procedures

Before EBUS-TBNA, the target lymph nodes for sampling were selected according to enlarged mediastinal or hilar lymph node with a short axis > 10 mm in thorax CT or maximum standardized uptake (SUV_{max}) value > 2.5 in PET-CT. The lymph node map was determined according to the classification proposed by the International Association for the Study of Lung Cancer^[15].

EBUS-TBNA was performed in an outpatient setting using a flexible bronchoscope (BF-UC260F-OL8; Olympus, Tokyo, Japan) by 1 of 2 experienced endoscopists (QW or SJL). First, local anesthesia was applied *via* aerosol inhalation and intratracheal spray of 2% lidocaine. Then, the EBUS scope was introduced, and all reachable lymph node stations were examined. Sampling was performed from mediastinal and hilar lymph nodes that had been previously identified as suspicious by imaging and were able to be accessed by EBUS-TBNA. For each target lesion, real-time punctures were made using a standard 22-gauge needle (ECHO-HD-22-EBUS-O; Cook Medical, Bloomington, IN, United States). Attempts were made to acquire both cytological and histological specimens during sampling, if possible (Figure 1). Major complications (*e.g.*, serious

hemorrhage [> 100 mL], pneumothorax, and post-procedure infection) that occurred during and/or after surgery were recorded.

Pathological examinations

The cytological samples were prepared as air-dried smears on glass slides and in liquid fixative for thinprep cytological test. The histological specimens were fixed in formalin solution, and immunohistochemistry was performed when necessary (Figure 1).

The specimens obtained from EBUS-TBNA were defined as positive for malignancy, negative for non-malignancy (tuberculosis, sarcoidosis, etc), and without definite diagnosis. In cases of no definite diagnosis, mediastinoscopy, video-assisted thoracic surgery (commonly known as VATS), or repeated EBUS-TBNA was recommended; patients who refused were subject to at least 12 mo of radiological and clinical follow-up every 3 mo. Follow-up was censored on December 31, 2018.

Statistical analyses

SPSS 20.0 software (IBM Corp., Armonk, NY, United States) was used for the statistical analyses. Continuous variables are presented as either means \pm standard deviations (normal distribution, Kolmogorov-Smirnov test) or median (minimum-maximum) (non-normal distribution). Categorical variables are presented as numbers and percentages and were analyzed using either the χ^2 test or the Fisher's exact test, as appropriate. Sensitivity for malignancy, negative predictive value (NPV) for non-malignancy, and overall accuracy were calculated. Predictors of malignant lymphadenopathy were modeled using logistic regression (enter method); variables with $P < 0.05$ in univariate analyses were included in multivariate analysis. $P < 0.05$ was considered statistically significant.

RESULTS

Patient characteristics

A total of 1204 patients who underwent EBUS-TBNA were reviewed. After exclusion of 1124 cases [without known extrathoracic malignancy ($n = 1112$), with previously or concurrent lymphoma or leukemia ($n = 7$), with sites of puncture not at the intrathoracic lymph node ($n = 3$), and with follow-up of less than 12 mo ($n = 2$)], 80 fulfilled the eligibility criteria (Figure 2). Among the cases included in the study (Table 1), the most common extrathoracic malignancies (65%) were breast, colorectal, and gastric cancers.

Details of the EBUS-TBNA procedures

The 80 included patients had a total of 123 enlarged lymph nodes sampled by EBUS-TBNA (median of one lymph node every patient; range: 1-4). Cytological specimens were successfully acquired from all 80 patients, and histological specimens were available for 74 (92.5%). The paratracheal region was the most common site (42.2%) for puncture (Table 2). The median number of puncture times was two for each lymph node (range: 1-6).

Pathology of lymph nodes

EBUS-TBNA diagnosed intrathoracic nodal metastasis from extrathoracic malignancy in 47 (58.8%) patients, and intrathoracic lymphadenopathy due to primary lung cancer in 13 (16.3%) patients (including 8 Lung adenocarcinomas, 3 Lung squamous carcinomas, and 2 small cell lung cancers). Regarding the 9 patients with benign diagnosis, sarcoidosis was diagnosed in 3, caseous granulomatosis with suspected tuberculosis in 3, and non-caseating granulomatous inflammation in 3. For the last 3, the inflammation was clinically inconsistent with sarcoidosis because of negative fungal and mycobacterial cultures. All 3 refused further interventional procedures, and after a median follow-up of 15 (14-19) mo, clinical and radiological results suggested benign behavior (all enlarged lymph nodes remained stable).

EBUS-TBNA found normal lymph node tissue or non-specific tissue for pathologic diagnosis in 11 (13.8%) patients. Six of them received surgical intervention (four mediastinoscopy procedures and two VATS procedures) to obtain pathologic results.

Reactive change was found in 2 patients, tuberculosis was found in 1, squamous carcinoma of lung with hilar lymph node metastasis in 1, rectal cancer lung metastasis with nodal involvement in 1, and renal cell cancer with mediastinal nodal involvement in 1. The remaining 5 patients who refused surgical intervention received periodical clinical and radiologic follow-up. One patient had progressive lymphadenopathy that was clinically considered metastasis from extrathoracic malignancy (colon cancer) and accepted the recommendation of systemic chemotherapy by multidisciplinary team. The remaining 4 patients showed a favorable outcome (1 was stable and 3 showed regressive lymphadenopathy) during a median follow-up of 16 (13-18) mo.

Diagnostic accuracy

Ultimately, 50 (62.5%) patients were diagnosed with extrathoracic malignancy with intrathoracic lymph nodes metastasis and 14 (17.5%) with primary lung cancer with nodal involvement; the remaining 16 (20.0%) patients exhibited benign behavior including tuberculosis, sarcoidosis and reactive lymphadenitis, or who had benign follow-up. The diagnostic sensitivity, NPV, and accuracy of EBUS-TBNA for intrathoracic lymphadenopathy in patients with extrathoracic malignancy were 93.8% ($n = 60/64$), 80.0% ($n = 16/20$), and 95.0% ($n = 76/80$), respectively.

Predictors of malignant lymphadenopathy

Univariate analyses revealed that longer short axis of the lymph node and synchronous lung lesion were associated with the presence of metastatic lymphadenopathy ($P = 0.003$ and $P = 0.001$, respectively). In the logistic regression multivariate model, longer short axis of the lymph node (odds ratio [OR]: 1.200, 95% confidence interval [CI]: 1.024-1.407) and synchronous lung lesion (OR: 19.449, 95%CI: 1.875-201.753) were independently associated with malignant intrathoracic lymphadenopathy ($P = 0.024$ and $P = 0.013$, respectively; Table 3).

Univariate analyses demonstrated that no characteristics of the lymph nodes themselves and EBUS-TBNA were associated with the yield of malignant intrathoracic lymphadenopathy (Table 4).

Safety of the ESD procedure

No major complications occurred during the EBUS-TBNA procedures. Twelve (15.0%) patients experienced transient hypoxemia and all recovered after immediate increase of oxygen flow.

DISCUSSION

EBUS-TBNA for the diagnosis of mediastinal and hilar lymph is poorly studied in patients with extrathoracic malignancies. This study's evaluation of the value of EBUS-TBNA for the diagnosis of enlarged intrathoracic lymph nodes in patients with extrathoracic malignancies suggested that the procedure is simple and accurate for diagnosis of intrathoracic lymphadenopathy with extrathoracic malignancy. Moreover, there were no complications.

With the rapid development of cancer therapies, overall survival of patients has improved significantly^[16]. Regular follow-up and surveillance are essential and, unfortunately, the detection of intrathoracic lymphadenopathy is not uncommon, due in part to improvements in imaging technologies. Up to 30% of extrathoracic malignancies may lead to intrathoracic lymph node metastasis^[17]. The most common solid malignancies responsible include breast, colorectal, head and neck, melanoma, kidney, and stomach cancers^[14,18,19]. This study showed that breast, colorectal, gastric, and renal cell cancers accounted for 73.4% (59/80) of the extrathoracic malignancies in our patients; of note, however, differences in other cancer types might simply be due to the differences in cancer types treated at our hospital.

Primary lung cancer with nodal involvement accounts for the majority of mediastinal and hilar lymphadenopathy cases^[20], but Mehta *et al*^[13] showed a high frequency of benign diagnoses. EBUS-TBNA is considered safe and feasible for tissue sampling with

access to both mediastinal and hilar lymph nodes, which is recommended by both the American College of Chest Physicians and European Society of Thoracic Surgeons in primary lung cancer^[7,8]. In a meta-analysis that included nine studies and 1066 patients, EBUS-TBNA had pooled sensitivity of 90%, accuracy of 96%, and NPV of 93% for mediastinal staging^[21]. EBUS-TBNA also showed an acceptable diagnostic ability in determining intrathoracic lymphadenopathy with extrathoracic malignancy. In a meta-analysis of six studies (553 patients) by Yang *et al*^[17], EBUS-TBNA provided pooled sensitivity of 85%, accuracy of 85%, and negative likelihood ratio of 16%. The present study revealed sensitivity, accuracy, and NPV of 93.8%, 95.0%, and 80.0%, respectively, by EBUS-TBNA for intrathoracic lymphadenopathy in patients with extrathoracic malignancy, in accordance with a previous study^[17]. As intrathoracic metastasis always indicates an advanced stage of the primary extrathoracic malignancy, early and accurate identification of the nature of the lymph node is important for staging, treatment strategy, and prognosis. Considering the promising diagnostic value of EBUS-TBNA, the procedure could be recommended as the first diagnostic procedure for mediastinal and hilar lymphadenopathies seen in extrathoracic malignancies.

Regarding granulomatous lymphadenitis, the diagnostic yield of EBUS-TBNA seems efficacious. A meta-analysis of 15 studies with 553 patients found that the pooled diagnostic accuracy for sarcoidosis was 79% by EBUS-TBNA^[22]. Concerning tuberculous lymphadenitis diagnosis, in a meta-analysis of 14 studies with 684 patients, EBUS-TBNA showed a pooled diagnostic yield of 80%^[23]. In regard to EBUS-TBNA for non-caseating granulomatous, more attention should be paid. Sanz-Santos *et al*^[14] reported a case of non-caseating granulomatous detected by EBUS-TBNA that was diagnosed with lymphoma after 6 mo of follow-up. Kitamura *et al*^[24] reported that 6% of patients with thoracic malignancy may have sarcoid-like reactions in non-metastatic lymph nodes. The present study found 3 patients with tuberculosis and 3 with sarcoidosis, but we could not obtain specimens for the 3 granulomatous cases and their benign behavior was merely determined through follow-up. These results suggest that in the case of the possibility of undiscovered malignancy, all granulomatous

lymphadenitis cases diagnosed by EBUS-TBNA without obvious infection or sign of sarcoidosis should be followed with more attention or surgical interventions should be advised. Evison *et al*^[25] suggested that for suspicious mediastinal and/or hilar lymph nodes but negative EBUS-TBNA, follow-up rather than resampling could be an appropriate approach.

The radiologic diagnosis of nodal involvement is mostly based on morphologic changes such as increase in size, coexistence of pulmonary lesions on CT scans, and lymph nodes > 2 cm in short axis^[26]. In this study, both univariate and multivariate regression suggested that the diameter of the lymph node and synchronous lung lesion can be malignancy indicators. As intrathoracic lymph node metastasis is relatively common in primary lung cancer^[4], patients with a history of extrathoracic malignancy, exhibiting intrathoracic lymphadenopathy combined with lung lesion, should be distinguished between distal metastasis and extrathoracic lesion and between intrathoracic metastasis and lung cancer. In such conditions, EBUS-TBNA may provide help in pathologic evidence acquisition.

PET-CT is widely used in cancer staging and distant metastasis detection. Generally, a SUV_{max} value > 2.5 could be clinically correlated with the risk of malignancy, and a SUV_{max} value > 6.3 is considered malignant with sensitivity and specificity of 70.6% and 83.3%, respectively^[27]. In the present study, no significant relationship was found between the SUV_{max} value and the presence of malignancy. This may be due to the possibility that increased SUV_{max} values can also be seen in benign pathologies involving inflammation such as sarcoidosis and infectious disease^[5]. Moreover, not all patients in our study underwent PET-CT, which may have also caused data bias.

As with other interventional procedures, the high diagnostic yield of EBUS-TBNA is associated not only with lesion-related factors such as size, location, and metabolic activity but also with procedure-related factors such as the operator's experience, number of passes, and number of lymph nodes sampled^[28]. In this study, none of the above factors showed a significant relationship with accurate diagnosis in malignancy. That may be caused by mixture of multiple lesion categories.

This study had a couple of ³⁵limitations. First, it was a retrospective study, which intrinsically has ^a certain selection bias. Second, several cases without definite diagnosis by EBUS-TBNA did not have histological confirmation. In the future, a well-designed prospective study should overcome the limitations above.

CONCLUSION

¹In conclusion, EBUS-TBNA is a simple and accurate procedure for the diagnosis of intrathoracic lymphadenopathy with extrathoracic malignancy. Its application resulted in no major complications.

ARTICLE HIGHLIGHTS

Research background

¹²Endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA) is an established technique ^{for the diagnosis of mediastinal} and hilar lymphadenectasis ⁱⁿ primary ^{lung cancer}, but is poorly studied in patients with extrathoracic malignancies.

Research motivation

Regular follow-up and surveillance are essential in cancer patients, ¹and the detection of intrathoracic ^{lymphadenopathy in those} ^{with extrathoracic malignancies is} not ^{uncommon}. EBUS-TBNA is recommended for tissue sampling both ^{in mediastinal and} ^{hilar lymph nodes in lung cancer}. ²⁷Data on the usefulness ^{of this technique in patients} ^{with} extrathoracic malignancies remain ^{limited}. ¹⁹

Research objectives

²⁰In this study, we describe ^{our experience with the use} ^{of EBUS-TBNA in patients with} extrathoracic malignancies ^{due to} intrathoracic lymphadenopathy. ⁶

Research methods

The results of the sample acquired by EBUS-TBNA were defined as positive for malignancy, negative for non-malignancy (tuberculosis, sarcoidosis, etc), and without definite diagnosis. Sensitivity, negative predictive value (NPV) for malignancy, and overall accuracy were calculated. Complications were recorded.

Research results

The diagnostic sensitivity, NPV, and accuracy of EBUS-TBNA for intrathoracic lymphadenopathy in patients with extrathoracic malignancy were 93.8% ($n = 60/64$), 80.0% ($n = 16/20$), and 95.0% ($n = 76/80$), respectively. Longer short axis of the lymph node ($P = 0.024$) and synchronous lung lesion ($P = 0.013$) were independently associated with malignant intrathoracic lymphadenopathy. No major complication was observed.

Research conclusions

EBUS-TBNA is a simple and accurate procedure for the diagnosis of intrathoracic lymphadenopathy with extrathoracic malignancy. Its application resulted in no major complications.

Research perspectives

This retrospective study demonstrates that EBUS-TBNA is effective and safe for diagnosis of intrathoracic lymphadenopathy in patients with extrathoracic malignancy. Additional prospective studies are warranted to establish standards for higher diagnostic yield.

ACKNOWLEDGEMENTS

We thank Zhonghu He for his generous help with the data analyses.

30%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|----------|--|-----------------------|
| 1 | www.ncbi.nlm.nih.gov
<small>Internet</small> | 141 words — 4% |
| <hr/> | | |
| 2 | discovery.ucl.ac.uk
<small>Internet</small> | 89 words — 3% |
| <hr/> | | |
| 3 | cyberleninka.org
<small>Internet</small> | 68 words — 2% |
| <hr/> | | |
| 4 | Bo Yang, Fang Li, Weiwei Shi, Hui Liu, Shengjie Sun, Guoqing Zhang, Shunchang Jiao. "Endobronchial ultrasound-guided transbronchial needle biopsy for the diagnosis of intrathoracic lymph node metastases from extrathoracic malignancies: A meta-analysis and systematic review", <i>Respirology</i> , 2014
<small>Crossref</small> | 50 words — 2% |
| <hr/> | | |
| 5 | Neal Navani, Matthew Nankivell, Ian Woolhouse, Richard N. Harrison et al. "Endobronchial Ultrasound-Guided Transbronchial Needle Aspiration for the Diagnosis of Intrathoracic Lymphadenopathy in Patients with Extrathoracic Malignancy: A Multicenter Study", <i>Journal of Thoracic Oncology</i> , 2011
<small>Crossref</small> | 40 words — 1% |
| <hr/> | | |
| 6 | www.science.gov
<small>Internet</small> | 39 words — 1% |

-
- 7 Kemal Can Tertemiz, Aylin Ozgen Alpaydin, Volkan Karacam. "The role of endobronchial ultrasonography for mediastinal lymphadenopathy in cases with extrathoracic malignancy", Surgical Endoscopy, 2016
Crossref 31 words — 1%
-
- 8 "INVITED ABSTRACTS", Journal of Thoracic Oncology, 2013
Crossref 30 words — 1%
-
- 9 Ravindra M. Mehta, Pavankumar Biraris, Shekhar Patil, Abhinav Singla, Kumar Kallur, Stefano Gasparini. "Utility of EBUS-TBNA in PET-positive mediastinal lymph nodes in subjects with extra-thoracic malignancy", PLOS ONE, 2019
Crossref 29 words — 1%
-
- 10 "Interventions in Pulmonary Medicine", Springer Science and Business Media LLC, 2018
Crossref 26 words — 1%
-
- 11 worldwidescience.org
Internet 26 words — 1%
-
- 12 www.pubfacts.com
Internet 26 words — 1%
-
- 13 docplayer.net
Internet 22 words — 1%
-
- 14 Hongbin Hu, Sheng An, Tong Sha, Feng Wu, Yinghui Jin, Lulan Li, Zhenhua Zeng, Jie Wu, Zhongqing Chen. "Association between dexmedetomidine administration and outcomes in critically ill patients with sepsis-associated acute kidney injury", Journal of Clinical Anesthesia, 2022
Crossref 21 words — 1%

15	assets.researchsquare.com Internet	20 words — 1%
16	bsdwebstorage.blob.core.windows.net Internet	20 words — 1%
17	journals.sagepub.com Internet	19 words — 1%
18	app.trdizin.gov.tr Internet	18 words — 1%
19	journal.publications.chestnet.org Internet	18 words — 1%
20	"Abstracts", Journal of Thoracic Oncology, 2009 Crossref	17 words — 1%
21	f6publishing.blob.core.windows.net Internet	17 words — 1%
22	www.aging-us.com Internet	17 words — 1%
23	Dongkai Shan, Yipu Ding, Xi Wang, Zinuan Liu et al. "Incremental diagnostic value of perivascular fat attenuation index for identifying hemodynamically significant ischemia with severe calcification", Research Square Platform LLC, 2022 Crossref Posted Content	16 words — < 1%
24	ERER, Onur Fevzi, ANAR, Ceyda, EROL, Serhat and ÖZKAN, Serir. "The utility of EBUS-TBNA in mediastinal or hilar lymph node evaluation in extrapulmonary malignancy", TÜBİTAK, 2016. Publications	15 words — < 1%

25 Kazuhiro Yasufuku, Jocelyne Fleury Feith. 15 words — < 1 %
"Cytological specimens obtained by
endobronchial ultrasound-guided transbronchial needle
aspiration: Sample handling and role of rapid on-site
evaluation", Annales de Pathologie, 2012

Crossref

26 Kuruswamy Thurai Prasad, Valliappan Muthu, 15 words — < 1 %
Inderpaul Singh Sehgal, Sahajal Dhooria et al.
"Endosonographic characteristics of mediastinal lymph nodes
for predicting malignancy in high tuberculosis burden settings:
a study of 774 subjects", Expert Review of Respiratory Medicine,
2022

Crossref

27 "Abstracts", Journal of Thoracic Oncology, 2011 14 words — < 1 %

Crossref

28 "EANM'15", European Journal of Nuclear Medicine 14 words — < 1 %
and Molecular Imaging, 2015

Crossref

29 www.dovepress.com 13 words — < 1 %

Internet

30 www.mdpi.com 13 words — < 1 %

Internet

31 M. Talaeepour, F. Moattar ., F. Atabi ., S. Borhan 12 words — < 1 %
Azad ., A.R. Talaeepour .. "Investigation on Radon
Concentration in the Tehran Subway Stations, in Regard with
Environmental Effects", Journal of Applied Sciences, 2006

Crossref

32 A. Christine Argento, Daniel L. Gilstrap, Scott 10 words — < 1 %
Shofer, Kamran Mahmood, Kimberly Blackwell,
Momen M. Wahidi. "Endobronchial Ultrasound-guided

Transbronchial Needle Aspiration in the Diagnosis of Breast Cancer Thoracic Metastases and Detection of Receptor Discordance", Journal of Bronchology & Interventional Pulmonology, 2018

Crossref

33 R. C. Rintoul. "Endobronchial and endoscopic ultrasound-guided real-time fine-needle aspiration for mediastinal staging", European Respiratory Journal, 2005

9 words — < 1%

Crossref

34 www.researchsquare.com

Internet

9 words — < 1%

35 www.frontiersin.org

Internet

8 words — < 1%

36 www.thieme-connect.com

Internet

8 words — < 1%

37 Faeq R. Kukhon, Xinyue Lan, Scott A. Helgeson, Vichaya Arunthari, Sebastian Fernandez - Bussy, Neal M. Patel. "Occult Lymph Node Metastasis in Radiologic Stage I Non - small Cell Lung Cancer: The Role of Endobronchial Ultrasound", The Clinical Respiratory Journal, 2021

7 words — < 1%

Crossref

38 Navani, N., D. R. Lawrence, S. Kolvekar, M. Hayward, D. McAsey, G. Kocjan, M. Falzon, A. Capitanio, P. Shaw, S. Morris, R. Z. Omar, and S. M. Janes. "EBUS-TBNA Prevents Mediastinocopies in the Diagnosis of Isolated Mediastinal Lymphadenopathy: A Prospective Trial", American Journal of Respiratory and Critical Care Medicine, 2012.

7 words — < 1%

Crossref

39 Sanz-Santos, Jose, Beatriz Cirauqui, Estefania Sanchez, Felipe Andreo, Pere Serra, Eduard Monso, Eva Castellà, Mariona Llatjós, Miguel Mesa, Juan Ruiz-Manzano, and Rafael Rosell. "Endobronchial ultrasound-guided transbronchial needle aspiration in the diagnosis of intrathoracic lymph node metastases from extrathoracic malignancies", Clinical & Experimental Metastasis, 2013. 7 words — < 1%
Crossref

40 T. Fujiwara. "The Utility of Sonographic Features During Endobronchial Ultrasound-Guided Transbronchial Needle Aspiration for Lymph Node Staging in Patients With Lung Cancer: A Standard Endobronchial Ultrasound Image Classification System", Chest, 09/01/2010 7 words — < 1%
Crossref

41 "Flexible Bronchoscopy", Wiley, 2020 6 words — < 1%
Crossref

42 &NA;, . "15th World Conference on Lung Cancer :", Journal of Thoracic Oncology, 2013. 6 words — < 1%
Crossref

43 Kazuhiro Yasufuku. "EBUS-TBNA Bronchoscopy", Endobronchial Ultrasound, 2009 6 words — < 1%
Crossref

EXCLUDE QUOTES OFF

EXCLUDE SOURCES OFF

EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE MATCHES OFF