

World Journal of *Psychiatry*

World J Psychiatr 2018 September 20; 8(3): 75-107



**EDITORIAL**

- 75 Successful treatment of nightmares may reduce psychotic symptoms in schizophrenia
Seeman MV
- 79 Psycho-emotional content of illness narrative master plots for people with chronic illness: Implications for assessment
Soundy A

MINIREVIEWS

- 83 Single men seeking adoption
Seeman MV

ORIGINAL ARTICLE**Observational Study**

- 88 Drinking plain water is associated with decreased risk of depression and anxiety in adults: Results from a large cross-sectional study
Haghighatdoost F, Feizi A, Esmailzadeh A, Rashidi-Pourfard N, Keshteli AH, Roohafza H, Adibi P

SYSTEMATIC REVIEW

- 97 Antidepressant foods: An evidence-based nutrient profiling system for depression
LaChance LR, Ramsey D

LETTERS TO THE EDITOR

- 105 Psychic euosmia and obsessive compulsive personality disorder
Pasquini M, Maraone A, Roselli V, Tarsitani L

ABOUT COVER

Editorial Board Member of *World Journal of Psychiatry*, Rami Bou Khalil, MD, MSc, Attending Doctor, Lecturer, Department of Psychiatry, Saint Joseph University, Beirut, Lebanon

AIM AND SCOPE

World Journal of Psychiatry (*World J Psychiatr*, *WJP*, online ISSN 2220-3206, DOI: 10.5498) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJP covers topics concerning behavior and behavior mechanisms, psychological phenomena and processes, mental disorders, behavioral disciplines and activities, adjustment disorders, anxiety disorders, delirium, dementia, amnesic disorders, cognitive disorders, dissociative disorders, eating disorders, factitious disorders, impulse control disorders, mental disorders diagnosed in childhood, mood disorders, neurotic disorders, personality disorders, schizophrenia and disorders with psychotic features, sexual and gender disorders, sleep disorders, somatoform disorders, and substance-related disorders. Priority publication will be given to articles concerning diagnosis and treatment of psychiatric diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, differential diagnosis, imaging tests, pathological diagnosis, molecular biological diagnosis, immunological diagnosis, genetic diagnosis, functional diagnostics, and physical diagnosis; and comprehensive therapy, drug therapy, surgical therapy, interventional treatment, minimally invasive therapy, and robot-assisted therapy.

We encourage authors to submit their manuscripts to *WJP*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

World Journal of Psychiatry is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), China National Knowledge Infrastructure (CNKI), and Superstar Journals Database.

EDITORS FOR
THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Ying-Na Bian*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Ying Dou*
Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL
World Journal of Psychiatry

ISSN
ISSN 2220-3206 (online)

LAUNCH DATE
December 31, 2011

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/2220-3206/editorialboard.htm>

EDITORIAL OFFICE
Jin-Lei Wang, Director
World Journal of Psychiatry
Baishideng Publishing Group Inc
8226 Regency Drive, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242

Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
8226 Regency Drive,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
September 20, 2018

COPYRIGHT
© 2018 Baishideng Publishing Group Inc. Articles pub-

lished by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f6publishing.com>

Observational Study

Drinking plain water is associated with decreased risk of depression and anxiety in adults: Results from a large cross-sectional study

Fahimeh Haghighatdoost, Awat Feizi, Ahmad Esmailzadeh, Nafiseh Rashidi-Pourfard, Ammar Hassanzadeh Keshteli, Hamid Roohafza, Payman Adibi

Fahimeh Haghighatdoost, Food Security Research Center and Department of Community Nutrition, School of Nutrition and Food Science, Isfahan University of Medical Sciences, Isfahan 81746-73461, Iran

Awat Feizi, Psychosomatic Research Center, Integrative Functional Gastrointestinal Research Center and Biostatistics and Epidemiology Department, School of Health Isfahan University of Medical Sciences, Hezarjarib, Isfahan 81746-73461, Iran

Ahmad Esmailzadeh, Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran 14176-53761, Iran

Nafiseh Rashidi-Pourfard, Shahid Motahari Hospital, Foolad-shahr, Isfahan 81746-73461, Iran

Ammar Hassanzadeh Keshteli, Department of Medicine, Faculty of Medicine & Dentistry, University of Alberta, Edmonton, AL 54321, Canada

Hamid Roohafza, Cardiac Rehabilitation Research Center, Isfahan Cardiovascular Research Institute Isfahan University of Medical Sciences, Isfahan 81746-73461, Iran

Payman Adibi, Integrative Functional Gastrointestinal Research Center and Gastroenterology Section, Department of Internal Medicine, Isfahan University of Medical Sciences, Isfahan 81746-73461, Iran

ORCID number: Fahimeh Haghighatdoost (0000-0003-4766-6267); Awat Feizi (0000-0002-1930-0340); Ahmad Esmailzadeh (0000-0002-8735-6047); Nafiseh Rashidi-Pourfard (0000-0003-0800-9408); Ammar Hassanzadeh Keshteli (0000-0001-7375-6210); Hamid Roohafza (0000-0003-3582-0431); Payman Adibi (0000-0001-6411-5235).

Author contributions: Haghighatdoost F and Feizi A contributed to statistical analysis, data interpretation and manuscript drafting; Esmailzadeh A contributed to SEPAHAN study concepts and

design, data collection and drafting of the manuscript; Rashidi-Pourfard N contributed to statistical analysis, data interpretation and manuscript drafting; Keshteli AH, Roohafza H and Adibi P contributed to SEPAHAN study concepts and design, data collection and drafting of the manuscript; Feizi A supervised the current secondary study; all authors approved the final version of the manuscript.

Institutional review board statement: The study protocol was approved by the bioethics committee of Isfahan University of Medical Sciences (#189069, #189082, #189086).

Informed consent statement: All study participants provided an informed consent statement.

Conflict-of-interest statement: None of the authors had any personal or financial conflicts of interest.

STROBE statement: The guidelines of the STROBE statement have been adopted.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Awat Feizi, PhD, Full Professor, Psychosomatic Research Center, Integrative Functional Gastrointestinal Research Center and Biostatistics and Epidemiology Department, School of Health Isfahan University of Medical Sciences, Hezarjarib, Isfahan 81746-73461, Iran. awat_feizi@hlth.mui.ac.ir
Telephone: +98-313-792 3250

Fax: +98-313-7923232

Received: February 14, 2018

Peer-review started: February 14, 2018

First decision: March 2, 2018

Revised: April 23, 2018

Accepted: June 9, 2018

Article in press: June 9, 2018

Published online: September 20, 2018

Abstract

AIM

To investigate the relation between plain water drinking and risk of depression and anxiety among a large sample of Iranian adults.

METHODS

A total of 3327 Iranian general adults were included in this cross-sectional study. Validated Iranian version of the Hospital Anxiety and Depression Scale was used to assess anxiety and depression. Water consumption was assessed by asking about the number of glasses of water that consumed daily. Water consumption was categorized into < 2, 2-5, and ≥ 5 glasses of water/d.

RESULTS

In the crude model, the lowest level of water drinking (< 2 glasses/d) compared with reference group (≥ 5 glasses/d) doubled the risk of depression and anxiety ($P < 0.0001$). After adjusting potential confounders, this inverse link remained significant for depression (OR: 1.79; 95%CI: 1.32, 2.42; $P < 0.0001$), but not for anxiety (OR: 1.49; 95%CI: 0.98, 2.25; $P = 0.109$). In stratified analyses by sex, after controlling for potential confounders, water drinking < 2 glasses/d was associated with 73% and 54% increment in the risk of depression in men and women, respectively ($P < 0.05$), whilst no significant association was observed for anxiety either in men or in women.

CONCLUSION

We found inverse associations between plain water consumption and depression. Also, these findings showed a tended risky association, but not statistically significant, between lower levels of water consumption and anxiety. These findings warrant evaluation in prospective and clinical trials studies to establish the plausible role of water in mental health status.

Key words: Water; Anxiety; Depression; Psychological disorders; Iranian

© The Author(s) 2018. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Evidence supports the relation between water consumption and health outcomes. Validated Iranian version of the Hospital Anxiety and Depression Scale was used to assess anxiety and depression.

Water consumption was assessed by asking about the number of glasses of water that consumed daily. After adjusting potential confounders, an inverse link was observed between water and depression, but not for anxiety, though in the crude model both disorders were inversely related to water drinking. These findings warrant evaluation in prospective studies to establish the plausible role of water in mental health status.

Haghighatdoost F, Feizi A, Esmaillzadeh A, Rashidi-Pourfard N, Keshteli AH, Roohafza H, Adibi P. Drinking plain water is associated with decreased risk of depression and anxiety in adults: Results from a large cross-sectional study. *World J Psychiatr* 2018; 8(3): 88-96 Available from: URL: <http://www.wjgnet.com/2220-3206/full/v8/i3/88.htm> DOI: <http://dx.doi.org/10.5498/wjp.v8.i3.88>

INTRODUCTION

The prevalence of common psychological disorders has been increasing over recent decades^[1,2]. Approximately, 29.2% of people are suffering from one of the common mental disorders worldwide^[3]. Brain disorders, including both mental and neurologic disorders account for one-third of the economic cost of all diseases^[4].

Drinking plenty of water is publicly believed to be useful for health and has been recommended in various dietary guidelines. Findings from a systematic review revealed that increased water consumption had a weight-reduction effect^[5], and it has widely been known as an approach in weight-loss programs^[6]. In addition, several studies have reported a bidirectional link between excess body weight^[7,8] or diabetes^[9] and mental disorders. It has been indicated that pathogenic substrates are the same in both metabolic and brain disorders; and therefore, the term of "metabolic-mood syndrome" has been suggested^[10]. Lifestyle changes may to some extent explain the concurrency of obesity and mental disorders. Replacing water with sugar sweetened beverages might be the reason for inverse link between water consumption and obesity^[11,12]. There is also evidence indicating the higher risk for mental disorders especially depression with higher consumption of sugar sweetened beverages^[13], as well as higher glycaemic index diets^[14,15].

Overall, accumulating evidence suggests that the beneficial relation between water ingestion and mental disorders is possible. However, the knowledge regarding beneficial effects of water consumption in mental health is rare. Water facilitates signaling pathway and nutrients delivery to the brain, removes toxins and inflammatory markers and provides energy sources for brain, and thereby improves brain function. Although several studies have shown that water insecurity is associated with psychological distress and anxiety^[16,17], to the best of our knowledge, there is no study assessing the association between the water quantity and common

mental disorders including depression and anxiety. In the current study, therefore, we aimed to evaluate whether decreased plain water intake is associated with higher risk of mental disorders and whether it is gender-specific.

MATERIALS AND METHODS

Subjects

This cross-sectional study was carried out within the framework of the cross-sectional study on the Epidemiology of psychological, Alimentary health and Nutrition (SEPAHAN) project. The main aim of this project was to investigate the relationship between functional gastrointestinal disorders and lifestyle, nutritional factors and psychological profiles. The details of the project have been provided in Alibi *et al.*^[18] (2012). Briefly, the participants of the study were selected using multistage cluster and convenience sampling method amongst non-academic Isfahanian adults working at Isfahan University of Medical Sciences (IUMS) in 20 cities across Isfahan province. Each city was considered as a first stage's cluster and health centers affiliated to IUMS in each city were considered as second stage clusters and among them, considering the full coverage of geographic regions, some centers were selected randomly and those participants who were willing to participate in SEPAHAN study were recruited. In order to increase the participation rate and the data collection accuracy, the project was conducted in two phases. In the first phase, 10087 questionnaires were distributed to the participants, collecting information about anthropometric measures, demographic profile, lifestyle, nutritional factors and physical activity. 8691 completed questionnaires were returned in the first phase (response rate: 86.16%). In the second phase, psychological data were collected (response rate: 64.6%). Finally, after matching returned questionnaires in phase 2 with their equivalents in phase 1, we reached 4763 questionnaires. Of these, 2904 persons had complete information about both water consumption and psychological profile which were included in the statistical analysis. The study protocol was reviewed and approved by the ethical committee of Isfahan University of Medical Sciences.

Water consumption was assessed by asking about the average number of glasses of plain water that usually consumed in a day by each participant. The possible items to answer were < 2 glasses/d, 2-5 glasses/d and ≥ 5 glasses/d.

Mental disorders assessment

A validated Iranian version of the Hospital Anxiety and Depression scale (HADS) was used to assess anxiety and depression^[19]. HADS is a simple psychological questionnaire includes two separate parts to screen the severity of anxiety and depression. Each part includes 7 questions with a four-point rating scale (0-3); higher scores indicate greater degree of anxiety or depression.

For both disorders, the score range is from 0 to 21. To identify the presence of either disorder, score 8 was considered as the cut point. Therefore, score 8 or greater were considered to have depression or anxiety and scores ≤ 7 were considered normal^[20].

Covariates assessment

Dietary intakes of participants were assessed using a valid and reliable 106-item dish-based food frequency questionnaire^[21]. Body mass index (BMI) was calculated as weight (kg) divided by height squared (m^2). The current level of participants' physical activity was assessed using General Practice Physical Activity Questionnaire^[22] and participants were categorized as physically moderately active, active, moderately inactive and inactive. Because of close relation between gastrointestinal disorders and psychological health, we considered functional gastrointestinal disorders (FGID) as an important covariate in our analysis. Suffering from gastrointestinal disorders was assessed using a valid and modified Iranian version of ROME III questionnaire^[23]. FGID was defined as suffering from at least one of the following main gastrointestinal disorders: gastroesophageal reflux, dyspepsia, irritable bowel syndrome and constipation.

Statistical analysis

General characteristics in categories of plain water intake were reported as means and standard error (SE) or percentage for continuous variables and categorical variables, respectively. To examine the differences across plain water categories, analysis of variance (ANOVA) for continuous variables and χ^2 test for categorical variables were used. All dietary intakes were adjusted for age (yr) and weight (kg) by analysis of covariance (ANCOVA). Multiple logistic regression was used to estimate odds ratios (OR) (95%CI) for the presence of depression and anxiety across categories of plain water intake in crude and multivariable-adjusted models. In adjusted models, we controlled confounding impacts of age, sex, marital status (married, single, divorced, widowed), educational level (less than 12 years, 12-16 years and more than 16 years), BMI (continuous), smoking (non- and ex-smokers vs. current smokers), physical activity (moderately active and active vs. moderately inactive and inactive), FGIDs (Yes/No), intake of anti-psychotic medicines (Yes/No), and dietary intakes which play role in mental health [magnesium, riboflavin, pyridoxine, folate, cobalamin, Docosahexaenoic acid (DHA), Eicosapentaenoic acid (EPA), energy, fibre and caffeine]. All confounders, including energy intake, were included in the statistical analysis as covariates. We performed stratified analyses, applying the above-mentioned models, by sex to evaluate potential modifying effect of sex related to mental health status. The adjusted models were also controlled for the same potential confounders, mentioned above. In all logistic regression analyses, *P* for linear trends was determined by Mantel-Haenszel

Table 1 General characteristic of participants across categories of plain water consumption

Variables	Plain water drinking			P value ¹
	< 2 glasses/d	2-5 glasses/d	≥ 5 glasses/d	
Participants (n)	854	1764	709	
Age (yr) ²	37.2 ± 0.3	36.1 ± 0.2	34.8 ± 0.3	< 0.0001
BMI (kg/m ²) ²	24.7 ± 0.1	24.7 ± 0.10	25.5 ± 0.2	< 0.0001
Anxiety score ²	4.0 ± 0.1	3.5 ± 0.10	3.0 ± 0.1	< 0.0001
Depression score ² n (%)	6.8 ± 0.1	6.1 ± 0.09	5.3 ± 0.1	< 0.0001
Anxious	136 (16.2)	244 (14.1)	61 (8.7)	< 0.0001
Depressed	303 (36.0)	478 (27.7)	151 (21.6)	< 0.0001
Male	242 (28.3)	757 (42.9)	387 (54.6)	< 0.0001
Marital status				0.035
Married	680 (81.4)	1431 (82.9)	549 (78.8)	
Single	135 (16.2)	271 (15.7)	139 (19.9)	
Other	20 (2.4)	25 (1.4)	9 (1.3)	
Anti-psychotic medicines use	56 (6.6)	91 (5.2)	39 (5.5)	0.342
Current smokers	147 (17.2)	243 (13.8)	70 (9.9)	< 0.0001
Moderately active and active	303 (39.6)	729 (45.7)	350 (56.1)	< 0.0001
Educational level				0.005
≤ 12 yr	333 (39.8)	633 (36.9)	301 (43.1)	
12-16 yr	438 (52.4)	956 (55.6)	347 (49.7)	
≥ 16 yr	65 (7.8)	129 (7.5)	50 (7.2)	
FGID ³ (%)	488 (57.1)	892 (50.6)	321 (45.3)	< 0.0001
Nutrients ⁴				
Energy (kcal/d)	2306.1 ± 30.1	2385.8 ± 20.7	2487.9 ± 34.7	0.001
Protein (% of total calorie)	84.3 ± 1.2	88.5 ± 0.8	93.2 ± 1.4	< 0.0001
Fat (% of total calorie)	95.5 ± 1.3	98.7 ± 0.9	103.1 ± 1.5	0.001
Carbohydrate (g/d)	285.3 ± 4.2	294.5 ± 2.9	306.3 ± 4.9	0.006
Fibre (g/d)	22.0 ± 0.2	22.5 ± 0.1	23.0 ± 0.2	0.003
Caffeine (mg/d)	106.7 ± 3.1	96.7 ± 2.2	96.0 ± 3.7	0.022
Magnesium (mg/d)	314.1 ± 4.3	327.2 ± 3.0	349.6 ± 5.0	< 0.0001
Thiamin (mg/d)	1.8 ± 0.03	1.8 ± 0.02	1.9 ± 0.03	0.062
Riboflavin (mg/d)	1.8 ± 0.03	1.9 ± 0.02	2.0 ± 0.03	< 0.0001
Pyridoxine (mg/d)	1.9 ± 0.03	2.0 ± 0.02	2.1 ± 0.03	< 0.0001
Folate (mg/d)	562.4 ± 5.8	573.8 ± 4.0	600.2 ± 6.7	0.801
Cobalamine (mg/d)	2.8 ± 0.05	3.0 ± 0.03	3.1 ± 0.05	< 0.0001
DHA (g/d) ³	0.2 ± 0.01	0.2 ± 0.01	0.2 ± 0.01	< 0.0001
EPA (g/d) ³	0.06 ± 0.003	0.1 ± 0.002	0.1 ± 0.003	< 0.0001
Food groups				
Fruits (g/d)	285.8 ± 8.7	322.4 ± 6.0	351.1 ± 10.1	< 0.0001
Vegetables (g/d)	215.4 ± 4.8	241.4 ± 3.3	264.7 ± 5.5	< 0.0001
Nuts, legumes and soy (g/d)	55.5 ± 1.5	56.8 ± 1.0	60.9 ± 1.7	0.047
White meat (g/d)	59.4 ± 1.8	64.03 ± 1.2	67.5 ± 2.02	0.009
Red meat (g/d)	75.4 ± 1.8	78.4 ± 1.2	84.5 ± 2.06	0.004
Refined grains (g/d)	401.6 ± 7.1	390.7 ± 5.4	389.4 ± 9.11	0.473
Whole grains (g/d)	39.5 ± 2.9	40.8 ± 1.1	52.1 ± 3.40	0.007

¹Derived from one way ANOVA and chi-square test for continuous and categorical variables, respectively and analysis of covariance (ANCOVA) for nutrients and food groups; ²Values are means ± SEs; ³FGID defined as suffering from at least one of the following gastrointestinal disorders: Gastroesophageal Reflux, dyspepsia, irritable bowel syndrome and constipation; ⁴The nutrients were adjusted for age and body weight. DHA: Docosahexaenoic acid; EPA: Eicosa pentaenoic acid.

extension chi-square test. All statistical analyses were done using Statistical Package for Social Sciences (SPSS, Inc., Chicago IL, United States; version 20). $P < 0.05$ was considered significant in all statistical analyses.

RESULTS

In terms of water intake, participants were categorized into three major classes (< 2 glasses, 2-5 glasses, ≥ 5 glasses/d). Those consumed more water had higher BMI ($P < 0.0001$) and were more probably to be younger, male, physically active (all $P < 0.0001$) and single ($P < 0.05$), but less educated compared with those

who consumed less water ($P = 0.005$). Greater plain water drinking was associated with lower prevalence of smoking, anxiety, depression ($P < 0.0001$). Consistently, those consumed greater plain water had lower anxiety and depression score compared with those in the lowest category of plain water intake (< 2 glasses/d) ($P < 0.0001$). Water consumption was strongly associated with increased risk of FGID ($P < 0.0001$) (Table 1). Greater plain water consumption was associated with higher intakes of energy, protein, fat, carbohydrate, fibre, magnesium, riboflavin, pyridoxine, cobalamin, docosahexaenoic acid (DHA), eicosa pentaenoic acid (EPA), fruits, vegetables and red meat (P for all <

Table 2 General characteristics of participants based on categories of anxiety and depression *n* (%)

	Depression		<i>P</i> value ¹	Anxiety		<i>P</i> value ¹
	Yes (28%)	No (72%)		Yes (13.1%)	No (86.9%)	
Age (yr) ²	36.3 ± 0.3	36.1 ± 0.2	0.562	35.4 ± 0.2	36.3 ± 0.3	0.020
BMI (kg/m ²) ²	24.8 ± 0.1	24.9 ± 0.1	0.712	25.0 ± 0.2	24.9 ± 0.1	0.636
Male	285 (30.2)	1082 (46.0)	< 0.0001	117 (26.1)	1251 (43.9)	< 0.0001
Anti-psychotic medicines	106 (11.2)	76 (3.2)	< 0.0001	70 (15.6)	113 (4.0)	< 0.0001
Current smokers	164 (17.4)	290 (12.3)	< 0.0001	84 (18.8)	370 (13.0)	0.001
Moderately active and active	338 (39.7)	1028 (48.8)	< 0.0001	155 (38.5)	1212 (47.4)	0.001
Educational level			< 0.0001			< 0.0001
< 12 yr	153 (16.7)	236 (10.2)		84 (19.3)	306 (11.0)	
12-16 yr	715 (78.1)	1876 (81.4)		337 (77.5)	2255 (80.9)	
≥ 16 yr	48 (5.2)	194 (8.4)		14 (3.2)	228 (8.2)	
Marital status			0.001			0.001
Married	733 (79.4)	1901 (82.4)		361 (83.2)	2274 (81.3)	
Single	163 (17.7)	380 (16.5)		58 (13.4)	486 (17.4)	
Other	27 (2.9)	26 (1.1)		15 (3.5)	38 (1.4)	
FGID ³	665 (70.5)	1022 (43.4)	< 0.0001	357 (79.7)	1329 (46.6)	< 0.0001
Water drinking			< 0.0001			< 0.0001
< 2 glasses/d	303 (32.5)	539 (23.1)		136 (30.8)	706 (25.0)	
2-5 glasses/d	478 (51.3)	1246 (53.4)		244 (55.3)	1481 (52.4)	
≥ 5 glasses/d	151 (16.2)	548 (23.5)		61 (13.8)	639 (22.6)	

¹Derived from independent t-test and chi-square test for continuous and categorical variables, respectively; ²Values are means ± SEs or percent; ³FGID defined as suffering from at least one of the following gastrointestinal disorders: Gastroesophageal Reflux, dyspepsia, irritable bowel syndrome and constipation.

0.01). Tiamin, folate, white meat, refined grains, whole grains, nuts, legumes and soy consumptions were not significantly different across the categories of plain water intake.

Table 2 presents the general characteristics of study population stratified by the status of anxiety or depression. Individuals, who were anxious, but not depressed, were younger than healthy subjects. The prevalence of both depression and anxiety was higher among women, anti-psychotic medicines users, smokers and individuals who suffered from FGID. Conversely, healthy subjects were more probably to be physically active or moderately active, have higher education levels and drink more glasses of water. Whilst depression was less prevalent among married individuals, anxiety was more prevalent.

Crude and multivariable-adjusted ORs (95%CI) of depression and anxiety across the categories of plain water intake are illustrated in Table 3. Compared with the reference group (≥ 5 glasses/d), lower levels of water consumption (< 2 glasses/d) was associated with a greater chance of having depression (OR: 2.04; 95%CI: 1.62, 2.56; *P* < 0.0001) and anxiety (OR: 2.02; 95%CI: 1.46, 2.78; *P* < 0.0001) in crude model. Adjustment for multiple potential confounders slightly weakened these associations, but remained strongly significant for depression (*P* < 0.0001). However, after adjustment for dietary intakes, the lower levels of plain water consumption was a risk factor for anxiety however the significant link disappeared (for < 2 glasses/d: OR: 1.49; 95%CI: 0.98, 2.25, and for 2-5 glasses/d: OR: 1.58, 95%CI: 1.08, 2.30; *P* = 0.109; vs ≥ 5 glasses/d).

In all crude and adjusted models, lower levels of water consumption (< 2 glasses/d) compared with the

reference group (≥ 5 glasses/d) was associated with a greater chance of having depression either in men or in women not only in crude but also in fully adjusted models (2-5 glasses/d: OR: 1.54; 95%CI: 1.00, 2.36 and < 2 glasses/d: OR: 1.73; 95%CI: 1.02, 2.93; *P* = 0.04) for men and in women (2-5 glasses/d: OR: 1.18 ; 95%CI: 0.86, 1.63 and < 2 glasses/d: OR: 1.54; 95%CI: 1.09, 2.15; *P* = 0.007). In the crude model drinking lower levels of water was associated with increased risk of anxiety in men (2-5 glasses/d: OR: 2.01; 95%CI: 1.20, 3.34 and < 2 glasses/d: OR: 1.83; 95%CI: 0.97, 3.43; *P* = 0.041) and in women (2-5 glasses/d: OR: 1.42 ; 95%CI: 0.98, 2.05 and < 2 glasses/d: OR: 1.59; 95%CI: 1.08, 2.34; *P* = 0.026); nevertheless, after adjustment for various confounders the significance associations disappeared in both genders.

DISCUSSION

In this analysis of a large cross-sectional study of general adults, lower daily plain water intake was associated with increased risk of depression and anxiety in the crude model. Although controlling for potential confounders attenuated these associations, the inverse link for depression remained strongly significant, whilst anxiety risk tended to be higher for lower water intake in the final model.

To our knowledge, this was the first investigation to examine the association of plain water consumption and common mental disorders, although the linkage of water/fluid consumption with obesity^[6,12,24], coronary diseases^[25], hyperglycaemia or diabetes^[26,27], cancer^[28] and mortality^[29,30] has been examined in earlier studies. Some of these studies^[6,12,24,25], but not all^[26-30], confirm

Table 3 Multivariable-adjusted ORs (and 95%CI) for depression and anxiety across categories of plain water consumption in the whole population and stratified by sex

	Plain water drinking			<i>P</i> trend ¹
	< 2 glasses/d	2-5 glasses/d	≥ 5 glasses/d	
Depression				
Crude model	2.04 (1.62, 2.56)	1.39 (1.13, 1.71)	1 (reference)	< 0.0001
Model 1 ²	1.75 (1.35, 2.26)	1.30 (1.02, 1.64)	1 (reference)	< 0.0001
Model 2 ²	1.84 (1.37, 2.45)	1.39 (1.07, 1.80)	1 (reference)	< 0.0001
Model 3 ²	1.84 (1.40, 2.49)	1.41 (1.07, 1.84)	1 (reference)	< 0.0001
Model 4 ²	1.79 (1.32, 2.42)	1.37 (1.04, 1.80)	1 (reference)	< 0.0001
Anxiety				
Crude model	2.02 (1.46, 2.78)	1.73 (1.29, 2.32)	1 (reference)	< 0.0001
Model 1 ²	1.53 (1.07, 2.17)	1.44 (1.04, 1.99)	1 (reference)	0.026
Model 2 ²	1.58 (1.06, 2.36)	1.60 (1.11, 2.30)	1 (reference)	0.029
Model 3 ²	1.56 (1.03, 2.35)	1.63 (1.12, 2.40)	1 (reference)	0.066
Model 4 ²	1.49 (0.98, 2.25)	1.58 (1.08, 2.30)	1 (reference)	0.109
Men (<i>n</i> = 1386)				
Depression				
Crude model	1.77 (1.19, 2.63)	1.37 (0.99, 1.90)	1 (reference)	0.004
Model 1 ²	2.00 (1.26, 3.20)	1.45 (0.98, 2.13)	1 (reference)	0.003
Model 2 ²	1.95 (1.18, 3.21)	1.59 (1.06, 2.38)	1 (reference)	0.008
Model 3 ²	1.73 (1.02, 2.92)	1.54 (1.01, 2.36)	1 (reference)	0.037
Model 4 ²	1.73 (1.02, 2.93)	1.54 (1.00, 2.36)	1 (reference)	0.040
Anxiety				
Crude model	1.83 (0.97, 3.43)	2.01 (1.20, 3.34)	1 (reference)	0.041
Model 1 ²	2.36 (1.17, 4.74)	1.55 (0.87, 2.78)	1 (reference)	0.016
Model 2 ²	2.15 (1.02, 4.54)	1.59 (0.86, 2.95)	1 (reference)	0.042
Model 3 ²	1.76 (0.81, 3.84)	1.55 (0.82, 2.94)	1 (reference)	0.147
Model 4 ²	1.74 (0.80, 3.80)	1.52 (0.80, 2.88)	1 (reference)	0.161
Women (<i>n</i> = 1941)				
Depression				
Crude model	1.75 (1.30, 2.35)	1.26 (0.95, 1.67)	1 (reference)	<0.0001
Model 1 ²	1.61 (1.17, 2.21)	1.21 (0.90, 1.63)	1 (reference)	0.001
Model 2 ²	1.61 (1.16, 2.23)	1.21 (0.89, 1.65)	1 (reference)	0.002
Model 3 ²	1.61 (1.15, 2.26)	1.22 (0.89, 1.67)	1 (reference)	0.003
Model 4 ²	1.54 (1.09, 2.15)	1.18 (0.86, 1.63)	1 (reference)	0.007
Anxiety				
Crude model	1.59 (1.08, 2.34)	1.42 (0.98, 2.05)	1 (reference)	0.026
Model 1 ²	1.35 (0.89, 2.05)	1.41 (0.95, 2.07)	1 (reference)	0.264
Model 2 ²	1.39 (0.91, 2.13)	1.44 (0.97, 2.15)	1 (reference)	0.233
Model 3 ²	1.37 (0.88, 2.13)	1.44 (0.95, 2.18)	1 (reference)	0.288
Model 4 ²	1.30 (0.83, 2.02)	1.40 (0.92, 2.12)	1 (reference)	0.420

¹Derived from a Mantel-Haenszel extension chi-square test; ²Model 1: Adjusted for age, sex (in the whole population), marital status, educational level, model 2: Further adjustment for BMI, smoking, physical activity, model 3: Additional control for functional gastrointestinal disorders (FGID) and anti-psychotic medicines, model 4: Further control for magnesium, riboflavin, pyridoxine, folate, cobalamin, DHA and EPA, energy, fibre and caffeine. DHA: Docosahexaenoic acid; EPA: Eicosa pentaenoic acid.

the beneficial effects of water drinking in reducing diseases' risk. Therefore, due to bidirectional link between metabolic status and mental health^[10], it might be concluded that water consumption can affect mental disorders risk *via* affecting metabolic status.

The bidirectional link between obesity and mental disorders is based on some shared peripheral and central pathological pathways, as well as genetic and environmental risk factors^[10]. However, in line with some studies^[26,31-33], we found that higher water consumption was associated with greater BMI. Therefore, some other plausible mechanisms may explain this inverse link. Our results indicate that higher water intake is associated with various healthy behaviours such as being more physically active and consuming greater amounts of nutrients particularly those involved in the

nervous system (*e.g.*, riboflavin, magnesium, pyridoxine and cobalamin), which is consistent with previous studies^[31,34]. Beneficial effects of these nutrients in neurotransmitters synthesis and transportation, as well as activity of many enzymes in the nervous system have been shown earlier^[35,36]. Nevertheless, even after controlling for dietary intakes, depression was significantly related to water consumption, and anxiety tended to be higher in those who drank less water. A possible reason for the inverse link between water consumption and the risk of depression might be the decreased activity of the sympathetic nervous system by drinking water which reduces plasma levels of norepinephrine^[37]. Elevated level of norepinephrine is a characteristic of psychosomatic depression^[38] which may induce noradrenergic-vasopressinergic activation,

and consequently the activation of the hypothalamic-pituitary-adrenal (HPA) axis. Increased vasopressinergic activation of the HPA axis has been suggested as a plausible mechanism in all depressive disorders^[38], which can be involved in mental symptoms production. In spite of similar link between water drinking and depression risk in men and women, the linkage for anxiety was stronger in men than women, though adjustment for antipsychotic medicines and FGIDs eliminated it. This difference between men and women might be related to sex-differences in the use of coping strategies^[39]. However, the reasons for this difference are not clear and require further investigation.

In the current analysis, water consumption was evaluated using a direct question regarding the average amount of daily consumption of water, and data regarding other beverages intake and moisture contents of food were not considered. In addition, we provided predefined categories of plain water intake for participants. These factors may lead to misclassification of participants and decrease the reliability of our findings. However, it should be taken into account that noncalorically sweetened beverages are not public among Iranians and they are not used generally. Moreover, tea and coffee are mostly consumed with sugar. Therefore, since all other beverages, which consumed by this population, had calorie and we controlled the confounding effect of energy in our analysis, it could be concluded that our findings have enough precise. The consistence between our study's results and available evidence regarding the health beneficial of water intake might be further reason to confirm the accuracy of our findings.

The limitations of the current study are using self-administered questionnaires for evaluating the study variables. However, the validity of all used instruments has been approved among Iranians. Although using self-reported tools of habitual fluid/water intake is a common limitation in this area of research, the validity of self-reported measures has been approved in earlier studies^[40,41]. In addition, no ideal method has been determined to assess beverage consumption and applying 24-h recall or frequency questionnaire beside a computer program on food composition is the existing recommend method in this regard^[42]. The cross-sectional design is another limitation that does not allow causal inferences. Indeed, it is possible that individuals with mental disorders tend to drink less water. Nevertheless, such changes would have weakened the associations identified. Therefore, the true estimates are probably even stronger than those we found. Moreover, since we studied the participants who are staffs of IUMS, although they were not academic and medical experts, this sample may not be representative of the entire general population and therefore the results of our study may not be generalizable to other individuals in our society. The strengths of this study are the large sample size and controlling various confounders' effects.

In conclusion, we found inverse associations between plain water consumption and common psycho-

logical disorders. After controlling for various potential confounders, this association was stronger for depression, whilst for anxiety tended to be higher. These findings warrant evaluation in prospective studies to establish the plausible role of water in the mental health status.

ARTICLE HIGHLIGHTS

Research background

Drinking a plenty of plain water has been known as a healthy behavior.

Research motivation

The association between drinking water and mental disorders has not been investigated.

Research objectives

We aimed to assess whether drinking water is associated with mental disorders, and also examine the sex-specific associations.

Research methods

Three thousand three hundred and twenty-seven adults in a cross-sectional study were categorized into three groups according the amount of water drinking (< 2, 2-5, and \geq 5 glasses of water/d). The risk of depression and anxiety was evaluated across different categories.

Research results

After controlling for various potential confounders, we found inverse associations between plain water consumption and depression, whilst for anxiety tended to be significant. These findings warrant evaluation in prospective studies to establish the plausible role of water in the mental health status.

Research conclusions

There were inverse associations between plain water consumption and common psychological disorders.

Research perspectives

Prospective studies to establish the plausible role of water consumption levels in the mental health status are needed.

REFERENCES

- 1 **Murphy JM**, Horton NJ, Laird NM, Monson RR, Sobol AM, Leighton AH. Anxiety and depression: a 40-year perspective on relationships regarding prevalence, distribution, and comorbidity. *Acta Psychiatr Scand* 2004; **109**: 355-375 [PMID: 15049772 DOI: 10.1111/j.1600-0447.2003.00286.x]
- 2 **Mojtabai R**. National trends in mental health disability, 1997-2009. *Am J Public Health* 2011; **101**: 2156-2163 [PMID: 21940913 DOI: 10.2105/AJPH.2011.300258]
- 3 **Berry CL**, Looker T, Germain J. The growth and development of the rat aorta. I. Morphological aspects. *J Anat* 1972; **113**: 1-16 [PMID: 4648481 DOI: 10.1093/ije/dyu038]
- 4 **Olesen J**, Leonardi M. The burden of brain diseases in Europe. *Eur J Neurol* 2003; **10**: 471-477 [PMID: 12940825 DOI: 10.1046/j.1468-1331.2003.00682.x]
- 5 **Muckelbauer R**, Sarganas G, Gruneis A, Müller-Nordhorn J. Association between water consumption and body weight outcomes: a systematic review. *Am J Clin Nutr* 2013; **98**: 282-299 [PMID: 23803882 DOI: 10.3945/ajcn.112.055061]
- 6 **Stookey JD**, Constant F, Popkin BM, Gardner CD. Drinking water is associated with weight loss in overweight dieting women independent of diet and activity. *Obesity* (Silver Spring) 2008; **16**: 2481-2488 [PMID: 18787524 DOI: 10.1038/oby.2008.409]
- 7 **Luppino FS**, de Wit LM, Bouvy PF, Stijnen T, Cuijpers P, Penninx BW, Zitman FG. Overweight, obesity, and depression: a systematic

- review and meta-analysis of longitudinal studies. *Arch Gen Psychiatry* 2010; **67**: 220-229 [PMID: 20194822 DOI: 10.1001/archgenpsychiatry.2010.2]
- 8 **McElroy SL**, Kotwal R, Malhotra S, Nelson EB, Keck PE, Nemeroff CB. Are mood disorders and obesity related? A review for the mental health professional. *J Clin Psychiatry* 2004; **65**: 634-651, quiz 730 [PMID: 15163249 DOI: 10.4088/JCP.v65n0507]
 - 9 **Hemmy Asamsama O**, Lee JW, Morton KR, Tonstad S. Bidirectional longitudinal study of type 2 diabetes and depression symptoms in black and white church going adults. *J Diabetes Metab Disord* 2015; **14**: 25 [PMID: 25897418 DOI: 10.1186/s40200-015-0150-5]
 - 10 **Mansur RB**, Brietzke E, McIntyre RS. Is there a "metabolic-mood syndrome"? A review of the relationship between obesity and mood disorders. *Neurosci Biobehav Rev* 2015; **52**: 89-104 [PMID: 25579847 DOI: 10.1016/j.neubiorev.2014.12.017]
 - 11 **Popkin BM**. Contemporary nutritional transition: determinants of diet and its impact on body composition. *Proc Nutr Soc* 2011; **70**: 82-91 [PMID: 21092363 DOI: 10.1017/S0029665110003903]
 - 12 **Fresán U**, Gea A, Bes-Rastrollo M, Ruiz-Canela M, Martínez-González MA. Substitution Models of Water for Other Beverages, and the Incidence of Obesity and Weight Gain in the SUN Cohort. *Nutrients* 2016; **8**: [PMID: 27809239 DOI: 10.3390/nu8110688]
 - 13 **Guo X**, Park Y, Freedman ND, Sinha R, Hollenbeck AR, Blair A, Chen H. Sweetened beverages, coffee, and tea and depression risk among older US adults. *PLoS One* 2014; **9**: e94715 [PMID: 24743309 DOI: 10.1371/journal.pone.0094715]
 - 14 **Gangwisch JE**, Hale L, Garcia L, Malaspina D, Opler MG, Payne ME, Rossom RC, Lane D. High glycemic index diet as a risk factor for depression: analyses from the Women's Health Initiative. *Am J Clin Nutr* 2015; **102**: 454-463 [PMID: 26109579 DOI: 10.3945/ajcn.114.103846]
 - 15 **Haghighatdoost F**, Azadbakht L, Keshteli AH, Feinle-Bisset C, Daghighzadeh H, Afshar H, Feizi A, Esmailzadeh A, Adibi P. Glycemic index, glycemic load, and common psychological disorders. *Am J Clin Nutr* 2016; **103**: 201-209 [PMID: 26607943 DOI: 10.3945/ajcn.114.105445]
 - 16 **Stevenson EG**, Ambelu A, Caruso BA, Tesfaye Y, Freeman MC. Community Water Improvement, Household Water Insecurity, and Women's Psychological Distress: An Intervention and Control Study in Ethiopia. *PLoS One* 2016; **11**: e0153432 [PMID: 27124391 DOI: 10.1371/journal.pone.0153432]
 - 17 **Aihara Y**, Shrestha S, Sharma J. Household water insecurity, depression and quality of life among postnatal women living in urban Nepal. *J Water Health* 2016; **14**: 317-324 [PMID: 27105416 DOI: 10.2166/wh.2015.166]
 - 18 **Adibi P**, Keshteli AH, Esmailzadeh A, Afshar H, Roohafza H, Bagherian-Sararoudi R, Daghighzadeh H, Soltanian N, Feinle-Bisset C, Boyce P, Talley N. The study on the epidemiology of psychological, alimentary health and nutrition (SEPAHAN): Overview of methodology. *J Res Med Sci* 2012; **17**: S292-S298
 - 19 **Montazeri A**, Vahdaninia M, Ebrahimi M, Jarvandi S. The Hospital Anxiety and Depression Scale (HADS): translation and validation study of the Iranian version. *Health Qual Life Outcomes* 2003; **1**: 14 [PMID: 12816545 DOI: 10.1186/1477-7525-1-14]
 - 20 **Montazeri A**, Harirchi AM, Shariati M, Garmaroudi G, Ebadi M, Fateh A. The 12-item General Health Questionnaire (GHQ-12): translation and validation study of the Iranian version. *Health Qual Life Outcomes* 2003; **1**: 66 [PMID: 14614778 DOI: 10.1186/1477-7525-1-66]
 - 21 **Keshteli A**, Esmailzadeh A, Rajaie S, Askari G, Feinle-Bisset C, Adibi P. A Dish-based Semi-quantitative Food Frequency Questionnaire for Assessment of Dietary Intakes in Epidemiologic Studies in Iran: Design and Development. *Int J Prev Med* 2014; **5**: 29-36 [PMID: 24554989]
 - 22 **NHS**. The General Practice Physical Activity Questionnaire (GPPAQ). 2008. Available from: URL: <http://www.enhertscg.nhs.uk/sites/default/files/pathways/General%20Practice%20Physical%20Activity%20Questionnaire%20%28GPPAQ%29.docx>
 - 23 **Gholamrezaei A**, Zolfaghari B, Farajzadegan Z, Nemati K, Daghighzadeh H, Tavakkoli H, Emami MH. Linguistic validation of the Irritable Bowel Syndrome-Quality of Life Questionnaire for Iranian patients. *Acta Med Iran* 2011; **49**: 390-395 [PMID: 21874644]
 - 24 **Vij VA**, Joshi AS. Effect of excessive water intake on body weight, body mass index, body fat, and appetite of overweight female participants. *J Nat Sci Biol Med* 2014; **5**: 340-344 [PMID: 25097411 DOI: 10.4103/0976-9668.136180]
 - 25 **Chan J**, Knutsen SF, Blix GG, Lee JW, Fraser GE. Water, other fluids, and fatal coronary heart disease: the Adventist Health Study. *Am J Epidemiol* 2002; **155**: 827-833 [PMID: 11978586 DOI: 10.1093/aje/155.9.827]
 - 26 **Pan A**, Malik VS, Schulze MB, Manson JE, Willett WC, Hu FB. Plain-water intake and risk of type 2 diabetes in young and middle-aged women. *Am J Clin Nutr* 2012; **95**: 1454-1460 [PMID: 22552035 DOI: 10.3945/ajcn.111.032698]
 - 27 **Roussel R**, Fezeu L, Bouby N, Balkau B, Lantieri O, Alhenc-Gelas F, Marre M, Bankir L, D.E.S.I.R. Study Group. Low water intake and risk for new-onset hyperglycemia. *Diabetes Care* 2011; **34**: 2551-2554 [PMID: 21994426 DOI: 10.2337/dc11-0652]
 - 28 **Di Maso M**, Bosetti C, Taborelli M, Montella M, Libra M, Zucchetto A, Turati F, Parpinel M, Negri E, Tavani A, Serraino D, Ferraroni M, La Vecchia C, Polesel J. Dietary water intake and bladder cancer risk: An Italian case-control study. *Cancer Epidemiol* 2016; **45**: 151-156 [PMID: 27821348 DOI: 10.1016/j.canep.2016.09.015]
 - 29 **Kant AK**, Graubard BI. A prospective study of water intake and subsequent risk of all-cause mortality in a national cohort. *Am J Clin Nutr* 2017; **105**: 212-220 [PMID: 27903521 DOI: 10.3945/ajcn.116.143826]
 - 30 **Palmer SC**, Wong G, Iff S, Yang J, Jayaswal V, Craig JC, Rochtchina E, Mitchell P, Wang JJ, Strippoli GF. Fluid intake and all-cause mortality, cardiovascular mortality and kidney function: a population-based longitudinal cohort study. *Nephrol Dial Transplant* 2014; **29**: 1377-1384 [PMID: 24398890 DOI: 10.1093/ndt/gft507]
 - 31 **Yang M**, Chun OK. Consumptions of plain water, moisture in foods and beverages, and total water in relation to dietary micronutrient intakes and serum nutrient profiles among US adults. *Public Health Nutr* 2015; **18**: 1180-1186 [PMID: 24507693 DOI: 10.1017/S136898001400007X]
 - 32 **Kant AK**, Graubard BI, Atchison EA. Intakes of plain water, moisture in foods and beverages, and total water in the adult US population--nutritional, meal pattern, and body weight correlates: National Health and Nutrition Examination Surveys 1999-2006. *Am J Clin Nutr* 2009; **90**: 655-663 [PMID: 19640962 DOI: 10.3945/ajcn.2009.27749]
 - 33 **Langdon SW**, Dennee-Sommers B. Exploring the relationships between self-objectification, rationales, and use of water as a strategy for appetite suppression. *Psychol Health Med* 2010; **15**: 17-25 [PMID: 20391221 DOI: 10.1080/13548500903431519]
 - 34 **Gazan R**, Sondey J, Maillot M, Guelinckx I, Lluh A. Drinking Water Intake Is Associated with Higher Diet Quality among French Adults. *Nutrients* 2016; **8**: [PMID: 27809236 DOI: 10.3390/nu8110689]
 - 35 **Casper RC**. Diet and mental health: an up-to-date analysis. *World Rev Nutr Diet* 2011; **102**: 98-113 [PMID: 21865824 DOI: 10.1159/000327798]
 - 36 **Ananth J**, Yassa R. Magnesium in mental illness. *Compr Psychiatry* 1979; **20**: 475-482 [PMID: 487803 DOI: 10.1016/0010-440X(79)90034-8]
 - 37 **Jordan J**, Shannon JR, Grogan E, Biaggioni I, Robertson D. A potent pressor response elicited by drinking water. *Lancet* 1999; **353**: 723 [PMID: 10073520 DOI: 10.1016/S0140-6736(99)99015-3]
 - 38 **Goekoop JG**, de Winter RF, Wolterbeek R, Van Kempen GM, Wiegant VM. Increased plasma norepinephrine concentration in psychotic depression. *Ther Adv Psychopharmacol* 2012; **2**: 51-63 [PMID: 23983957 DOI: 10.1177/2045125312436574]
 - 39 **Kelly MM**, Tyrka AR, Price LH, Carpenter LL. Sex differences in the use of coping strategies: predictors of anxiety and depressive symptoms. *Depress Anxiety* 2008; **25**: 839-846 [PMID: 17603810 DOI: 10.1002/da.20341]
 - 40 **Jimoh FO**, Bunn D, Hooper L. Assessment of a Self-Reported Drinks Diary for the Estimation of Drinks Intake by Care Home Residents: Fluid Intake Study in the Elderly (FISE). *J Nutr Health Aging* 2015;

- 19: 491-496 [PMID: 25923476 DOI: 10.1007/s12603-015-0458-3]
- 41 **Nissensohn M**, López-Ufano M, Castro-Quezada I, Serra-Majem L. Assessment of beverage intake and hydration status. *Nutr Hosp* 2015; **31** Suppl 3: 62-69 [PMID: 25719773 DOI: 10.3305/nh.2015.31. sup3.8753]
- 42 **Sidossis LS**. Assessment of energy expenditure and energy intake in children: is it possible? *Curr Op Clin Nutr Metab Care* 2003; **6**: 499 [DOI: 10.1097/00075197-200309000-00002]

P- Reviewer: Chakrabarti S, Hosak L, Khajehei M **S- Editor:** Ji FF
L- Editor: A **E- Editor:** Bian YN





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

