



Noninvasive assessment of fibrosis and steatosis in alcoholic liver diseas



网页 图片 新闻 视频 购物 更多 搜索工具

找到约 1,800 条结果 (用时 0.72 秒)

Google 学术: Noninvasive assessment of fibrosis and steatosis in alcoholic liver disease using Unidimensional Transient Elastography (Fibroscan®).

Transient elastography: a new noninvasive method for ... - Sandrin - 被引用次数: 1514

Noninvasive assessment of liver fibrosis by ... - Ziol - 被引用次数: 1184

... failure of liver stiffness measurement using FibroScan ... - Foucher - 被引用次数: 341

25 - EASL's International Liver Congress 2015

<https://ilc-congress.eu/cpgs-non-invasive/files/basic.../page25,ht...> 翻译此页

Assessment of asymptomatic liver fibrosis in alcoholic patients using [185] ... Ther 2008:1188-1198. for non-invasive assessment of liver fibrosis and steatosis in NAFLD. ... Liver fibrosis in viral hepatitis: noninvasive assessment with acoustic [187] ... to unidimensional transient elastography in the noninvasive assessment of ...

Noninvasive assessment of hepatic fibrosis: Ultrasound ...

www.uptodate.com/.../noninvasive-assessment-of-hepatic-fibro... 翻译此页

2015年3月18日 - Liver disease is an important problem worldwide. ... This is true for a wide range of disorders, including viral hepatitis, alcoholic and nonalcoholic ... Assessment of liver fibrosis using transient elastography in patients with alcoholic liver disease, ... Non-invasive diagnosis of liver steatosis using controlled ...

Methodology - EASL - Clinical Practice Guidelines

www.easl.eu/research/our.../non-invasive...liver-disease.../3 翻译此页

How does transient elastography (TE) perform for staging liver fibrosis? ... What are the indications for non-invasive tests for staging liver disease in non-alcoholic fatty liver

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 19467

Manuscript Type: TOPIC HIGHLIGHTS

Noninvasive assessment of alcoholic liver disease using unidimensional transient elastography (Fibroscan®)

Monica Lupsor-Platon, Radu Badea

Abstract

Unidimensional transient elastography (TE) is a noninvasive technique, which has been increasingly used in the assessment of diffuse liver diseases. This paper focuses on reviewing the existing data on the use of TE in the diagnosis of fibrosis and in monitoring disease progression in alcoholic liver disease, on the factors that may influence the result of fibrosis prediction, and last but not least, on its potential use in

Match Overview

1	Internet 424 words crawled on 16-Jun-2015 www.easl.eu	4%
2	CrossCheck 414 words Pavlov, Chavdar S, Giovanni Casazza, Dimitrinka Nikolova, Emmanuel Tsochatzis, Andrew K Burroughs, Vladimir	4%
3	Internet 394 words crawled on 25-Mar-2014 www.ncbi.nlm.nih.gov	4%
4	Internet 195 words crawled on 14-Dec-2013 www.jgld.ro	2%
5	CrossCheck 115 words Eve Gelsi. "Effect of Detoxification on Liver Stiffness Assessed by Fibroscan® in Alcoholic Patients : EFFECT C ...	1%
6	CrossCheck 112 words Sebastian Mueller. "Liver stiffness: a novel parameter for the diagnosis of liver disease", Hepatic Medicine Eviden	1%
7	CrossCheck 85 words Nahon, P.. "Assessment of liver fibrosis using transient elastography in patients with alcoholic liver disease", Jo	1%
8	Publications 72 words Lupsor-Platon, Monica; Stefănescu, Horia; Mureșan, Daniel; Florea, Mira; Erzsébet Szász, Mária; Maniu, Anca an	1%



Noninvasive assessment of alcoholic liver disease using unidimensional

[网页](#) [图片](#) [新闻](#) [视频](#) [更多](#) [搜索工具](#)

找到约 9,290 条结果 (用时 0.63 秒)

Google 学术: Noninvasive assessment of alcoholic liver disease using unidimensional transient elastography (Fibroscan®)

Transient elastography: a new noninvasive method for ... - Sandrin - 被引用次数: 1586

... failure of liver stiffness measurement using FibroScan ... - Foucher - 被引用次数: 357

Noninvasive assessment of liver fibrosis by ... - Ziol - 被引用次数: 1221

Non-invasive assessment of liver fibrosis

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) [翻译此页](#)

作者: V Papastergiou - 2012 - 被引用次数: 28 - [相关文章](#)

Keywords: liver fibrosis, non-invasive, transient elastography, serum biomarkers, ... shown to be reliable in the setting of non-alcoholic fatty liver disease (NAFLD); using a ... Development of combined indices comprising Fibrotest and Fibroscan to unidimensional transient elastography in the noninvasive assessment of ...

Acoustic Radiation Force Impulse and Supersonic Shear ...

www.sciencedirect.com/science/article/.../S0301562913007540 [翻译此页](#)

作者: I Şporea - 2013 - 被引用次数: 23 - [相关文章](#)

To assess the severity of chronic liver diseases, the hepatologist can use invasive (liver ... Subsequently, transient elastography (TE), commercialized as the FibroScan, ... hepatitis B (Chon et al., 2012 and Marcellin et al., 2009) and non-alcoholic ... methods have been used for the non-invasive assessment of liver fibrosis.

Noninvasive assessment of hepatic fibrosis: Ultrasound ...

www.uptodate.com/.../noninvasive-assessment-of-hepatic-fibro... [翻译此页](#)

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

 包括专利 包含引用 创建快讯

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

Transient elastography: a new noninvasive method for assessment of hepatic fibrosis

researchgate.net 中的 [PDF]

L Sandrin, B Fourquet, JM Hasquenoph, S Yon... - *Ultrasound in medicine ...*, 2003 - Elsevier
 ... Furthermore, due to the limited size of liver samples and the subjective assessment made by ...
 Therefore, there is an increasing need for alternative noninvasive methods to estimate the grade ...
 health problem and hepatic fibrosis grade is a key parameter to assess prognosis and ...

被引用次数：1599 相关文章 所有 12 个版本 引用 保存

Prevalence and factors associated with failure of liver stiffness measurement using FibroScan in a prospective study of 2114 examinations

J Foucher, L Castéra, PH Bernard... - *European journal of ...*, 2006 - journals.lww.com
 ... Zeitschrift Fur Gastroenterologie Transient elastography (FibroScan) for the non-invasive
 assessment of liver ... Disease Diagnosis of Nonalcoholic Fatty Liver Disease: Invasive versus
 Noninvasive Wieckowska, A ... Use of elastometry (FibroScan((R))) for the non-invasive staging ...

被引用次数：359 相关文章 所有 6 个版本 引用 保存

Noninvasive assessment of liver fibrosis by measurement of stiffness in patients with chronic hepatitis C

wiley.com 中的 [HTML]

M Ziol, A Handra-Luca, A Kettaneh, C Christidis... - ..., 2005 - Wiley Online Library
 ... 3-7 Transient elastography is a new technique that rapidly and noninvasively measures mean ...
 here need to be confirmed by independent validation studies, the simple, noninvasive, and well ...
 Transient elastography: a new non-invasive method for assessment of hepatic fibrosis. ...

被引用次数：1226 相关文章 所有 8 个版本 引用 保存

Magnetic resonance elastography for the noninvasive staging of liver fibrosis

L Huwart, C Sempoux, E Vicaut, N Salameh, L Annet... - *Gastroenterology*, 2008 - Elsevier
 ... To assess the quality of the model (and thus the robustness of our conclusions), we ... The results
 of our study show that among 3 noninvasive methods (MR elastography, ultrasound ... imaging for
 the detection of hepatocellular carcinomas and steatosis and the assessment of liver ...

被引用次数：402 相关文章 所有 10 个版本 引用 保存

Noninvasive assessment of liver fibrosis by measurement of stiffness in patients with nonalcoholic fatty liver disease (NAFLD)

M Yoneda, H Mawatari, K Fuiita, H Endo, H Iida... - ... and Liver Disease. 2008 - Elsevier