[PDF] Impact of adjuvant chemotherapy on patients with ... https://wjso.biomedcentral.com/track/pdf/10.1186/s12957-018-1455-x

rectal cancer improved disease-free and overall survival of patients with ypT0–2 ypN0 tumor stages in our study. In particular, ypT2 ypN0 patients seem to profit from adjuvant treatment. Keywords: Rectal neoplasms, Neoadjuvant therapy, Chemoradiotherapy, Adjuvant chemotherapy, Disease-free survival

Role of adjuvant chemotherapy in locally advanced rectal ... https://www.ncbi.nlm.nih.gov/pubmed/28865435

Sep 02, 2017 · 1. BMC Cancer. 2017 Sep 2;17(1):615. doi: 10.1186/s12885-017-3624-7. Role of adjuvant chemotherapy in locally advanced rectal cancer with ypT0-3N0 after preoperative chemoradiation therapy and surgery.

Cited by: 6 Author: Chang Gon Kim, Chang Gon Kim, Joong ...

Publish Year: 2017

Impact of adjuvant chemotherapy on patients with ypT0-2 ...

https://wjso.biomedcentral.com/articles/10.1186/s12957-018-1455-x •

To investigate the importance of adjuvant chemotherapy in locally advanced rectal cancer (\geq cT3 or N+) staged ypT0-2 ypN0 on final histological work-up after neoadjuvant chemoradiation and radical resection. The clinical course of patients with rectal cancer and ypT0-2 ypN0 stages after neoadjuvant chemoradiation and radical resection was analyzed from 1999 to 2012.

Cited by: 1 Author: Christian Galata, Kirsten Merx, Sabine M...


Publish Year: 2018


Is adjuvant chemotherapy necessary for patients with ypT0 ...

https://pubmed.ncbi.nlm.nih.gov/30430016

Background and objective: The benefit from adjuvant chemotherapy for patients treated with neoadjuvant chemoradiotherapy (NCRT) and curative surgery remains controversial, particularly among those responding well to NCRT. This retrospective study aimed to clarify the benefits of adjuvant chemotherapy in terms of the oncological outcomes of patients with ypT0-2N0 rectal cancer after ...

Rectal cancer patients with downstaging after neoadjuvant ... https://pubmed.ncbi.nlm.nih.gov/32647668





Evaluating the benefit of adjuvant chemotherapy in patients with y

===





ALL

IMAGES

VIDEOS

13,800 Results

Any time ▼

Impact of adjuvant chemotherapy on patients with ypT0-2 ...

https://wjso.biomedcentral.com/articles/10.1186/s12957-018-1455-x •

Aug 02, 2018 · Conclusion. Administration of adjuvant chemotherapy after neoadjuvant chemoradiation and radical resection in rectal cancer improved disease-free and overall survival of patients with ypT0–2 ypN0 tumor stages in our study. In particular, ypT2 ypN0 patients seem to profit from adjuvant treatment.

Cited by: 3 Author: Christian Galata, Kirsten Merx, Sabine M...

Publish Year: 2018 Estimated Reading Time: 11 mins

Role of Adjuvant Chemotherapy in ypT0-2N0 Patients ...

https://www.sciencedirect.com/science/article/pii/S0360301615001984

Jul 01, 2015 - To explore the role of adjuvant chemotherapy for patients with ypT0-2N0 rectal cancer treated by preoperative chemoradiation therapy (PCRT) and radical resection. Patients and Methods A national consortium of 10 institutions was formed, and patients with ypT0-2N0 mid- and low-rectal cancer after PCRT and radical resection from 2004 to 2009 were included.

Cited by: 14 Author: In Ja Park, Dae Yong Kim, Hee Cheol Ki...

Publish Year: 2015

[PDF] Impact of adjuvant chemotherapy on patients with ...

https://wjso.biomedcentral.com/track/pdf/10.1186/s12957-018-1455-x

survival in ypT0-1 ypN0 patients showed no correlation to the administration of adjuvant chemotherapy. Conclusion: Administration of adjuvant chemotherapy after neoadjuvant chemoradiation and radical resection in rectal cancer improved disease-free and overall survival of patients with ypT0-2 ypN0 tumor stages in our study.

Impact of adjuvant chemotherapy on patients with ypT0-2 ... europepmc.org/abstract/MED/30071852 •

Aug 02, 2018 · Fifty-four patients with adjuvant (ACT) and 50 patients without adjuvant chemotherapy (NACT) after neoadjuvant chemoradiation followed by radical resection for rectal cancer were included in the analysis. Mean follow-up was 68 ± 33.7 months. One patient without adjuvant chemotherapy and none in the ACT group developed a local recurrence.

Cited by: 3 Author: Christian Galata, Kirsten Merx, Sabine M...

Publish Year: 2018