

学术搜索

文章

我的图书馆

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

[Performance of 18Fluorodeoxyglucose-Positron Emission Tomography and Somatostatin Receptor Scintigraphy for High Ki67 \(\$\geq 10\%\$ \) Well-Differentiated Endocrine ...](#)

R Abgral, S Leboulleux, D Déandreis... - The Journal of ..., 2010 - press.endocrine.org

Objective: The purpose of this prospective study was to compare the performance of ^{111}In -octreotide somatostatin receptor scintigraphy (SRS) and ^{18}F fluorodesoxyglucose positron emission tomography (FDG-PET) in aggressive well-differentiated endocrine carcinoma (WDEC) defined by a high Ki67 ($\geq 10\%$). Methods: Eighteen consecutive patients explored in a single hospital between November 2003 and 2008 for high Ki67 ($\geq 10\%$) WDEC ...

被引用次数: 41 相关文章 所有 5 个版本 引用 保存

[\[HTML\] endocrine.org](#)

以上显示的是最相近的搜索结果。 [查看全部搜索结果](#)

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 32410

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Performance of 18-fluoro-2-deoxyglucose positron emission tomography for esophageal cancer screening

Sekiguchi M *et al.* FDG-PET for esophageal cancer screening

Masau Sekiguchi, Takashi Terauchi, Yasuo Kakugawa, Naoki Shimada, Yutaka Saito, Takahisa Matsuda

Masau Sekiguchi, Yasuo Kakugawa, Takahisa Matsuda, Cancer Screening Center, National Cancer Center Hospital, Tokyo 104-0045, Japan

Match Overview

1	Internet 100 words crawled on 20-Feb-2017 researchonline.jcu.edu.au	2%
2	Internet 64 words crawled on 16-Jul-2016 www.wjgnet.com	2%
3	Crossref 21 words Kakugawa, Yasuo. "New reduced volume preparation reg... men in colon capsule endoscopy", World Journal of Gastro	1%
4	Crossref 20 words Sekiguchi, Masau, Yasuo Kakugawa, Takashi Terauchi, Mi nori Matsumoto, Hiroshi Saito, Yukio Muramatsu, Yutaka S	<1%

学术搜索

文章

我的图书馆

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

Performance of 18fluorodeoxyglucose-positron emission tomography and somatostatin receptor scintigraphy for high Ki67 ($\geq 10\%$) well-differentiated endocrine ...

R Abgral, S Leboulleux, D Déandreis... - The Journal of ..., 2010 - press.endocrine.org

Objective: The purpose of this prospective study was to compare the performance of ^{111}In -octreotide somatostatin receptor scintigraphy (SRS) and 18fluorodesoxyglucose positron emission tomography (FDG-PET) in aggressive well-differentiated endocrine carcinoma (WDEC) defined by a high Ki67 ($\geq 10\%$). Methods: Eighteen consecutive patients explored in a single hospital between November 2003 and 2008 for high Ki67 ($\geq 10\%$) WDEC ...

被引用次数: 46 相关文章 所有 5 个版本 引用 保存

[\[HTML\]](#) [endocrine.org](#)

以上显示的是最相近的搜索结果。 [查看全部搜索结果](#)

[All](#)[Images](#)[News](#)[Shopping](#)[Videos](#)[More](#)[Settings](#)[Tools](#)

About 29,300 results (0.93 seconds)

Scholarly articles for Performance of 18-fluoro-2-deoxyglucose positron emission tomography for esophageal cancer screening

... -fluoro-2-deoxyglucose positron emission tomography ... - Han - Cited by 32

... emission tomography and computed tomography ... - Lardinois - Cited by 117

... cancer with integrated positron-emission tomography ... - Lardinois - Cited by 1494

Role of (18F) 2-fluoro-2-deoxyglucose positron emission tomography ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC)

by EC Smyth - 2011 - Cited by 47 - Related articles

Dec 14, 2011 - Whole-body **positron emission tomography** (PET) scanning after the ... Role of PET in staging the depth of disease-**esophageal carcinoma** metastatic disease, where its **performance** is superior than in the detection of the depth of the PET-CT detected 2/20 cancers from 2861 patients **screened** giving a ...

F-18-fluoro-2-deoxyglucose positron emission tomography (PET) and ...

<https://systematicreviewsjournal.biomedcentral.com/articles/10.1186/2046-4053-1-62>

by MA Schröer-Günther - 2012 - Cited by 18 - Related articles

Dec 13, 2012 - F-**18-fluoro-2-deoxyglucose positron emission tomography** (PET) ... Computed tomography Staging Recurrence **Esophageal** neoplasms Systematic review. Introduction. Malignant melanoma accounts for about 5% of all skin **cancers** [1]. ... Assessment Database were **screened** to identify systematic reviews.

Role of fluoro-2-deoxyglucose positron emission tomography for the ...

onlinelibrary.wiley.com/doi/10.1111/j.1759-7714.2011.00060.x/full

by K Isobe - 2011 - Cited by 1 - Related articles

Nov 1, 2011 - **Fluoro-2-deoxyglucose** (FDG) **positron emission tomography** (PET) is useful for the ... of PET has not yet been established for the **screening** of lung cancer patients for ... rectal cancer in one), gastric cancer in one, **esophageal cancer** in one, ... Male, 18. Female, 7. ECOG Performance status 0-12 1-10 0-0