

87866_Auto_EditedC.docx

*Case Control Study***Childhood asthma biomarkers including Zinc: An exploratory cross-sectional study**

Hoda Atef Abdelsattar Ibrahim, Mona Mohsen, Boles Salep Aziz Hanna, Dina Mahmoud, Khaled Mohamed Abdelhamid El-Khashab

Abstract**BACKGROUND**

Childhood bronchial asthma (BA) is a chronic inflammatory respiratory disease. Nutritional conditions, including zinc deficiency, can affect such allergic disorders.

AIM

To outline the difference in serum zinc levels between asthmatic children and healthy controls.

METHODS

A cross-sectional study was carried out at Children's Hospital, Cairo University, investigating serum zinc levels in children with BA ($n = 40$) and healthy children ($n = 21$). Other markers included serum ferritin, iron, hemoglobin (Hb), and immunoglobulin E (IgE) levels. Independent t -tests and Mann-Whinny tests were used for comparisons. The Kruskal-Wallis test was applied to compare serum ferritin and IgE levels with regard to asthma severity. Spearman's rank correlation was performed to explore the relationship between serum ferritin levels and both iron and Hb levels in asthmatic children.

RESULTS

Children with BA had higher levels of zinc, yet the difference was not significant ($P = 0.115$). Serum ferritin and IgE levels were significantly higher in asthmatic children ($P = 0.006$ and 0.001 , respectively), yet their levels did not differ significantly by severity ($P = 0.623$ and 0.126 , respectively). There was a nonsignificant weak correlation between serum ferritin levels and both serum iron and Hb levels.

CONCLUSION

Serum zinc levels do not seem to differ between asthmatic children and healthy children. Serum ferritin levels may be a marker of asthma control. Serum IgE levels are not markers of asthma severity.

Key Words: Children; Asthma; Zinc; Ferritin

Atef Abdelsattar Ibrahim H, Mohsen M, Salep Aziz Hanna B, Mahmoud D, Mohamed Abdelhamid El-Khashab K. Childhood asthma biomarkers including Zinc: An exploratory cross-sectional study. *World J Clin Pediatr* 2023; In press

Core Tip: Serum zinc levels were higher in asthmatic children than in nonasthmatic children. However, the difference was not significant. Serum ferritin levels were significantly higher in asthmatic children, which may be due to its immunosuppressive properties. Serum ferritin should not be considered in the diagnosis of iron deficiency anemia in asthmatic children. Serum immunoglobulin E should not be applied to diagnose the severity of childhood asthma. Further studies that track biomarkers such as ferritin during asthma progression are needed.

INTRODUCTION

The most prevalent chronic respiratory condition in children is ⁵ bronchial asthma (BA). BA is a chronic inflammatory disease of the lungs that causes airway inflammation and bronchial hyperreactivity; it may also be described as intermittent, reversible airway blockages^[1]. Due to its immune-modulating properties, zinc has attracted much attention in relation to asthma and airway inflammation. Zinc is a crucial trace element for human metabolism and helps regulate gene expression, enzyme activity, and protein structure. Additionally, it is crucial for immune system regulation and functions as an antioxidant, anti-inflammatory, and antiapoptotic agent^[2].

In the presence of continuous inflammation, free serum ferritin levels are elevated; in addition, free serum ferritin has a protective function in redox biology and iron homeostasis. In contrast, new research reveals that ferritin may have a causal role in the inflammatory pathology of illness, including rheumatologic, immunologic, neoplastic, and infectious diseases, and ferritin levels may be fundamental in the pathology of disease and help in predicting prognosis in addition to tracking disease activity^[3].

Since the beginning of the 20th century, it has been known that immunoglobulin E (IgE) is different from other immunoglobulin isotypes in that it may both trigger ⁵ extremely rapid pathological reactions and serve as a highly sensitive immunological amplifier. Furthermore, it is well known that patients with atopic diseases have higher IgE levels and that IgE serves as a vital link between the adaptive immune system's role in antigen recognition ¹¹ and the effector functions of mast cells and basophils at mucosal and cutaneous sites of environmental exposure. Due to these roles, IgE has become a ⁵ desirable target for pharmacological intervention, and IgE blocking has clinical potential in a wide range of therapeutic fields^[4,5].

³³ Our study aimed to identify the difference in serum zinc levels between asthmatic children and healthy controls. Additionally, other labs of interest, such as serum ferritin and IgE levels, were studied. Moreover, the possible role of these findings as markers of controlled asthma was investigated.

MATERIALS AND METHODS

This exploratory cross-sectional study was carried out at Children's Hospital Cairo University between May 2022 and October 2022. Sixty-one children were enrolled [40 cases (asthmatic), 21 controls (non-asthmatic)]. The control group consisted of healthy children of comparable age and sex to the cases who had no disease based on history and physical examination and no history of BA or zinc deficiency. All asthmatic children who were followed in the asthma clinic, aged from 5-12 years, and whose parents or caregivers approved participation were included. Owing to the absence of the reliability of lung function tests in children under five, as they are rarely practical^[6], children aged less than 5 years were excluded. Data on sociodemographic and clinical characteristics such as body mass index (BMI) and degree of asthma severity were collected. In addition, laboratory findings such as serum zinc, albumin, ferritin, and IgE levels were recorded and compared between cases and controls.

Case definition

The diagnosis of BA was considered using the medical history, family history, clinical examination, and symptoms including episodic dyspnea, coughing, wheezing, and tightness in the chest, as well as laboratory findings. The results of pulmonary function tests allowed for the confirmation of the diagnosis of BA and a determination of the severity and reversibility of airflow restriction^[7,8]. Malnutrition was defined using World Health Organization definitions^[9-12].

Control of potential bias

As our primary outcome was to compare cases and controls with regard to zinc levels, we essentially excluded participants with a drug history of zinc supplementation. We also excluded cases with a drug history of iron therapy, as our secondary outcome was to examine the difference between the two groups with regard to iron hemostasis.

Children admitted to the hospital were excluded because in-hospital admission could negatively affect their nutritional status and zinc level. Likewise, children with comorbidities that could affect their nutritional status were excluded.

Sample size calculation

The primary objective of the current study was to compare serum zinc levels between asthmatic children and healthy controls. Umar *et al*^[13] reported that the mean serum zinc level in BA patients was 79.63 ± 9.62 $\mu\text{g/dL}$, while it was 93.27 ± 12.21 $\mu\text{g/dL}$ in healthy controls. G*Power software (version 3.1.9.2) was used to estimate the required sample size. The alpha was set as 0.05, the power ($1-\beta$) was set as 0.99, and the case-to-control ratio was set as 2:1. Considering a nonparticipation rate of 10%, the minimum required sample size for the study was 60 patients, including 40 cases, and 20 controls.

Ethical concerns

All procedures were carried out in line with the ethical standards of the responsible committee on human experimentation and with the Helsinki Declaration of 2013 and were approved by the Research Ethics Committee of the Faculty of Medicine, Cairo University. The ethical approval number is MS-587-2021.

Statistical analysis

Data are statistically described in terms of the mean \pm SD, the median and interquartile range, or frequencies (number of cases) and percentages when appropriate. Tests of normality were performed for all the numerical variables of interest using the Kolmogorov-Smirnov/Shapiro-Wilk test. The comparison of numerical variables between cases and controls was performed using the independent samples *t*-test for the parametric data and the Mann-Whitney *U* test for nonparametric statistics. When the analyses between more than 2 groups (between children with mild, moderate, and severe asthma) were needed, the Kruskal-Wallis test for nonparametric data was performed (as in the comparison regarding ferritin and IgE levels). Cross-tabulation was applied to

compare the categorical variables using chi-square and Fisher's exact tests depending on whether more than 20% of the cells had expected cell counts less than 5. Spearman's rank correlation was applied if one or both of the numerical data of interest were not parametrically distributed, as in the case of the correlation between serum ferritin levels and both serum iron and hemoglobin levels. A two-sided P value less than or equal to 0.05 was considered statistically significant. A graphical presentation was also used to illustrate the difference between medians using a box plot and the difference between means using error plot graphs. Furthermore, a scattered plot was applied to clearly illustrate the possible linear relation between serum iron and ferritin levels. All statistical calculations were performed using the computer program IBM SPSS (Statistical Package for the Social Science; IBM Corp, Armonk, NY, United States) release 27 for Microsoft Windows.

RESULTS

Our study aimed to describe the difference in zinc levels between children with BA and healthy controls without BA.

Preliminary analysis

Matching between cases and controls regarding age, sex, BMI, and the coexistence of malnutrition was performed (Table 1). Table 1 shows no significant differences, with P values more than 0.05

The normal distribution of the numerical variables of interest was tested using the Kolmogorov-Smirnov/Shapiro-Wilk test to identify the possible statistical methods of choice (Table 2). Age, serum ferritin levels, IgE levels, and BMI scores were not normally distributed. On the other hand, serum zinc, iron, hemoglobin (Hb) and albumin levels were normally distributed.

Sociodemographic criteria

Table 3 illustrates the sociodemographic criteria of the study participants. Among the study participants, male sex generally predominated, specifically in children with BA (Figure 1).

Characteristics of the study participants

Table 4 shows the distribution and level of significance of the biochemical laboratory assessments between cases and controls. The mean \pm SD serum zinc level for asthmatic *vs* nonasthmatic children was 94.4 ± 24.7 and 85.2 ± 19 , respectively, with no significant difference ($P = 0.115$). Similarly, no significant differences were observed regarding serum iron, Hb, and albumin levels in cases *vs* controls ($P = 0.389$, 0.857 , and 0.391 , respectively). In contrast, serum levels of IgE and ferritin showed significant differences between cases and controls, with P values of 0.001 and 0.006 , respectively.

Our primary objective was to compare zinc levels between asthmatic and nonasthmatic children. The mean difference was higher in asthmatic children, as shown in Figure 2.

Upon determining the serum levels of ferritin, median differences were higher in children with BA, as shown in Figure 3. This may disclose its role in inflammation, as higher serum ferritin levels do not necessarily mean higher serum iron levels. Spearman's rank correlation was performed and revealed a weak nonsignificant correlation between both serum ferritin and iron levels ($r_s = -0.077$, $P = 0.637$), as shown in Table 5, which also yielded a similar weak nonsignificant correlation between serum ferritin and Hb levels ($r_s = 0.204$, $P = 0.208$). In addition, a weak relation was observed in the scatter plot for ferritin and iron levels, as shown in Figure 4.

Upon checking the median difference in IgE levels, children with BA showed higher levels, as shown in Figure 5. However, serum IgE levels did not differ significantly in regard to the degree of asthma in children with BA, as shown in Table 6 and Figure 6. Likewise, serum ferritin levels did not show a significant difference regarding the grades of asthma severity (Table 6).

DISCUSSION

Our study included 61 children: 40 diagnosed with BA and 21 without BA. We examined zinc levels between the two groups. Yousef *et al*^[14] detected significantly diminished levels in cases with BA in comparison to controls, and the P value was < 0.01 . In addition, higher levels of serum zinc in control children were observed in another former study, yet the difference was not significant ($P = 0.388$)^[15]. Our study showed a different result. For example, mean values of serum zinc were higher in cases with BA in our study. This difference may be because all cases were receiving asthma therapy such as inhaled steroids. Rahman *et al*^[16] proposed that after steroid therapy, stimulation of glutathione (GSH) synthesis in the liver occurs due to a decrease in the generation of reactive oxygen species by neutrophils. Reduced GSH and GSH disulfide (GSSG) are critical modulators of both the rate of zinc transfer and the ultimate number of zinc atoms transferred. GSSG increases the rate of zinc transfer by 3-fold, and its concentration is the major determinant for efficient zinc transfer^[17]. In another study, Raevé *et al*^[18] showed that after corticosteroid therapy, macrophage oxidant production decreased, and the number of oxidant-generating cells present in the asthmatic airway mucosa also decreased; hence, the enhancement of GSH synthesis in the liver could subsequently occur. Considering that lower serum zinc levels indicate higher asthma severity^[19], our finding of higher zinc levels in asthmatic children may suggest that lower serum zinc levels may indicate poor compliance with therapy and vice versa.

Other biochemical laboratory test results were assessed, all of which showed no significant difference between cases and controls except for serum ferritin and serum IgE levels. Serum ferritin appears to be a better biomarker for inflammation than iron status^[20]. This may be the reason for the significantly elevated levels of ferritin in children with BA compared to controls in our study ($P = 0.006$). There is mounting evidence that circulating ferritin levels might not only reflect the acute phase response but also play a crucial role in inflammation. Its secretion is regulated via proinflammatory cytokines, and ferritin has immunosuppressive effects that are probably mediated by binding to its receptor. Although it is commonly accepted that circulating ferritin levels may reflect an

acute phase response, the explanation for how and why serum ferritin is increased is unknown^[21]. Higher ferritin does not essentially equal iron overload^[22]. Ferritin can be a good biomarker of appropriate *vs* excessive inflammation, and previous research found that high ferritin in severe coronavirus disease 2019 pneumonia patients is associated with improved outcomes following steroid treatment^[23]. Another study found that low ferritin levels in the course of steroid therapy were linked to greater mortality^[24]. Therefore, our study may increase attention toward the possible use of ferritin as a marker of asthma control after steroid therapy. In addition, serum ferritin is not significantly correlated with Hb and iron. However, using serum ferritin as a marker of iron hemostasis or iron deficiency anemia in asthmatic children may be controversial.

Allergic diseases involving asthma are characterized by an increase in serum IgE levels^[25,26]. Our study showed that there was a significant increase in serum IgE levels in patients with BA. However, there was no significant difference regarding the degrees of severity. Sandeep *et al*^[27] reported a similar finding. It may be suggested that levels of IgE are quite high at the local inflammation site and that the serum levels do not essentially reflect the levels in the lungs or bronchus. Moreover, IgE is bound to mast cells with rather high affinity, and hence, circulating IgE may not provide conclusive evidence of the severity of inflammation^[28].

CONCLUSION

Serum zinc levels did not show a significant difference between asthmatic children and nonasthmatic children. Serum ferritin may be a marker of controlled asthma. Serum IgE levels should not be used to stratify asthmatic children according to severity.

ARTICLE HIGHLIGHTS

Research background

Zinc levels might differ in asthmatic children.

Research motivation

The possible role of the biochemical nutritional assessment including zinc to be a biomarker for asthma severity.

Research objectives

To outline the difference in ¹⁶zinc levels between asthmatic and healthy children.

Research ¹⁵methods

A cross-sectional study was carried out investigating serum ¹⁶zinc levels in asthmatic and healthy children.

Research results

Zinc levels weren't different. Ferritin levels were significantly higher in cases with bronchial asthma.

Research conclusions

Ferritin could be used as a future biomarker for asthma controller therapy.

Research perspectives

Further studies investigating the possible role of ferritin and other possible biomarkers for asthma severity should be outlined.

24%

SIMILARITY INDEX

PRIMARY SOURCES

1	link.springer.com Internet	69 words — 3%
2	cyberleninka.org Internet	57 words — 2%
3	www.ncbi.nlm.nih.gov Internet	54 words — 2%
4	www.researchgate.net Internet	50 words — 2%
5	respiratory-research.biomedcentral.com Internet	38 words — 1%
6	www.frontiersin.org Internet	28 words — 1%
7	jmisr.researchcommons.org Internet	27 words — 1%
8	mafiadoc.com Internet	27 words — 1%
9	sciencescholar.us Internet	25 words — 1%

-
- 10 Aida M. Yousef, Ekramy Elmorsy. "Serum zinc level in bronchial asthma", Egyptian Journal of Chest Diseases and Tuberculosis, 2017
Crossref 21 words — 1%
-
- 11 mcmed.us
Internet 19 words — 1%
-
- 12 tkb.dergisi.org
Internet 18 words — 1%
-
- 13 www.karger.com
Internet 18 words — 1%
-
- 14 eprints.hud.ac.uk
Internet 17 words — 1%
-
- 15 Amira M. M. Hamed, Yasser T. Kassem, Hamada K. Fayed, Ahmed M. Solaiman. "Serum zinc levels in hospitalized children with pneumonia: a hospital-based case-control study", Egyptian Journal of Bronchology, 2020
Crossref 15 words — 1%
-
- 16 E. Arik Yilmaz. "Erythrocyte zinc levels in children with bronchial asthma", Pediatric Pulmonology, 12/2011
Crossref 12 words — < 1%
-
- 17 www.mdpi.com
Internet 12 words — < 1%
-
- 18 Eirini Maratou, Dimitrios J Hadjidakis, Anastasios Kollias, Katerina Tsegka et al. "Studies of insulin resistance in patients with clinical and subclinical hypothyroidism", European Journal of Endocrinology, 2009
Crossref 11 words — < 1%

-
- 19 genesenvironment.biomedcentral.com 11 words — < 1%
Internet
-
- 20 repository.publisso.de 11 words — < 1%
Internet
-
- 21 www.imrpress.com 11 words — < 1%
Internet
-
- 22 SYH-JAE LIN. "Influence of Atopy and Concomitant Sinusitis on Serum Eosinophil Cationic Protein Levels in Asthmatic Children", *Pediatric Asthma Allergy & Immunology*, 01/2000 10 words — < 1%
Crossref
-
- 23 Jun-Hwa Lee. "Comparison of Serum Zinc Levels Measured by Inductively Coupled Plasma Mass Spectrometry in Preschool Children with Febrile and Afebrile Seizures", *Annals of Laboratory Medicine*, 2012 9 words — < 1%
Crossref
-
- 24 brieflands.com 9 words — < 1%
Internet
-
- 25 worldwidescience.org 9 words — < 1%
Internet
-
- 26 Beatrijs L. P. Bloemers, Louis Bont, Roel A. de Weger, Sigrid A. Otto, Jose A. Borghans, Kiki Tesselaar. "Decreased Thymic Output Accounts for Decreased Naive T Cell Numbers in Children with Down Syndrome", *The Journal of Immunology*, 2011 8 words — < 1%
Crossref
-
- 27 Bernd W. Scheithauer, Thomas A. Gaffey, Ricardo V. Lloyd, Thomas J. Sebo et al. "Pathobiology of Pituitary Adenomas and Carcinomas", *Neurosurgery*, 2006 8 words — < 1%

28 Hirotaka Yamashita. "Dermal mast cells play a central role in the incidence of scratching behavior in mice induced by multiple application of the hapten, 2,4,6-trinitrochlorobenzene", *Experimental Dermatology*, 6/2005

Crossref

8 words — < 1%

29 Zhinuan Hong, Zhixin Huang, Zhen Chen, Mingqiang Kang. "Prognostic Value of Carcinoembryonic Antigen Changes Before and After Operation for Esophageal Squamous Cell Carcinoma", *World Journal of Surgery*, 2022

Crossref

8 words — < 1%

30 f6publishing.blob.core.windows.net

Internet

8 words — < 1%

31 journalrepository.org

Internet

8 words — < 1%

32 researchonline.gcu.ac.uk

Internet

8 words — < 1%

33 www.ejca.eg.net

Internet

8 words — < 1%

34 Shaker, Olfat G., Nermin Abdel Hamid Sadik, and Nehal Abd El-Hamid. "Impact of single nucleotide polymorphism in tumor necrosis factor- α gene 308G/A in Egyptian asthmatic children and wheezing infants", *Human Immunology*, 2013.

Crossref

7 words — < 1%

EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE MATCHES OFF