

学术搜索

找到约 152 条结果 (用时0.06秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

包括专利

包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置, 指定搜索语言

Increased serum S100β protein concentrations following severe head injury in humans: a biochemical marker of brain death?

A Regner, M Kaufman, G Friedman, I Chemale - Neuroreport, 2001 - journals.lww.com
 ... Brain CT was performed within 6 h in all cases, ... I; Zanoni, C; Fossa, A; Gehrke, JT; Jotz, GP; Kaufmann, M; Simon, D; Regner, A Clinical Chemistry and ... Liver International Serum S100b (astrocyte-specific protein) is a useful marker of hepatic encephalopathy in patients with ...
 被引用次数: 47 相关文章 所有 5 个版本 引用 保存

Elevated serum levels of astroglial S100β in patients with liver cirrhosis indicate early and subclinical portal-systemic encephalopathy

J Wiltfang, W Nolte, M Otto, J Wildberg, E Bahn... - Metabolic brain ..., 1999 - Springer
 ... In all cases, a p value of less than 0.05 (two-sided level of significance) was considered ... Figure 1: Serum S100B was significantly increased in the presence of early clinically overt PSE (PSE ...
 Initial clinical experiences with TIPS (transjugular intrahepatic portasystemic stent- shunt ...
 被引用次数: 25 相关文章 所有 7 个版本 引用 保存

Sepsis-associated delirium

M Ebersoldt, T Sharshar, D Annane - Intensive care medicine, 2007 - Springer
 ... may be detected in almost all patients [1, 4-6]. The severity of SAD, whether assessed clinically [2, 3 ... examination prior to ICU discharge and showed ischemic and hemorrhagic lesions in 7% and 10% of cases, respectively. ... However, no clinical or experimental data are available ...
 被引用次数: 174 相关文章 所有 9 个版本 引用 保存

A novel marker for traumatic brain injury: CSF αII-spectrin breakdown product levels

NC Ringger, BE O'steen, JG Brabham... - Journal of ..., 2004 - online.liebertpub.com
 ... levels are a promising biomarker of injury and further study is warranted in clinical TBI. ... the National Institutes of Health guide- lines detailed in the Guide for the Care and Use of Lab ... The CCI model that we have used in these studies has some inherent variability in impact force ...
 被引用次数: 91 相关文章 所有 9 个版本 引用 保存

Biochemical markers of cerebrovascular injury in sleep apnoea syndrome

W Jordan, J Hagedohm, J Wiltfang... - European ..., 2002 - Eur Respiratory Soc
 ... by lumbar puncture is only indicated for diagnostic purposes in severe cases and is ... Patients with

area-c54.it 中的 [PDF]

researchgate.net 中的 [PDF]

ersjournals.com 中的 [HTML]

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 24571

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Clinical scenarios for the use of S100 β as a marker of hepatic encephalopathy

Duarte-Rojo A *et al.* S100 β as a marker of hepatic encephalopathy

Andrés Duarte-Rojo, Astrid Ruiz-Margáin, Ricardo U Macias-Rodriguez, Francisco Javier Cubero, José Estradas-Trujillo, Rosa Ma. Muñoz-Fuentes, Aldo Torre

Andrés Duarte-Rojo, University of Arkansas for Medical Sciences, Little Rock, AR 72205, United States

Astrid Ruiz-Margáin, Ricardo U Macias-Rodriguez, José Estradas-Trujillo, Rosa Ma. Muñoz-Fuentes, Aldo Torre, Department of Gastroenterology, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City 14000,

Match Overview

1	Internet 112 words crawled on 02-Mar-2016 www.wjnet.com	3%
2	Internet 42 words crawled on 01-Mar-2008 intl.ajronline.org	1%
3	CrossCheck 27 words Jon Sen. "S100B in neuropathologic states: The CRP of the brain?", <i>Journal of Neuroscience Research</i> , 05/15/2007	1%
4	Internet 22 words crawled on 31-Jul-2013 www.ncbi.nlm.nih.gov	1%

学术搜索

找到约 205 条结果 (用时0.07秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

包括专利

包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Increased serum S100β protein concentrations following severe head injury in humans: a biochemical marker of brain death?

A Regner, M Kaufman, G Friedman, I Chemale - Neuroreport, 2001 - journals.lww.com

... Brain CT was performed within 6 h in all cases. ... Key words: Brain death; Head injury; Human; S100β protein. ... da Rocha, AB; Schneider, RF; de Freitas, GR; Andre, C; Grivicich, I; Zaroni, C; Fossa, A; Gehrke, JT; Jotz, GP; Kaufmann, M; Simon, D; Regner, A Clinical Chemistry and ...

被引用次数: 47 相关文章 所有 5 个版本 引用 保存

Elevated serum levels of astroglial S100β in patients with liver cirrhosis indicate early and subclinical portal-systemic encephalopathy

J Wiltfang, W Nolte, M Otto, J Wildberg, E Bahn... - Metabolic brain ..., 1999 - Springer

... In all cases, a p value of less than 0.05 (two-sided level of significance) was considered ... Figure 1: Serum S100B was significantly increased in the presence of early clinically overt PSE (PSE ... Initial clinical experiences with TIPS (transjugular intrahepatic portosystemic stent-shunt ...

被引用次数: 26 相关文章 所有 7 个版本 引用 保存

[HTML] Sepsis-associated encephalopathy: not just delirium

FG Zampieri, M Park, FS Machado, LCP Azevedo - Clinics, 2011 - SciELO Brasil

... family and is associated with regulation of programmed cell death on various biological scenarios. ... brain imaging resonance in SAE and performed autopsies in selected cases 42 with ... Primeau F, Elie M. Use Of Medications With Anticholinergic Effect Predicts Clinical Severity Of ...

被引用次数: 46 相关文章 所有 13 个版本 引用 保存 更多

Sepsis-associated delirium

M Ebersoldt, T Sharshar, D Annane - Intensive care medicine, 2007 - Springer

... in almost all patients [1, 4-6]. The severity of SAD, whether assessed clinically [2, 3 ... If the clinical examination is unrevealing, the diagnosis of SAD should rely on laboratory ... brain injuries as suggested by postmortem studies showing ischemic lesions in all cases, hemorrhage in ...

被引用次数: 179 相关文章 所有 10 个版本 引用 保存

S100β protein: The preoperative new clinical indicator of brain damage in patients with fulminant hepatic failure

S Shiotani, M Shimada, Y Soejima, T Yoshizumi... - Transplantation ..., 2004 - Elsevier

... Please enable JavaScript to use all the features on this page. ... In FHF, we sometimes encounter cases where the EEG is able to return to normal after liver transplantation ... conclusion, these results suggest that serum S100β protein has the potential to be a new clinical indicator of ...

被引用次数: 2 相关文章 所有 7 个版本 引用 保存

... due to extreme environmental temperature by quantification of basic fibroblast growth factor

scielo.br 中的 [HTML]

area-c54.it 中的 [PDF]

全部 新闻 图片 视频 购物 更多 ▾ 搜索工具

找到约 226,000 条结果 (用时 0.56 秒)

Hepatic encephalopathy - Wikipedia, the free encyclopedia

https://en.wikipedia.org/wiki/Hepatic_encephalopathy ▾ 翻译此页

The mildest form of **hepatic encephalopathy** is difficult to detect **clinically**, but may be ... In a small proportion of **cases**, the encephalopathy is caused directly by liver ... This is **used** in the treatment of refractory ascites, bleeding from oesophageal the prognosis is determined largely by other **markers of liver** failure, such as

Sepsis-associated encephalopathy: not just delirium - NCBI

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: FG Zampieri - 2011 - 被引用次数: 46 - 相关文章

Brain dysfunction, or sepsis-associated **encephalopathy** (SAE), has been ... well-established **clinical** or biological **markers** of damage to the brain during sepsis. ... Han et al. associated the **use** of anticholinergic drugs with symptoms of delirium. Increases in **S100 β** levels in critically ill patients without apparent brain ...

Minimal Hepatic Encephalopathy in Children: Evaluation ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: BR Foerster - 2009 - 被引用次数: 22 - 相关文章

2009年6月9日 - Minimal **hepatic encephalopathy** (MHE) in children is difficult to evaluate due to ... In some **cases**, minimal or "subclinical" encephalopathy occurs, is also traditionally **used** as a **clinical marker of hepatic encephalopathy**.

[PDF] Hepatic Encephalopathy in Chronic Liver Disease ... - AA...

<https://www.aasld.org/sites/.../hepaticencephenhanced.pdf> ▾ 翻译此页

作者: H Vilstrup - 2014 - 被引用次数: 1 - 相关文章

2014年7月8日 - This document was designed for **use** on a variety patients with advanced liver disease, presenting in approximately 4% of **cases**.61 ... Overt **hepatic encephalopathy** (OHE) is diagnosed by **clinical** criteria and ... rater reliability and thus are chosen as **marker** symptoms of OHE 67 Orientation or mixed