Liver injury in the era of COVID-19 - Abstract - Europe PMC https://europepmc.org/article/PMC/PMC7856845 •

Another cohort study of 342 patients with COVID-19 revealed a relationship between **hepatic steatosis** and both transaminitis and increased disease severity. However, **steatosis** did not predispose to the development of clinically relevant liver insufficiency during the course of infection [75].

STAY (GI) HEALTHY: COVID-19 and Gastrointestinal ...

https://europepmc.org/article/MED/33521703 -

Jan 21, 2021 - Strikingly, GI symptoms may be the only clinical manifestation of COVID-19 or precede other symptoms: in a **study** of 1141 cases, 16% presented with GI symptoms as only chief complaint; 22 in another **study** analyzing 138 consecutive hospitalized **patients**, 14 (10%) **patients** presented with diarrhea and nausea 1-2 days before the development of fever ...

Author: Edoardo Vespa, Nicola Pugliese, Fran... Publish Year: 2021

STAY (GI) HEALTHY: COVID-19 and Gastrointestinal ...

https://europepmc.org/article/PMC/PMC7825983 •

Jan 21, 2021 - Europe PMC is an archive of life sciences journal literature. Introduction. The coronavirus SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2), which has infected more than 70 million people worldwide, is the cause of COVID-19 (coronavirus disease 2019), a condition including both asymptomatic or pauci-symptomatic forms and rapidly progressive deadly forms.

Author: Edoardo Vespa, Nicola Pugliese, Fran... Publish Year: 2021

Pathophysiology of COVID-19: Mechanisms Underlying \dots

https://journals.physiology.org/doi/full/10.1152/physiol.00019.2020

Aug 12, 2020 · The involvement of the gastrointestinal (GI) tract and hepatic system in COVID-19 disease progression is being increasingly reported. The most common GI manifestations reported in both adult and especially pediatric COVID-19 patients include diarrhea, nausea, vomiting, and abdominal pain (16, 133, 157).

Cited by: 9 Author: Mary Kathryn Bohn, Alexandra Hall, Lusi...
Publish Year: 2020

(1 2 3 4 5 >

Hepatic and gastrointestinal disturbances in Egyptian patients infe

∷ ↓



IMAGES VIDEOS ALL

12,100 Results Any time ▼

Coronavirus Disease-2019 (COVID-19) and the Liver

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8111098

Apr 28, 2021 · Within a year of its emergence, coronavirus disease-2019 (COVID-19) has evolved into a pandemic. What has emerged during the past 1 year is that, apart from its potentially fatal respiratory presentation from which the severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2) derives its name, it presents with a myriad of gastrointestinal (GI) and liver manifestations.

Cited by: 1 Author: Anshuman Elhence, Manas Vaishnav, Sagn...

Publish Year: 2021

Gastrointestinal manifestations in COVID-19

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7989191

Introduction. Coronavirus disease 2019 (COVID-19) is caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). It is currently a pandemic and as of 26 December 2020 there have been >79 million cases worldwide and >1.7 million deaths. 1 Several vaccines have been developed to control the pandemic. 2 The earliest record of coronavirus infections among animals was in the late 1920s ..


Author: Jayani C Kariyawasam, Umesh Jayarajah, ...

Publish Year: 2021

Coronavirus Disease (COVID-19) and Digestive System ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8289353

Aug 01, 2021 · Evidence for the GI infection in subjects with COVID-19. In addition to the detection of the viral presence in hronchoalyzolar fluid enutum, and blood enerimens from nationts with COVID-19, live





Hepatic and gastrointestinal disturbances in Egyptian patients infer





ALL

IMAGES

VIDEOS

12,000 Results

Any time ▼

Coronavirus Disease-2019 (COVID-19) and the Liver

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8111098

Apr 28, 2021 · Within a year of its emergence, coronavirus disease-2019 (COVID-19) has evolved into pandemic. What has emerged during the past 1 year is that, apart from its potentially fatal respiratory presentation from which the severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2) derives its name, it presents with a myriad of gastrointestinal (GI) and liver manifestations.

Cited by: 1

Author: Anshuman Elhence, Manas Vaishnav, Sa...

Publish Year: 2021

Gastrointestinal manifestations in COVID-19

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7989191

GI manifestations are reported in 11.4-61.1% of individuals with COVID-19, with variable onset and severity. The majority of COVID-19-associated GI symptoms are mild and self-limiting and include anorexia, diarrhoea, nausea, vomiting and abdominal pain/discomfort. A minority of patients present wit an acute abdomen with aetiologies such as ...

Cited by: 1

Author: Jayani C Kariyawasam, Umesh Jayarajah...

Publish Year: 2021

Clinical Characteristics of COVID-19 Patients With ...

https://pubmed.ncbi.nlm.nih.gov/32287140

Objective: Since the outbreak of Coronavirus Disease 2019 (COVID-19) in December 2019, various digestive symptoms have been frequently reported in patients infected with the virus. In this study, we aimed to further investigate the prevalence and outcomes of COVID-19 patients with digestive symptoms. Methods: In this descriptive, cross-sectional, multicenter study, we enrolled confirmed ...

Cited by: 958

Author: Lei Pan, Lei Pan, Mi Mu, Pengcheng Yan...

Publish Year: 2020

COVID-19-associated gastrointestinal and liver injury ...

https://www.nature.com/articles/s41392-020-00373-7

Nov 02, 2020 - Besides common respiratory symptoms, some patients with COVID-19 experience gastrointestinal symptoms, such as diarrhea, nausea, vomiting, and loss of appetite.

Cited by: 30

Author: Peijie Zhong, Jing Xu, Dong Yang, Yue Sh ...

Publish Year: 2020