

Name of Journal: *World Journal of Orthopedics*

ESPS Manuscript NO: 29148

Manuscript Type: Observational cohort study

Tourniquets do not increase the total blood loss or re-amputation risk in transtibial amputations.

Christian Wied¹, Peter T. Tengberg¹, Gitte Holm¹, Thomas Kallemose^{1,2}, Nicolai B. Foss³, Anders Troelsen¹, Morten T. Kristensen^{1,4}.

Match Overview

1	Internet 69 words crawled on 26-Sep-2016 pmrc.dk	2%
2	Internet 52 words crawled on 14-Jun-2016 www.rehab.research.va.gov	1%
3	Internet 49 words crawled on 30-Dec-2014 support.sas.com	1%
4	Crossref 38 words "Scientific surgery", British Journal of Surgery , 10/2006	1%
5	Internet 32 words crawled on 15-Nov-2012 143.104.52.229	1%
6	Crossref 24 words Pedowitz, R.A. "Muscle injury induced beneath and distal to a pneumatic tourniquet: A quantitative animal study ...	1%

学术搜索

获得 8 条结果 (用时0.10秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Dutch evidence-based guidelines for amputation and prosthetics of the lower extremity: Amputation surgery and postoperative management. Part 1

J Geertzen, H van der Linde... - Prosthetics and ..., 2014 - poi.sagepub.com

... 8. Evidence suggests that a TTA with a 'sagittal flap' **does not** give superior results compared with ... The use of a **tourniquet** in TTA is recommended because **tourniquet** use results in less ... of a single information dossier should be considered so that each discipline **can** see which ...

相关文章 所有 6 个版本 引用 保存

[PDF] ASSESSMENT OF THE OUTCOME OF LOWER LIMB AMPUTATIONS AS SEEN IN KENYATTA NATIONAL HOSPITAL.

DRKO AWORI - surgery.uonbi.ac.ke

... 50 While it gives an anatomical representation, it **does not** assess the functional obstruction ... produce skin necrosis through **tourniquet** effect, or are used once wound healing is assured. ... start walking early with the physiotherapist. These aids **can** assist in control of stump ...

相关文章 引用 保存 更多

12 Amputation

IM Nordon, RJ Hinchliffe, KG Jones - High Risk Diabetic Foot: ..., 2010 - books.google.com

... In the presence of a functioning infrainguinal bypass graft, it is still advisable to avoid **tourniquet** use. ... Synovial fluid leak **does not** appear to be a significant impediment to healing. ... Similarly, severe infection **can** lead to wound dehiscence requiring further surgical correction (94). ...

相关文章 引用 保存

13 Major Lower Extremity Amputation

TA Kimball, MR Nehler - Critical Limb Ischemia, 2016 - books.google.com

... **Not** surprising both studies found that pneumatic **tourniquet** use lowered absolute **blood loss** and patient transfusion requirements. It is our practice to use **tourniquets** for BKAs as long as

Scholar

8 results (0.15 sec)

Articles

Case law

My library

Any time

Since 2016

Since 2015

Since 2012

Custom range

Sort by relevance

Sort by date

☒ Include patents

☒ Include citations

☒ Create alert

Dutch evidence-based guidelines for amputation and prosthetics of the lower extremity: Amputation surgery and postoperative management. Part 1

J Geertzen, H van der Linde... - Prosthetics and ..., 2014 - pol.sagepub.com

... 8. Evidence suggests that a TTA with a 'sagittal flap' **does not** give superior results compared with ... The use of a **tourniquet** in TTA is recommended because **tourniquet** use results in less ... of a single information dossier should be considered so that each discipline **can** see which ...

Cited by 1 Related articles All 6 versions Cite Save

[PDF] ASSESSMENT OF THE OUTCOME OF LOWER LIMB AMPUTATIONS AS SEEN IN KENYATTA NATIONAL HOSPITAL

DRKO AWORI - surgery.uonbi.ac.ke

... He however, found out that majority of patients **do not** fit the absolute criteria yet a decision to salvage or amputate the limb has to be made. ... produce skin necrosis through **tourniquet** effect, or are used once wound healing is assured. ... These aids **can** assist in control of stump ...

Related articles Cite Save More

12 Amputation

IM Nordon, RJ Hinchliffe, KG Jones - High Risk Diabetic Foot: ..., 2010 - books.google.com

... are **not** warranted (level B). The converse is also valid, younger healthy patients **do** very well ... of a functioning infrainguinal bypass graft, it is still advisable to avoid **tourniquet** use. ... Efficient holistic care by the multidisciplinary team ensures that the patient **can** move on from this ...

Related articles Cite Save

13 Major Lower Extremity Amputation

TA Kimball, MR Nahler - Critical Limb Ischemia, 2016 - books.google.com

... **Not** surprising both studies found that pneumatic **tourniquet** use lowered absolute **blood loss** and patient ... It is our practice to use **tourniquets** for BKAs as long as the patient's arterial ... Thereafter the fasciocutaneous flaps **can** be reflected proximally to enable the remainder of the ...

Related articles Cite Save

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 1,730 条结果 (用时 0.88 秒)

Blood loss in total knee arthroplasty: an analysis of risk factors - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: N Prasad - 2007 - 被引用次数: 155 - 相关文章

The amount of blood loss in a primary cemented total knee arthroplasty (TKA) ... The post-operative transfusion trigger can be brought to 8.0 g% in a ... and less healthy, which increases the chance of complications during donation, ... We did not use a re-infusion drain or intra-operative cell savers during our study period.

缺少字词: transtibial amputations

Management of the mangled extremity

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: ML Prasarn - 2012 - 被引用次数: 12 - 相关文章

2012年6月13日 - In patients with complete traumatic disruption and clearly irreparable ... Although the decision to amputate in the acute setting is difficult for the patient, ... As stated above, loss of plantar sensation alone does not necessary indicate ... It has been shown that transtibial amputations have significantly better ...

Complications in Vascular and Endovascular Surgery: How to avoid ...

<https://books.google.com.hk/books?isbn=1908986808> - 翻译此页

Jonathan J Earnshaw,, Michael Wyatt, - 2012 - Medical

Antiplatelet agents do not need to be stopped perioperatively, as these rarely cause ... as well as increasing the risk of operative blood loss and the incidence of stump haematoma.... It is desirable to avoid the use of a tourniquet unless there is normal ... Flexion deformity in the limb to be amputated should be addressed ...