

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 55804

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Evaluation of intrahepatic manifestation and distant extrahepatic disease in alveolar echinococcosis

Tilmann Graeter, Hai-Hua Bao, Rong Shi, Wen-Ya Liu, Wei-Xia Li, Yi Jiang, Julian Schmidberger, Eleonore Brumpt, Eric Delabrousse, Wolfgang Kratzer; the XUUB consortium

Abstract

Match Overview

1	Internet 82 words crawled on 14-Jun-2020 www.wjgnet.com	2%
2	Crossref 37 words Tilmann Graeter, Haihua Bao, Eric Delabrousse, Eleonore Brumpt et al. "Hepatic alveolar echinococcosis: Comparative c...	1%
3	Internet 31 words crawled on 05-Dec-2019 www.edp-open.org	1%



ALL

IMAGES

VIDEOS

24,600 Results

Any time ▾

Innovation in hepatic alveolar echinococcosis imaging ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4273719>

Hepatic Alveolar Echinococcosis (HAE), caused by larvae of **Echinococcus multilocularis**, is a rare but potentially **lethal parasitic disease**. The first diagnostic suspicion is usually based on **hepatic ultrasound exam** performed because of abdominal symptoms or in the context of a general checkup; **HAE diagnosis** may thus also be an incidental finding on imaging.

Cited by: 28

Author: Wenya Liu, Éric Delabrousse, Oleg Blag...

Publish Year: 2014

Evaluation of hepatic cystic lesions - PubMed Central (PMC)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3691048>

Jun 21, 2013 · However, **echinococcosis** is considered to be an **emerging disease** in **Europe**[38,39]. Thus, CE and AE are **diseases** with a **considerable global disease impact**, as indicated by a substantial loss in **disability-adjusted** life years[38,40].

Cited by: 71

Author: Marten A Lantinga, Tom Jg Gevers, Joo...

Publish Year: 2013

Hepatic alveolar hydatid disease (Echinococcus ...

<https://www.sciencedirect.com/science/article/pii/S0009926019300613>

Alveolar hydatid disease or **alveolar echinococcosis (AE)** is caused by the **parasite Echinococcus multilocularis** and is increasingly seen as an **imported disease** in non-endemic areas such as the UK. It is rare compared to **cystic echinococcosis (CE)**, but like CE commonly affects the liver.

Author: M.D. Chouhan, M.D. Chouhan, E. Wil... **Publish Year:** 2019

Extrahepatic Primary Adrenal Alveolar Echinococcosis: A ...

https://www.researchgate.net/publication/249965566_Extrahepatic_Primary_Adrenal_al...

Alveolar echinococcosis (AE) primarily affects the liver and extrahepatic disease is considered



Evaluation of intrahepatic manifestation and distant extr.  

Sign in



ALL IMAGES VIDEOS

25,700 Results Any time ▾

Innovation in hepatic alveolar echinococcosis imaging ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4273719>

Hepatic **Alveolar Echinococcosis** (HAE), caused by larvae of *Echinococcus multilocularis*, is a rare but potentially lethal parasitic **disease**. The first diagnostic suspicion is usually based on hepatic ultrasound exam performed because of abdominal symptoms or in the context of a general checkup; HAE diagnosis may thus also be an incidental finding on imaging.

Cited by: 30 **Author:** Wenya Liu, Éric Delabrousse, Oleg Blago...
Publish Year: 2014

Alveolar Echinococcosis: Spectrum of Findings at Cross ...

<https://pubs.rsna.org/doi/10.1148/rg.327125708>

Nov 01, 2012 · **Alveolar echinococcosis** is a rare **parasitic disease** caused by the **fox tapeworm** *Echinococcus multilocularis*, which is endemic in many parts of the world. Without timely **diagnosis** and **therapy**, the pr...

Cited by: 106 **Author:** Mecit Kantarci, Ummugulsum Bayraktuta...
Publish Year: 2012

Alveolar Echinococcosis in Children

Search Tools

Turn off Hover Translation (关闭取词)



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

27,200 Results

Any time ▾

[Innovation in hepatic alveolar echinococcosis imaging ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4273719>

Hepatic **Alveolar Echinococcosis** (HAE), caused by larvae of *Echinococcus multilocularis*, is a rare but potentially lethal parasitic **disease**. The first diagnostic suspicion is usually based on hepatic ultrasound exam performed because of abdominal symptoms or in the context of a general checkup; HAE diagnosis may thus also be an incidental finding on imaging.

Cited by: 30

Author: Wenya Liu, Éric Delabrousse, Oleg Blago...

Publish Year: 2014

[Alveolar echinococcosis-spreading disease challenging ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3710431>

Jul 14, 2013 · INTRODUCTION. The fox-tapeworm *Echinococcus multilocularis* (*E. multilocularis*) is the causative agent of **alveolar echinococcosis** (AE), a potentially deadly parasitic **disease**. AE is prevalent in the northern hemisphere and central Europe is an endemic focus[1-3]. In the era prior to anthelmintic treatment the cumulative lethality for AE was about 90% ten years after a diagnosis has been ...

Cited by: 11

Author: Georgi Atanasov, Christoph Benckert, Ar...

Publish Year: 2013

[Multimodality imaging in diagnosis and management of ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4859741>

Epidemiology and pathophysiology. In humans, the *E. granulosus* cestodes causes cystic **echinococcosis** (CE), *E. multilocularis* cestodes causes **alveolar echinococcosis** (AE), and *E. vogeli* and *E. oligarthrus* cause polycystic **echinococcosis** (also termed as neotropical **echinococcosis**). Although CE is common worldwide, AE is a **manifestation** seen only in the northern hemisphere.

Cited by: 18

Author: Mesut Bulakçı, Merve Gülbiz Kartal, Sabr...

Publish Year: 2016

[Alveolar echinococcosis: Evaluation of therapeutic strategies](#)

<https://www.researchgate.net/publication/262795896...>

Cystic **echinococcosis** (CE) and **alveolar echinococcosis** (AE) are chronic, complex and neglected **diseases**. Their treatment depends on a number of factors related to the lesion, setting and patient.