

Name of journal: *World Journal of Clinical Infectious Diseases*

ESPS Manuscript NO: 17056

Columns: Original article

Basic Study

Tuf mRNA rather than 16S rRNA is associated with culturable *Staphylococcus aureus*

Anne JM Loonen, Petra FG Wolffs, Maikel de Bresser, Maurice Habraken, Cathrien A Bruggeman, Mirjam HA Hermans, Adriaan JC van den Brule

Abstract

AIM: To study the presence of various nucleic acids targets of *Staphylococcus aureus* during bacterial growth and antibiotic induced killing in relation to viability.

METHODS: *S. aureus* was cultured to log phase and spiked in Todd-Hewitt (TH)

Match Overview

1

Internet 20 words
crawled on 25-Mar-2010
www.epa.gov

<1%

2

CrossCheck 19 words
L. Birch. "A comparison of nucleic acid amplification techniques for the assessment of bacterial viability", Letters in Applied

<1%

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 43,100 条结果 (用时 0.59 秒)

Google 学术: Tuf mRNA rather than 16S rRNA is associated with culturable Staphylococcus aureus

Specific metabolic activity of ripening bacteria ... - [Falentín](#) - 被引用次数: 36

... Phytoplasma australiense"(subgroup tuf-Australia I; rp- ... - [Tran-Nguyen](#) - 被引用次数: 122

Diagnostic microbial microarrays in soil ecology - [Sessitsch](#) - 被引用次数: 73

Current Perspectives on Viable but Non-Culturable (VBNC ...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: T Ramamurthy - 2014 - 被引用次数: 7 - [相关文章](#)

2014年7月31日 - In VBNC cells, the expression levels of ribosome-associated proteins such ... The VBNC *Staphylococcus aureus* displayed mutational inactivation of ... on agar plates, can recover their ability if plated on matrices other than agar. ... state, 16S rRNA and mRNA for tuf, rpoS, and relA have been targeted (77).

Sensitive Quantitative Detection of Commensal Bacteria by ...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: K Matsuda - 2007 - 被引用次数: 146 - [相关文章](#)

2006年10月27日 - Using group- or species-specific primers for 16S or 23S rRNA, analytical ... in good agreement with the RT-qPCR counts rather than with the CFU

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 37,700 条结果 (用时 0.73 秒)

Google 学术 : Tuf mRNA rather than 16S rRNA is associated with culturable Staphylococcus aureus

Specific metabolic activity of ripening bacteria ... - [Falentin](#) - 被引用次数 : 36

... Phytoplasma australiense"(subgroup [tuf-Australia I](#); rp- ... - [Tran-Nguyen](#) - 被引用次数 : 127

Diagnostic microbial microarrays in soil ecology - [Sessitsch](#) - 被引用次数 : 74

Current Perspectives on Viable but Non-Culturable (VBNC ...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者 : T Ramamurthy - 2014 - 被引用次数 : 7 - [相关文章](#)

2014年7月31日 - In VBNC cells, the expression levels of ribosome-associated proteins such ... The VBNC *Staphylococcus aureus* displayed mutational inactivation of on agar plates, can recover their ability if plated on matrices other than agar. ... state, 16S rRNA and mRNA for tuf, rpoS, and relA have been targeted (77).

[PDF] Download as PDF - InTech

www.intechopen.com/download/pdf/37259 ▾ [翻译此页](#)

2012年5月30日 - involved in the generation of the functional properties. In addition, the abundance of specific mRNA transcripts can be quantified by reverse transcription