

Long-term rare giant sialolithiasis for 30 years: A case report and review of literature

Jit-Swen Mao, Yu-Chien Lee, Jessie Chao-Yun Chi, Wan-Ling Yi, Yung-An Tsou, Chia-Der Lin, Chih-Jaan Tai, Liang-Chun Shih

Specialty type: Medicine, research and experimental

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Corvino A, Italy;
Ozcan C, Turkey

Received: May 4, 2023

Peer-review started: May 4, 2023

First decision: May 25, 2023

Revised: June 1, 2023

Accepted: July 6, 2023

Article in press: July 6, 2023

Published online: August 6, 2023



Jit-Swen Mao, Yu-Chien Lee, Wan-Ling Yi, Yung-An Tsou, Chia-Der Lin, Chih-Jaan Tai, Liang-Chun Shih, Department of Otorhinolaryngology-Head and Neck Surgery, China Medical University Hospital, Taichung 400, Taiwan

Jessie Chao-Yun Chi, Department of Otorhinolaryngology, Head and Neck Surgery, Taichung Hospital, Ministry of Health and Welfare, Taichung 404, Taiwan

Yung-An Tsou, Liang-Chun Shih, Department of Otorhinolaryngology-Head and Neck Surgery, Asia University Hospital, Taichung 404, Taiwan

Yung-An Tsou, Chia-Der Lin, Chih-Jaan Tai, Liang-Chun Shih, School of Medicine, China Medical University, Taichung 404, Taiwan

Corresponding author: Liang-Chun Shih, Department of Otorhinolaryngology-Head and Neck Surgery, China Medical University Hospital, No. 2 Yuh-Der Road, Taichung 400, Taiwan. entdrshih7111@gmail.com

Abstract

BACKGROUND

Sialolithiasis is one of the most common salivary gland disorders, most commonly affecting the submandibular gland. Submandibular sialolithiasis can be treated using non-invasive conservative measures and invasive treatments. Treatment selection was based on the ductal system anatomy and the size and location of the stones. This study aimed to review the updates on sialolithiasis treatment and compare the different management strategies of the variables.

CASE SUMMARY

This report presents a case of a long-term, rare, and giant sialolithiasis within the submandibular gland parenchyma for 30 years in an older adult. Our patient presented with painless right submandibular swelling. Computed tomography revealed a calcified mass measuring 35 mm × 20 mm within the right submandibular gland. In this case, the infection and fibrosis of the affected gland and size of the stone did not provide us with other alternatives except for the excision of the involved gland. Thus, right submandibular sialoadenectomy was performed *via* the transcervical approach. After the surgery, the patient recovered without any complaints, side effects, or complications.

CONCLUSION

Tailored management is important for preserving gland function, maintaining low risk, and reducing patient discomfort.

Key Words: Huge sialolithiasis; Submandibular gland; Treatment; Complications; Case report

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Giant sialoliths are uncommon, and their long duration is rarely documented. This report presents a case of a long-term, rare, and giant sialolithiasis within the submandibular gland parenchyma for 30 years in an older adult. We reviewed 44 cases of giant sialoliths that were reported between 1981 and 2022. Based on the comparative table, our case is the second largest with the longest duration. This case highlights the importance of proper diagnosis and treatment.

Citation: Mao JS, Lee YC, Chi JCY, Yi WL, Tsou YA, Lin CD, Tai CJ, Shih LC. Long-term rare giant sialolithiasis for 30 years: A case report and review of literature. *World J Clin Cases* 2023; 11(22): 5382-5390

URL: <https://www.wjgnet.com/2307-8960/full/v11/i22/5382.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v11.i22.5382>

INTRODUCTION

Sialoliths are calcified salivary stones that can form within the secretory system of the salivary glands. Although various hypotheses have been proposed, the exact etiology of these sialoliths remains unclear. Most patients with sialolithiasis present with symptoms such as local pain, focal swelling, trismus, xerostomia, and lumps in the throat. Inflammation with oozing pus may be noted in some cases due to secondary infection.

The submandibular gland is the most commonly affected site for sialoliths, accounting for approximately 80%-90% of cases[1]. The rate of giant sialolith formation within the submandibular gland is high, and several contributing factors have been proposed, including a longer Wharton's duct, larger duct caliber, and tortuous course of the Wharton's duct accompanied by a slower salivary flow rate within the submandibular gland compared with that in other salivary ducts. The slower salivary flow rate is caused by saliva in the submandibular glands flowing against gravity, and the dependent position of the gland makes it more prone to stasis. Additionally, owing to the higher calcium and phosphate levels, the saliva within the submandibular glands tends to be more alkaline and may induce sialolithiasis[2].

Sialoliths usually range from 1 mm to 10 mm in size, and those measuring 15 mm in one dimension are classified as giant sialoliths[3]. In this report, we present a case of giant right submandibular sialolithiasis that was treated by surgical ablation for approximately 30 years and compare various giant sialolith cases and their different treatments with respect to variables such as size, duration, and symptom presentation.

CASE PRESENTATION

Chief complaints

A 75-year-old woman presented to our institution with painless right submandibular swelling with a persistence of 30 years.

History of present illness

The clinical presentation was mostly asymptomatic, with only slight facial asymmetry and occasional right-sided aural fullness. However, the indurated mass began to manifest focal swelling 1 mo prior, with intermittent severe pain that radiated to the ipsilateral neck and contralateral face. The additional onset of dysphagia, odynophagia, and a subsequent decrease in appetite were also noted but without associated postprandial swelling or xerostomia.

History of past illness

The patient had no known medical history.

Personal and family history

The patient denied alcohol or cigarette use and confirmed a family history of sialolithiasis.

Physical examination

Bimanual palpation revealed a raised right erythematous submandibular mass measuring 40 mm × 30 mm overlying the skin. The swelling was indurated, tender on palpation, and firm in consistency. No other obvious cervical lymphadenopathies could be palpated. Intraorally, no stones or purulent discharge was secreted from the Wharton's duct.

Laboratory examinations

White blood cells count levels were elevated (13800/ μ l).

Imaging examinations

Computed tomography (CT) revealed a calcified mass measuring 35 mm \times 20 mm within the right submandibular gland (Figure 1).

FINAL DIAGNOSIS

Combined with the patient's medical history, the final diagnosis was submandibular sialolithiasis.

TREATMENT

Excision of the right submandibular gland and sialoliths was performed using a standard extraoral approach. Intraoperative findings revealed that the right submandibular gland was indurated and filled with giant sialoliths. Neighboring structures, such as the hypoglossal, lingual, and marginal mandibular nerves, were preserved (Figure 2).

OUTCOME AND FOLLOW-UP

The postoperative was uneventful, with no complications. Histopathological examination confirmed the diagnoses of sialolithiasis and chronic sialadenitis. A giant, yellow, hard sialolith measuring 25 mm \times 17 mm \times 17 mm was noted. Microscopically, the submandibular gland showed varied dilatations of the salivary ducts (ductal ectasia). Ulcerative changes in the dilated ductal epithelium with areas of squamous metaplasia, periductal fibrosis, and moderately mixed acute and chronic inflammatory infiltrates with lymphocytic aggregate foci were also observed.

DISCUSSION


Several treatment modalities for submandibular sialolithiasis exist; however, the primary goal is to preserve gland function, maintain low risk, and reduce patient discomfort[4]. Non-invasive conservative measures for submandibular sialolithiasis include glandular focal massage and trans-oral removal. Glandular focal massage is important during or after the administration of sialogogues, and irrigation is primarily indicated in cases of small intraductal stones. Transoral removal can be considered if stones are palpated bimanually and/or localized within the perihilar region of the gland using ultrasonography[4,5]. If a superimposed infection is suspected, antibiotics should be prescribed and, when warranted, combined with simple sialolithotomy[5,6].

Invasive management of submandibular sialolithiasis consists of open surgery (which can be performed with transoral duct surgery[7-9] or a transcervical approach to the submandibular space[10], extracorporeal shock wave lithotripsy[11-15], interventional sialendoscopy, and intraductal shock-wave lithotripsy[16-19]. These invasive methods are recommended for stones > 7 mm with symptoms[20]. Treatment selection is mostly based on ductal system anatomy and the location of the stones[7].

Transoral duct surgery may be considered for stones of all sizes in the papilla, owing to the narrow ostium of the papilla and stones in the distal and middle ducts. Small mobile stones (not exceeding 3-5 mm) in the papilla distal and middle ducts can also be extracted using interventional sialendoscopy. The proximal duct, however, is curved and narrow; thus, interventional sialendoscopy is difficult to perform. In cases where the stones are immobile, impacted, palpable, or large, transoral duct surgery is recommended, although mechanical fragmentation or intraductal shockwave lithotripsy followed by fragment extraction is also an option. Intraparenchymal stones are rare; however, if small, mobile stones (not exceeding 3-5 mm) can be visualized during endoscopy, interventional sialendoscopy, including mechanical fragmentation or intraductal shock wave lithotripsy, is recommended. Transoral duct surgery with submandibulectomy is indicated for large palpable stones (10 mm) located posthilar to the intraparenchymal region. Finally, salivary gland massage combined with a sour diet and sialogogue use is recommended after the removal of salivary stones[5].

In summary, transoral duct surgery offers better accessibility to the duct system and remains the preferred treatment modality; however, intraductal shock wave lithotripsy has gradually replaced transoral duct surgery in cases of smaller salivary stones. Compared with an earlier algorithm for the management of submandibular stones published in 2009, the indications for extracorporeal shockwave lithotripsy have since been adjusted, and the newly developed intraductal shockwave lithotripsy is also available. Although extracorporeal shockwave lithotripsy is considered less important, the procedure remains an option for intraparenchymal stones that are not visible or accessible. Finally, botulinum toxin injections may be considered in situations where the patient is deemed inoperable[7].

Among all the invasive treatment methods, adenectomy has a higher success rate than other techniques. Open surgery via the transcervical approach has a lower risk of infection because the oral space does not communicate with the neck space, whereas the intraoral approach avoids facial scarring. However, open surgical approaches are associated with



DOI: 10.12998/wjcc.v11.i22.5382 Copyright ©The Author(s) 2023.

Figure 1 Axial view of the head and neck computed tomography with contrast showing an irregular opacity of 35 mm x 20 mm in the parenchyma of the right submandibular gland.

several complications. Examples include postoperative neurological damage, particularly hypoglossal, lingual, and marginal mandibular nerve injuries, hemorrhage, hematoma, and disfiguring scars. Temporary marginal mandibular nerve damage is the most common complication of transcervical submandibular gland surgery. This complication was reported to be 36% by Smith *et al*[21], and permanent damage was reported to be as high as 12% by De *et al*[22]. In the literature, Beahm *et al*[23] reported rates between 7.7% and 36% in a previous literature review. To reduce the risk of marginal mandibular nerve injury in the transcervical surgical approach to the submandibular gland, three surgical maneuvers are recommended: (1) The gland was identified to be lower than the hyoid bone, and the marginal mandibular nerve was not examined; (2) The marginal mandibular nerve is located at the level at which it leaves, tracks, and protects the parotid tail; and (3) The facial veins are located at a lower level and ligated, and the nerve is suspended by elevation; thus, the marginal mandibular nerve is removed from the surgical area.

However, Smith *et al*[21] reported that when a lower approach was used, no permanent marginal mandibular nerve palsy occurred, although 36% of the nerves were temporarily dysfunctional. This may have been due to stretching during the lower surgical approach to the glands. The hypoglossal nerve is partially sheltered by the posterior belly of the digastric muscle, which is located in the inferomedial portion of the lower one-third of the submandibular glands. This may explain why hypoglossal nerve damage has been reported very rarely (0%-1.4%)[24] compared with marginal mandibular nerve injuries. In this case report, no permanent or temporary complications occurred during the postoperative follow-up. Although the submandibular glands contribute 69% of the salivary secretion volume in the resting state[25], xerostomia was not reported in our case. This could be explained by the fact that other salivary glands were sufficiently functional to compensate for the resected glands.

Moreover, regarding the oncological risk to which patients with chronic inflammation and lithiasis are exposed[26], we reviewed some studies on salivary gland tumors. Most salivary gland tumors arise in the parotid gland (70%), followed by the minor salivary glands (22%) and submandibular glands (8%)[27]. Differential diagnoses for benign submandibular tumors include pleomorphic adenomas, Warthin's tumors, and oncocytomas. Histopathological characteristics can be used to distinguish between them. Pleomorphic adenomas are characterized by thick irregularly marginated capsules[28, 29]. Oncocytomas have thin capsules and monomorphic oncocytes without mitosis or necrosis[29,30]. The lymphatic population is the most common cytology and histology of Warthin's tumor[28,29]. Histopathological examination of our patient did not reveal the above characteristics, and the characteristics of malignant transformation include local invasion into the muscular, perineural, and lymphatic structures, as well as microscopic features, including nuclear atypia, cellular polymorphism, mitoses, and focal necrosis, were absent too. Thus, salivary gland tumors were excluded.

Owing to the similarities in the clinical presentation of benign salivary tumors and submandibular sialolithiasis, radiological imaging is essential to distinguish between the two entities. Ultrasound is recommended for the initial assessment of salivary gland abnormalities; however, ultrasound is insufficient because information on the surrounding structures cannot be provided. Its accuracy in the identification of benign salivary lesions can be increased using elastography or contrast-enhanced ultrasound; however, the European Federation of Societies for Ultrasound in Medicine and Biology do not recommend contrast-enhanced ultrasound for the characterization of salivary gland lesions in clinical practice. Thus, further studies are required to investigate the diagnostic roles of contrast-enhanced ultrasound and elastography in the evaluation of salivary gland lesions. Corvino *et al*[26] described the utility of CT and magnetic resonance imaging in the initial staging, histologic grading of salivary gland malignancies, and preoperative planning. Positron emission tomography with F-18 fluorodeoxyglucose is useful for the evaluation and clinical management of head and neck lesions[26].

Giant sialoliths are rare and classified as those measuring 15 mm in one dimension[3]. We compared cases of giant sialoliths in the submandibular gland from 1981 to 2022 and their different treatments according to variables such as size, duration, and symptom presentation. The cases and their details are listed in Table 1.

Table 1 Comparative table of various cases of giant submandibular sialolithiasis with different treatments

Ref.	Year	Size (mm)	Symptoms/signs	Duration	Treatment	Complication
Zakaria[35]	1981	33	Swelling in the right submandibular region, increasing pain during meals	20 yr	The right submandibular gland was excised	None
Naraynsingh[36]	1985	60	Painless hard left submandibular swelling	3 yr	Incision of left submandibular gland <i>via</i> a transcervical approach	None
Frame and Smith [37]	1986	30	Recurrent pain and swelling of the right side of the face	N/A	Excise the gland and calculus	None
		25	Slight discomfort in the area	N/A	Sialo-adenectomy <i>via</i> a standard incision	None
Kaltman and Eichner[38]	1987	45	Pain and swelling over the left submandibular area	15 yr	Standard extraoral approach	None
Raveenthiran and Hayavadana Rao [39]	2004	35	Painless swelling in the floor of her mouth	N/A	Transoral sialolithotomy	None
Yildirim[40]	2004	30	Huge, firm mass below the right angle of the mandible	N/A	Excised <i>via</i> extraoral approach	None
Chan and Patel[41]	2006	35	Painful mass in the left floor of the mouth	N/A	Transoral sialodochoplasty and removal calculus	None
Graziani <i>et al</i> [42]	2006	22	Asymptomatic	N/A	Transorally	None
Ledesma-Montes <i>et al</i> [43]	2007	36	Painful mass located on the right side of the floor of the mouth	12 yr	Surgical excision of the sialolith	None
Biddle and Arora [44]	2008	26	Persistent draining wound over the right submandibular region, odynophagia, and dysphagia were noted	2 yr	Excised <i>via</i> standard incision	None
Rai and Burman[45]	2009	72	Episodes of pain for last 2 yr, but severe pain and swelling in the left lower submandibular region	1 mo	Sialolithotomy with sialodochoplasty <i>via</i> an intraoral approach	None
Soares <i>et al</i> [46]	2009	25	Hard elongated mass along the right Wharton's duct and a reduced salivary flow	4 mo	Dissected with the patient under local anesthesia	None
Rivera-Serrano and Schaitkin[47]	2011	23	Chronic unrelenting purulent discharge from bilateral submandibular ducts	N/A	Transoral sialolithotomy approach	None
Krishnan <i>et al</i> [48]	2009	34	Recurrent pain and swelling	8 yr	Transoral approach	None
Emir <i>et al</i> [49]	2010	35	Mass on the right side of the floor of the mouth and fistulization of the Wharton duct into the oral cavity	N/A	Removed transorally from the opening of the fistula at the floor of the mouth	None
Shetty and Sharma [50]	2010	27	Firm painful swelling in the left submandibular region	1 mo	Gentle extraction	None
Boffano and Gallesio[51]	2010	22	Swelling and pain in the submandibular right region	N/A	Removal of the stone and sialodochoplasty were done <i>via</i> an intraoral approach	None
Arunkumar <i>et al</i> [52]	2015	20	Recurrent right submandibular swelling with pain during meals and gradual regression	8 mo	Removed intraorally	None
Pandarakalam <i>et al</i> [53]	2013	40	Swelling, gradually increased in size and became painful recently	4 yr	Excision of the right submandibular salivary gland	None
Iqbal <i>et al</i> [54]	2012	35	Hard swelling in the anterior oral floor of mouth	N/A	Intraoral incision	None
Ben-Shoshan and Lacroix[31]	2014	17	Mild pain in the submandibular area	1 yr	Fell out without any intervention	None
Gupta <i>et al</i> [3]	2013	28	Intermittent, dull aching pain, and swelling in left submandibular area	3 mo	Intraoral approach	None
Bhullar <i>et al</i> [55]	2015	31	Pain in the left submandibular region associated with meals	6 mo	Transoral sialolithotomy	None
Akinyamoju and Adisa[32]	2015	44	Hard, painless, left floor of mouth swelling, moderate pain at mealtimes	1 yr	Removed non-surgically	None

Shahoon <i>et al</i> [56]	2015	55	Difficulty breathing and a foul-tasting mouth since 21 days previously with history of right submandibular swelling episodes	21 d	Surgical removal	None
Arslan <i>et al</i> [57]	2015	35	Recurrent pain, swelling, and redness in the left submandibular area	6 mo	Submandibular gland resection	None
Oliveira <i>et al</i> [58]	2016	30	Pain over the right floor of mouth and submandibular region	N/A	Excision of the right submandibular gland and stone <i>via</i> standard extraoral approach	None
Weinberg and Albers[59]	2016	38	Swelling at the submandibular region, pus from Wharton's duct was noted	N/A	Excision	None
Gadve <i>et al</i> [60]	2016	25	Pain and swelling in the floor of mouth on the left side	1 mo	Remove the sialolith surgically	None
Goh <i>et al</i> [61]	2016	35	Left submandibular swelling, increasing pain, fever and discharge was noted	5 yr	Submandibulectomy with transcervical approach	None
Omezli <i>et al</i> [62]	2016	37	Painful and persistent purulent discharge from unilateral submandibular ducts	N/A	Intraoral approach	None
Lim <i>et al</i> [63]	2017	50	Right submandibular swelling, increased in size rapidly causing discomfort and pain	10 yr	Right submandibulectomy	None
Iwai <i>et al</i> [64]	2017	41	Right submandibular swelling several times	40 yr	Submandibular sialoadenectomy	None
Sakthivel <i>et al</i> [65]	2017	50	Recurrent pain and swelling in the right submandibular region	6 mo	Intraoral removal of stone	None
Rodrigues <i>et al</i> [33]	2017	45	Painless mass in the neck	3 yr	Removed without anesthesia or surgical excision	None
Singh <i>et al</i> [66]	2020	42	Swelling over right side of floor of mouth and pain during meals	2 yr	Sialolithotomy and marsupialization of Wharton's duct	None
Abraham <i>et al</i> [67]	2021	40	Hard mass beneath the tongue with occasional dull pain	N/A	Transoral surgical removal of the right submandibular calculus	None
Thong <i>et al</i> [68]	2021	25	Right submandibular swelling	2 wk	Excision of right submandibular gland	None
Kumar <i>et al</i> [69]	2021	28	Firm mass below the left angle of the mandible	4 yr	Excision of the left submandibular gland and stone <i>via</i> trans-cervical approach	None
Brooks <i>et al</i> [34]	2021	16&19	Firm, ovoid swelling in the floor of the mouth approximate to the left sublingual caruncle with thin, slightly pale overlying mucosa	N/A	Patient declined any additional diagnostic measures	None
Ungari <i>et al</i> [70]	2022	23	Painful swelling in the left side of the mouth floor	20 yr	Excision of the left submandibular gland and stone <i>via</i> trans-cervical approach	None
Mohsin <i>et al</i> [71]	2022	28	Moveable, firm, extensive non-tender swelling in the posterior floor of the mouth	15 yr	Incision of the right submandibular duct's longitudinal axis and removed the stones	None

Between 1981 and 2022, 44 giant sialoliths were noted. The mean giant sialoliths size is 34.70 mm (two cases were within the range of 15-19 mm; 14 cases within 20-29 mm; 15 cases within 30-39 mm; seven cases within 40-49 mm; three cases within 50-59 mm; one case within 60-69 mm and one case within the range 70-79 mm). The median duration of giant sialoliths was 2 years (11 cases were less than a year; 9 cases were within the range of 1-5 years; two cases were within 6-10 years; three cases were within the range of 11-15 years; two cases were within the range of 16-20 years; one case was within the range of 36-40 years; and 15 cases did not discuss the duration). Of the 43 tumors, 23 and 21 were located in the right and left submandibular glands, respectively. Of the giant sialoliths, 90.70% were surgically removed (incision or excision), either intra-orally or extra-orally (transcervical), whereas 9.30% were removed using other methods. The patients with giant sialoliths reported by Ben-Shoshan and Lacroix[31] did not undergo any intervention. Akinyamaju and Adisa[32] and Rodrigues *et al*[33] removed the stones non-surgically. Brooks *et al*[34] reported that a patient declined any additional diagnostic measures and was lost at follow-up. No postoperative complications were observed in these cases.

CONCLUSION

Giant sialoliths are rare, and their longer durations have rarely been documented. This report presents a case of long-term, rare, and giant sialolithiasis within the submandibular gland parenchyma for 30 years in an older adult. Our patient



Figure 2 Excised sialolith and right submandibular gland.

presented with typical clinical and radiographic findings. In this case, the infection and fibrosis of the affected gland and size of the stone provided us with no other alternatives except for the excision of the involved gland. Thus, right submandibular sialoadenectomy was performed *via* the transcervical approach. After the surgery, the patient recovered smoothly without any complaints, side effects, or complications. Based on the comparative table, this is the second largest case with a long duration that we would like to share. This case highlights the importance of proper diagnosis and treatment. Clinicians are advised to tailor management according to the patient and to keep in mind that the primary goal should always be to preserve gland function, maintain low risk, and reduce patient discomfort.

FOOTNOTES

Author contributions: Mao JS and Lee YC conceived the study and contributed to protocol development, data collection, data analysis, interpretation of results, and drafting of the article; Mao JS and Lee YC collected the data; Chi JCY, Yi WL, Tsou YA, Lin CD, Tai CJ, and Shih LC provided critical feedback, direction, and article revisions; Mao JC, Lee YC, and Shih LC contributed to the protocol development, interpreted the results, and finalized the manuscript; all authors have approved the final version of the manuscript submitted for publication; Mao JS and Lee YC contributed equally to this study.

Supported by The China Medical University Hospital, No. DMR-110-242 and No. DMR-110-057.

Informed consent statement: All study participants or their legal guardians provided written informed consent prior to study enrollment.

Conflict-of-interest statement: The authors declare no conflict of interest.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Taiwan

ORCID number: Jit-Swen Mao [0009-0002-1919-9278](https://orcid.org/0009-0002-1919-9278); Yu-Chien Lee [0009-0002-8140-9811](https://orcid.org/0009-0002-8140-9811); Yung-An Tsou [0000-0002-8698-069X](https://orcid.org/0000-0002-8698-069X); Liang-Chun Shih [0000-0002-3036-6156](https://orcid.org/0000-0002-3036-6156).

S-Editor: Lin C

L-Editor: A

P-Editor: Zhao S

REFERENCES

- 1 McKenna JP, Bostock DJ, McMenemy PG. Sialolithiasis. *Am Fam Physician* 1987; **36**: 119-125 [PMID: [3318353](https://pubmed.ncbi.nlm.nih.gov/3318353/)]
- 2 Burket LW, Greenberg MS, Glick M, Ship JA. *Burket's oral medicine*. 11th ed. United States: PMPH USA, 2008: 201-202

- 3 **Gupta A**, Rattan D, Gupta R. Giant sialoliths of submandibular gland duct: Report of two cases with unusual shape. *Contemp Clin Dent* 2013; **4**: 78-80 [PMID: 23853458 DOI: 10.4103/0976-237X.111599]
- 4 **Zenk J**, Constantinidis J, Al-Kadah B, Iro H. Transoral removal of submandibular stones. *Arch Otolaryngol Head Neck Surg* 2001; **127**: 432-436 [PMID: 11296054 DOI: 10.1001/archotol.127.4.432]
- 5 **Zenk J**, Gottwald F, Bozzato A, Iro H. [Submandibular sialoliths. Stone removal with organ preservation]. *HNO* 2005; **53**: 243-249 [PMID: 15184990 DOI: 10.1007/s00106-004-1110-4]
- 6 **Williams MF**. Sialolithiasis. *Otolaryngol Clin North Am* 1999; **32**: 819-834 [PMID: 10477789 DOI: 10.1016/s0030-6665(05)70175-4]
- 7 **Koch M**, Mantsopoulos K, Müller S, Sievert M, Iro H. Treatment of Sialolithiasis: What Has Changed? An Update of the Treatment Algorithms and a Review of the Literature. *J Clin Med* 2021; **11** [PMID: 35011971 DOI: 10.3390/jcm11010231]
- 8 **McGurk M**, Makdissi J, Brown JE. Intra-oral removal of stones from the hilum of the submandibular gland: report of technique and morbidity. *Int J Oral Maxillofac Surg* 2004; **33**: 683-686 [PMID: 15337182 DOI: 10.1016/j.ijom.2004.01.024]
- 9 **Capaccio P**, Torretta S, Ottaviani F, Sambataro G, Pignataro L. Modern management of obstructive salivary diseases. *Acta Otorhinolaryngol Ital* 2007; **27**: 161-172 [PMID: 17957846]
- 10 **Goh YH**, Sethi DS. Submandibular gland excision: a five-year review. *J Laryngol Otol* 1998; **112**: 269-273 [PMID: 9624377 DOI: 10.1017/s0022215100158323]
- 11 **Iro H**, Zenk J, Waldfahrer F, Benzel W, Schneider T, Eil C. Extracorporeal shock wave lithotripsy of parotid stones. Results of a prospective clinical trial. *Ann Otol Rhinol Laryngol* 1998; **107**: 860-864 [PMID: 9794616 DOI: 10.1177/000348949810701009]
- 12 **Capaccio P**, Ottaviani F, Manzo R, Schindler A, Cesana B. Extracorporeal lithotripsy for salivary calculi: a long-term clinical experience. *Laryngoscope* 2004; **114**: 1069-1073 [PMID: 15179215 DOI: 10.1097/00005537-200406000-00021]
- 13 **Escudier MP**, Brown JE, Drage NA, McGurk M. Extracorporeal shockwave lithotripsy in the management of salivary calculi. *Br J Surg* 2003; **90**: 482-485 [PMID: 12673752 DOI: 10.1002/bjs.4067]
- 14 **Zenk J**, Bozzato A, Winter M, Gottwald F, Iro H. Extracorporeal shock wave lithotripsy of submandibular stones: evaluation after 10 years. *Ann Otol Rhinol Laryngol* 2004; **113**: 378-383 [PMID: 15174765 DOI: 10.1177/000348940411300507]
- 15 **Katz P**. [New techniques for the treatment of salivary lithiasis: sialoendoscopy and extracorporeal lithotripsy: 1773 cases]. *Ann Otolaryngol Chir Cervicofac* 2004; **121**: 123-132 [PMID: 15223997 DOI: 10.1016/s0003-438x(04)95499-0]
- 16 **Marchal F**, Dulguerov P, Becker M, Barki G, Disant F, Lehmann W. Specificity of parotid sialendoscopy. *Laryngoscope* 2001; **111**: 264-271 [PMID: 11210873 DOI: 10.1097/00005537-200102000-00015]
- 17 **Marchal F**, Dulguerov P, Becker M, Barki G, Disant F, Lehmann W. Submandibular diagnostic and interventional sialendoscopy: new procedure for ductal disorders. *Ann Otol Rhinol Laryngol* 2002; **111**: 27-35 [PMID: 11800367 DOI: 10.1177/000348940211100105]
- 18 **Nahlieli O**, Shacham R, Bar T, Eliav E. Endoscopic mechanical retrieval of sialoliths. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2003; **95**: 396-402 [PMID: 12686923 DOI: 10.1067/moe.2003.145]
- 19 **Koch M**, Zenk J, Iro H. [Diagnostic and interventional sialoscopy in obstructive diseases of the salivary glands]. *HNO* 2008; **56**: 139-144 [PMID: 17676291 DOI: 10.1007/s00106-007-1563-3]
- 20 **Faizal B**, Gangadharan S, Thankappan K. Comparison between Sialendoscopy and Conventional Methods in the Treatment of Sialolithiasis. *Malays J Med Sci* 2017; **24**: 94-100 [PMID: 29386976 DOI: 10.21315/mjms2017.24.5.10]
- 21 **Smith WP**, Peters WJ, Markus AF. Submandibular gland surgery: an audit of clinical findings, pathology and postoperative morbidity. *Ann R Coll Surg Engl* 1993; **75**: 164-167 [PMID: 8323209]
- 22 **De M**, Singh PK, Johnson AP. Morbidity associated with submandibular gland excision: a retrospective analysis. *Internet J Anesthesiol* 2006; **1** [DOI: 10.5580/173d]
- 23 **Beahm DD**, Peleaz L, Nuss DW, Schaitkin B, Sedlmayr JC, Rivera-Serrano CM, Zanation AM, Walvekar RR. Surgical approaches to the submandibular gland: a review of literature. *Int J Surg* 2009; **7**: 503-509 [PMID: 19782158 DOI: 10.1016/j.ijsu.2009.09.006]
- 24 **Springborg LK**, Møller MN. Submandibular gland excision: long-term clinical outcome in 139 patients operated in a single institution. *Eur Arch Otorhinolaryngol* 2013; **270**: 1441-1446 [PMID: 22941392 DOI: 10.1007/s00405-012-2175-4]
- 25 **Schneyer LH**, Levin LK. Rate of secretion by individual salivary gland pairs of man under conditions of reduced exogenous stimulation. *J Appl Physiol* 1955; **7**: 508-512 [PMID: 14367237 DOI: 10.1152/jappl.1955.7.5.508]
- 26 **Corvino A**, Caruso M, Varelli C, Di Gennaro F, Pignata S, Corvino F, Vallone G, Catalano O. Diagnostic imaging of parotid gland oncocytoma: a pictorial review with emphasis on ultrasound assessment. *J Ultrasound* 2021; **24**: 241-247 [PMID: 32710434 DOI: 10.1007/s40477-020-00511-5]
- 27 **Lee TH**, Lin YS, Lee WY, Wu TC, Chang SL. Malignant transformation of a benign oncocytoma of the submandibular gland: a case report. *Kaohsiung J Med Sci* 2010; **26**: 327-332 [PMID: 20538238 DOI: 10.1016/S1607-551X(10)70047-4]
- 28 **Eneroth CM**. Oncocytoma of major salivary glands. *J Laryngol Otol* 1965; **79**: 1064-1072 [PMID: 5858167 DOI: 10.1017/s0022215100064811]
- 29 **Chen B**, Hentzelman JI, Walker RJ, Lai JP. Oncocytoma of the Submandibular Gland: Diagnosis and Treatment Based on Clinicopathology. *Case Rep Otolaryngol* 2016; **2016**: 8719030 [PMID: 27722003 DOI: 10.1155/2016/8719030]
- 30 **Brandwein MS**, Huvos AG. Oncocytic tumors of major salivary glands. A study of 68 cases with follow-up of 44 patients. *Am J Surg Pathol* 1991; **15**: 514-528 [PMID: 2031528 DOI: 10.1097/0000478-199106000-00002]
- 31 **Ben-Shoshan M**, Lacroix Y. A giant sialolith in a 16-year-old boy presenting to the emergency room. *Arch Dis Child* 2014; **99**: 883 [PMID: 24771306 DOI: 10.1136/archdischild-2013-305804]
- 32 **Akinyamoju AO**, Adisa AO. Non-surgical extraction of a massive sialolith in the Wharton's duct of a Nigerian. *Afr J Med Med Sci* 2015; **44**: 177-180 [PMID: 26937532]
- 33 **Rodrigues GHC**, Carvalho VJG, Alves FA, Domaneschi C. Giant submandibular sialolith conservatively treated. *Autops Case Rep* 2017; **7**: 9-11 [PMID: 28536681 DOI: 10.4322/acr.2017.005]
- 34 **Brooks JK**, Macauley MR, Price JB. Concurrent giant sialoliths within the submandibular gland parenchyma and distal segment of Wharton's duct: Novel case report. *Gerodontology* 2021; **38**: 437-440 [PMID: 33586807 DOI: 10.1111/ger.12544]
- 35 **Zakaria MA**. Giant calculi of the submandibular salivary gland. *Br J Oral Surg* 1981; **19**: 230-232 [PMID: 6945127 DOI: 10.1016/0007-117x(81)90009-3]
- 36 **Naraynsingh V**. Giant submandibular gland calculi. *J Oral Maxillofac Surg* 1985; **43**: 384-385 [PMID: 3857304 DOI: 10.1016/0278-2391(85)90263-0]

- 37 **Frame J.** W. and A. J. Smith. Large calculi of the submandibular salivary glands. *International journal of oral and maxillofacial surgery* 15.6 (1986): 769-771
- 38 **Kaltman S,** Eichner M. Giant sialolithiasis appearing as odontogenic infection. *J Am Dent Assoc* 1987; **115**: 425-426 [PMID: 3476663 DOI: 10.14219/jada.archive.1987.0245]
- 39 **Raveenthiran V,** Hayavadana Rao PV. Giant calculus in the submandibular salivary duct: report of the first prepubertal patient. *Pediatr Surg Int* 2004; **20**: 163-164 [PMID: 14760496 DOI: 10.1007/s00383-003-1113-3]
- 40 **Yildirim A.** A case of giant sialolith of the submandibular salivary gland. *Ear Nose Throat J* 2004; **83**: 360-361 [PMID: 15195885]
- 41 **Chan EK,** Patel ND. Giant calculus of the submandibular salivary duct. *Ear Nose Throat J* 2006; **85**: 306, 308 [PMID: 16771020]
- 42 **Graziani F,** Vano M, Cei S, Tartaro G, Mario G. Unusual asymptomatic giant sialolith of the submandibular gland: a clinical report. *J Craniofac Surg* 2006; **17**: 549-552 [PMID: 16770196 DOI: 10.1097/00001665-200605000-00027]
- 43 **Ledesma-Montes C,** Garcés-Ortiz M, Salcido-García JF, Hernández-Flores F, Hernández-Guerrero JC. Giant sialolith: case report and review of the literature. *J Oral Maxillofac Surg* 2007; **65**: 128-130 [PMID: 17174777 DOI: 10.1016/j.joms.2005.10.053]
- 44 **Biddle RJ,** Arora S. Giant Sialolith of the Submandibular Salivary Gland. *Radiol Case Rep* 2008; **3**: 101 [PMID: 27303513 DOI: 10.2484/rcr.2007.v3i2.101]
- 45 **Rai M,** Burman R. Giant submandibular sialolith of remarkable size in the comma area of Wharton's duct: a case report. *J Oral Maxillofac Surg* 2009; **67**: 1329-1332 [PMID: 19446227 DOI: 10.1016/j.joms.2008.11.014]
- 46 **Soares EC,** Costa FW, Pessoa RM, Bezerra TP. Giant salivary calculus of the submandibular gland. *Otolaryngol Head Neck Surg* 2009; **140**: 128-129 [PMID: 19130977 DOI: 10.1016/j.otohns.2008.08.018]
- 47 **Rivera-Serrano CM,** Schaitkin BM. Bilateral giant submandibular sialoliths and the role for salivary endoscopy. *Am J Otolaryngol* 2011; **32**: 85-87 [PMID: 20022665 DOI: 10.1016/j.amjoto.2009.09.011]
- 48 **Krishnan B,** Gehani RE, Shehumi MI. Submandibular giant sialoliths-2 case reports and review of the literature. *Indian J Otolaryngol Head Neck Surg* 2009; **61**: 55-58 [PMID: 23120671 DOI: 10.1007/s12070-009-0019-3]
- 49 **Emir H,** Kaptan ZK, Uzunkulaoglu H, Dogan S. A rare case of asymptomatic bilateral submandibular gland sialolithiasis: a giant, fistulized calculus on the right and multiple calculi on the left. *Ear Nose Throat J* 2010; **89**: 502-504 [PMID: 20981664]
- 50 **Shetty BN,** Sharma P. Unusual case of a projecting intraoral giant sialolith. *Indian J Surg* 2010; **72**: 155-157 [PMID: 23133232 DOI: 10.1007/s12262-010-0028-1]
- 51 **Boffano P,** Gallesio C. Surgical treatment of a giant sialolith of the Wharton duct. *J Craniofac Surg* 2010; **21**: 134-135 [PMID: 20072020 DOI: 10.1097/SCS.0b013e3181c46bc4]
- 52 **Arun Kumar KV,** Garg N, Kumar V. Oversized submandibular gland sialolith: a report of two cases. *J Maxillofac Oral Surg* 2015; **14**: 116-119 [PMID: 25838684 DOI: 10.1007/s12663-012-0361-y]
- 53 **Pandarakalam C,** Goebel WM, Seyer B. Chronic sclerosing sialadenitis or Küttner's tumor associated with a giant sialolith: a case report. *Oral Surg Oral Med Oral Pathol Radiol* 2013; **115**: e38-e40 [PMID: 23312536 DOI: 10.1016/j.oooo.2012.10.011]
- 54 **Iqbal A,** Gupta AK, Natu SS. Unusually large sialolith of Wharton's duct. *Ann Maxillofac Surg* 2012; **2**: 70-73 [PMID: 23483770 DOI: 10.4103/2231-0746.95327]
- 55 **Bhullar RS,** Dhawan A, Bhullar K, Malhotra S. Giant submandibular gland duct sialolith mimicking an impacted canine tooth. *Natl J Maxillofac Surg* 2015; **6**: 89-92 [PMID: 26668461 DOI: 10.4103/0975-5950.168219]
- 56 **Shahoon H,** Farhadi S, Hamed R. Giant sialoliths of Wharton duct: Report of two rare cases and review of literature. *Dent Res J (Isfahan)* 2015; **12**: 494-497 [PMID: 26604966 DOI: 10.4103/1735-3327.166238]
- 57 **Arslan S,** Vuralkan E, Çobanoğlu B, Arslan A, Ural A. Giant sialolith of submandibular gland: report of a case†. *J Surg Case Rep* 2015; **2015** [PMID: 25848088 DOI: 10.1093/jscr/rjv043]
- 58 **Oliveira Tde P,** Oliveira IN, Pinheiro EC, Gomes RC, Mainenti P. Giant sialolith of submandibular gland duct treated by excision and ductal repair: a case report. *Braz J Otorhinolaryngol* 2016; **82**: 112-115 [PMID: 26420563 DOI: 10.1016/j.bjorl.2015.03.013]
- 59 **Weinberg A,** Albers AE. Giant sialolith after submandibulectomy. *Pan Afr Med J* 2016; **25**: 228 [PMID: 28293344 DOI: 10.11604/pamj.2016.25.228.10717]
- 60 **Gadve V,** Mohite A, Bang K, Sheno SR. Unusual giant sialolith of Wharton's duct. *Indian J Dent* 2016; **7**: 162-164 [PMID: 27795655 DOI: 10.4103/0975-962X.174594]
- 61 **Goh LC,** Chitra BK, Shaariyah MM, Ng WS. Transcervical approach to the removal of a deep-seated giant submandibular calculus and the submandibular gland. *BMJ Case Rep* 2016; **2016** [PMID: 27793872 DOI: 10.1136/bcr-2016-217514]
- 62 **Omezli MM,** Ayranci F, Sadik E, Polat ME. Case report of giant sialolith (megalith) of the Wharton's duct. *Niger J Clin Pract* 2016; **19**: 414-417 [PMID: 27022811 DOI: 10.4103/1119-3077.179273]
- 63 **Lim EH,** Nadarajah S, Mohamad I. Giant Submandibular Calculus Eroding Oral Cavity Mucosa. *Oman Med J* 2017; **32**: 432-435 [PMID: 29026477 DOI: 10.5001/omj.2017.81]
- 64 **Iwai T,** Izumi T, Ohya T, Oguri S, Tohna I. Giant Sialolith of the Submandibular Gland. *J Clin Diagn Res* 2017; **11**: ZJ03-ZJ04 [PMID: 28969298 DOI: 10.7860/JCDR/2017/29383.10391]
- 65 **Sakthivel P,** Yogal R, Singh S, Sharma P, Singh CA. Giant Sialolith of Submandibular Duct. *JNMA J Nepal Med Assoc* 2017; **56**: 262-264 [PMID: 28746326]
- 66 **Singh AK,** Yeola M, Raghuvanshi P, Singh N. Giant Wharton's duct sialolithiasis causing sialo-oral fistula: a rare case report with literature review. *J Family Med Prim Care* 2020; **9**: 5793-5795 [PMID: 33532437 DOI: 10.4103/jfmpc.jfmpc_1525_20]
- 67 **Abraham ZS,** Mathias M, Kahinga AA. Unusual giant calculus of the submandibular duct: Case report and literature review. *Int J Surg Case Rep* 2021; **84**: 106139 [PMID: 34216915 DOI: 10.1016/j.ijscr.2021.106139]
- 68 **Thong HK,** Othman IA, Ramli R, Sabir Husin Athar PP. Painless Giant Submandibular Gland Sialolith: A Case Report. *Cureus* 2021; **13**: e19429 [PMID: 34926020 DOI: 10.7759/cureus.19429]
- 69 **Kumar ND,** Sherubin JE, Bagavathy K. Sialolithiasis: An Unusually Large Salivary Stone. *J Maxillofac Oral Surg* 2021; **20**: 227-229 [PMID: 33927489 DOI: 10.1007/s12663-019-01223-y]
- 70 **Ungari C,** Cicconetti A, Cerbelli E, Sulpasso A, Filiaci F. Giant submandibular sialolith: a case report. *Clin Ter* 2022; **173**: 217-221 [PMID: 35612333 DOI: 10.7417/CT.2022.2421]
- 71 **Mohsin SF,** Riyaz MA, Alqazlan AA. Detection and Management of Giant Submandibular Gland Sialolith. *Pak J Med Sci* 2022; **38**: 2030-2033 [PMID: 36246686 DOI: 10.12669/pjms.38.7.5988]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

