

Supplementary Table 1 ICD-10 codes

| | |
|---|------------------|
| Acquired absence of gallbladder | Z90.49 |
| Calculus of gallbladder without cholecystitis without obstruction | K80.20 |
| Simple fatty liver | K76.0 |
| Nonalcoholic steatohepatitis | K75.81 |
| Alcohol abuse | CM_ALCOHOL |
| Diabetes Mellitus | CM_DM CM_DMCX |
| Dyslipidemia | E78.5 |
| Hypertension | CM_HTN_C |
| Metabolic syndrome | E88.81 |
| Nicotine dependence | F17.2 |
| F17.20 Nicotine dependence, unspecified | |
| F17.200 uncomplicated | |
| F17.201 in remission | |
| F17.203 Nicotine dependence unspecified, with withdrawal | |
| F17.208 with other nicotine-induced disorders | |
| F17.209 with unspecified nicotine-induced disorders | |
| F17.21 Nicotine dependence, cigarettes | |
| F17.210 uncomplicated | |

| | |
|--|---------|
| <p>F17.211 in remission</p> <p>F17.213 with withdrawal</p> <p>F17.218 with other nicotine-induced disorders</p> <p>F17.219 with unspecified nicotine-induced disorders</p> <p>F17.22 Nicotine dependence, chewing tobacco</p> <p>F17.220 uncomplicated</p> <p>F17.221 in remission</p> <p>F17.223 with withdrawal</p> <p>F17.228 with other nicotine-induced disorders</p> <p>F17.229 with unspecified nicotine-induced disorders</p> <p>F17.29 Nicotine dependence, other tobacco product</p> <p>F17.290 uncomplicated</p> <p>F17.291 in remission</p> <p>F17.293 with withdrawal</p> <p>F17.298 with other nicotine-induced disorders</p> <p>F17.299 with unspecified nicotine-induced disorders</p> | |
| History of tobacco dependence | Z87.891 |

| | |
|-----------------|----------|
| Tobacco use NOS | Z72.0 |
| Obesity | CM_OBESE |