

World Journal of *Clinical Cases*

World J Clin Cases 2020 December 26; 8(24): 6213-6545



MINIREVIEWS

- 6213 Role of gut microbiome in regulating the effectiveness of metformin in reducing colorectal cancer in type 2 diabetes

Huang QY, Yao F, Zhou CR, Huang XY, Wang Q, Long H, Wu QM

ORIGINAL ARTICLE**Retrospective Cohort Study**

- 6229 Impact factors of lymph node retrieval on survival in locally advanced rectal cancer with neoadjuvant therapy

Mei SW, Liu Z, Wang Z, Pei W, Wei FZ, Chen JN, Wang ZJ, Shen HY, Li J, Zhao FQ, Wang XS, Liu Q

Retrospective Study

- 6243 Three-year follow-up of Coats disease treated with conbercept and 532-nm laser photocoagulation

Jiang L, Qin B, Luo XL, Cao H, Deng TM, Yang MM, Meng T, Yang HQ

- 6252 Virus load and virus shedding of SARS-CoV-2 and their impact on patient outcomes

Chen PF, Yu XX, Liu YP, Ren D, Shen M, Huang BS, Gao JL, Huang ZY, Wu M, Wang WY, Chen L, Shi X, Wang ZQ, Liu YX, Liu L, Liu Y

- 6264 Risk factors for *de novo* hepatitis B during solid cancer treatment

Sugimoto R, Furukawa M, Senju T, Aratake Y, Shimokawa M, Tanaka Y, Inada H, Noguchi T, Lee L, Miki M, Maruyama Y, Hashimoto R, Hisano T

- 6274 Cause analysis and reoperation effect of failure and recurrence after epiblepharon correction in children

Wang Y, Zhang Y, Tian N

Clinical Trials Study

- 6282 Effects of different acupuncture methods combined with routine rehabilitation on gait of stroke patients

Lou YT, Yang JJ, Ma YF, Zhen XC

Observational Study

- 6296 Application of endoscopic submucosal dissection in duodenal space-occupying lesions

Li XY, Ji KY, Qu YH, Zheng JJ, Guo YJ, Zhang CP, Zhang KP

- 6306 Early renal injury indicators can help evaluate renal injury in patients with chronic hepatitis B with long-term nucleos(t)ide therapy

Ji TT, Tan N, Lu HY, Xu XY, Yu YY

Prospective Study

- 6315** Neoadjuvant chemoradiotherapy plus surgery in the treatment of potentially resectable thoracic esophageal squamous cell carcinoma
Yan MH, Hou XB, Cai BN, Qu BL, Dai XK, Liu F

CASE REPORT

- 6322** Uterine rupture in patients with a history of multiple curettages: Two case reports
Deng MF, Zhang XD, Zhang QF, Liu J
- 6330** Pleural effusion and ascites in extrarenal lymphangiectasia caused by post-biopsy hematoma: A case report
Lin QZ, Wang HE, Wei D, Bao YF, Li H, Wang T
- 6337** Eighty-year-old man with rare chronic neutrophilic leukemia caused by CSF3R T618I mutation: A case report and review of literature
Li YP, Chen N, Ye XM, Xia YS
- 6346** Sigmoid colon duplication with ectopic immature renal tissue in an adult: A case report
Namgung H
- 6353** Paraplegia from spinal intramedullary tuberculosis: A case report
Qu LM, Wu D, Guo L, Yu JL
- 6358** Confocal laser endomicroscopy distinguishing benign and malignant gallbladder polyps during choledochoscopic gallbladder-preserving polypectomy: A case report
Tang BF, Dang T, Wang QH, Chang ZH, Han WJ
- 6364** Sclerosing stromal tumor of the ovary with masculinization, Meig's syndrome and CA125 elevation in an adolescent girl: A case report
Chen Q, Chen YH, Tang HY, Shen YM, Tan X
- 6373** Primary pulmonary malignant melanoma diagnosed with percutaneous biopsy tissue: A case report
Xi JM, Wen H, Yan XB, Huang J
- 6380** SRY-negative 45,X/46,XY adult male with complete masculinization and infertility: A case report and review of literature
Wu YH, Sun KN, Bao H, Chen YJ
- 6389** Refractory case of ulcerative colitis with idiopathic thrombocytopenic purpura successfully treated by Janus kinase inhibitor tofacitinib: A case report
Komeda Y, Sakurai T, Sakai K, Morita Y, Hashimoto A, Nagai T, Hagiwara S, Matsumura I, Nishio K, Kudo M
- 6396** Immunotherapies application in active stage of systemic lupus erythematosus in pregnancy: A case report and review of literature
Xiong ZH, Cao XS, Guan HL, Zheng HL

- 6408** Minimally invasive maxillary sinus augmentation with simultaneous implantation on an elderly patient: A case report
Yang S, Yu W, Zhang J, Zhou Z, Meng F, Wang J, Shi R, Zhou YM, Zhao J
- 6418** Congenital nephrogenic diabetes insipidus due to the mutation in *AVPR2* (c.541C>T) in a neonate: A case report
Lin FT, Li J, Xu BL, Yang XX, Wang F
- 6425** Primary gastric melanoma in a young woman: A case report
Long GJ, Ou WT, Lin L, Zhou CJ
- 6432** Extreme venous letting and cupping resulting in life-threatening anemia and acute myocardial infarction: A case report
Jang AY, Suh SY
- 6437** Novel conservative treatment for peritoneal dialysis-related hydrothorax: Two case reports
Dai BB, Lin BD, Yang LY, Wan JX, Pan YB
- 6444** Clinical characteristics of pulmonary cryptococcosis coexisting with lung adenocarcinoma: Three case reports
Zheng GX, Tang HJ, Huang ZP, Pan HL, Wei HY, Bai J
- 6450** Fracture of the scapular neck combined with rotator cuff tear: A case report
Chen L, Liu CL, Wu P
- 6456** Synchronous colonic mucosa-associated lymphoid tissue lymphoma found after surgery for adenocarcinoma: A case report and review of literature
Li JJ, Chen BC, Dong J, Chen Y, Chen YW
- 6465** Novel mutation in the *ASXL3* gene in a Chinese boy with microcephaly and speech impairment: A case report
Li JR, Huang Z, Lu Y, Ji QY, Jiang MY, Yang F
- 6473** Recurrent thrombosis in the lower extremities after thrombectomy in a patient with polycythemia vera: A case report
Jiang BP, Cheng GB, Hu Q, Wu JW, Li XY, Liao S, Wu SY, Lu W
- 6480** Status epilepticus as an initial manifestation of hepatic encephalopathy: A case report
Cui B, Wei L, Sun LY, Qu W, Zeng ZG, Liu Y, Zhu ZJ
- 6487** Delayed diagnosis of prosopagnosia following a hemorrhagic stroke in an elderly man: A case report
Yuan Y, Huang F, Gao ZH, Cai WC, Xiao JX, Yang YE, Zhu PL
- 6499** Oral myiasis after cerebral infarction in an elderly male patient from southern China: A case report
Zhang TZ, Jiang Y, Luo XT, Ling R, Wang JW
- 6504** Rare case of drain-site hernia after laparoscopic surgery and a novel strategy of prevention: A case report
Gao X, Chen Q, Wang C, Yu YY, Yang L, Zhou ZG

- 6511** Extracorporeal shock wave therapy treatment of painful hematoma in the calf: A case report
Jung JW, Kim HS, Yang JH, Lee KH, Park SB
- 6517** Takotsubo cardiomyopathy associated with bronchoscopic operation: A case report
Wu BF, Shi JR, Zheng LR
- 6524** Idiopathic adulthood ductopenia with elevated transaminase only: A case report
Zhang XC, Wang D, Li X, Hu YL, Wang C
- 6529** Successful endovascular treatment with long-term antibiotic therapy for infectious pseudoaneurysm due to *Klebsiella pneumoniae*: A case report
Wang TH, Zhao JC, Huang B, Wang JR, Yuan D
- 6537** Primary duodenal tuberculosis misdiagnosed as tumor by imaging examination: A case report
Zhang Y, Shi XJ, Zhang XC, Zhao XJ, Li JX, Wang LH, Xie CE, Liu YY, Wang YL

ABOUT COVER

Peer-Reviewer of *World Journal of Clinical Cases*, Dr. Adonis Protopapas is a gastroenterology Resident at the first Propaedeutic Department of Internal Medicine of the Aristotle University of Thessaloniki (Greece), located at the A.H.E.P.A Hospital. He earned his Bachelor's degree in 2015 from the Democritus University of Thrace, followed by three Master's of Science degrees, with specializations in clinic pharmacology, medical research methodology, and healthcare management. His research interests are mainly focused on the area of hepatology, although he also participates in various projects related to endoscopy and inflammatory bowel disease. He is particularly fascinated by research on cirrhosis and its complications. (L-Editor: Filipodia)

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

WJCC mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

INDEXING/ABSTRACTING

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, PubMed, and PubMed Central. The 2020 Edition of Journal Citation Reports® cites the 2019 impact factor (IF) for *WJCC* as 1.013; IF without journal self cites: 0.991; Ranking: 120 among 165 journals in medicine, general and internal; and Quartile category: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ji-Hong Liu*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jin-Lai Wang*.

NAME OF JOURNAL

World Journal of Clinical Cases

ISSN

ISSN 2307-8960 (online)

LAUNCH DATE

April 16, 2013

FREQUENCY

Semimonthly

EDITORS-IN-CHIEF

Dennis A Bloomfield, Sandro Vento, Bao-gan Peng

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/2307-8960/editorialboard.htm>

PUBLICATION DATE

December 26, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Observational Study

Early renal injury indicators can help evaluate renal injury in patients with chronic hepatitis B with long-term nucleos(t)ide therapy

Tong-Tong Ji, Ning Tan, Hai-Ying Lu, Xiao-Yuan Xu, Yan-Yan Yu

ORCID number: Tong-Tong Ji 0000-0001-9384-4285; Ning Tan 0000-0001-9917-2192; Hai-Ying Lu 0000-0003-3568-3502; Xiao-Yuan Xu 0000-0002-1759-4330; Yan-Yan Yu 0000-0002-7557-1305.

Author contributions: Ji TT

acquired the data and wrote the manuscript; Tan N analyzed the data and wrote the manuscript; Lu HY designed the research and reviewed and approved the final manuscript; Xu XY and Yu YY instructed the research.

Supported by the National 12th Five-Year Plan for Science and Technology, No. 2018ZX10725-506.

Institutional review board

statement: The study was reviewed and approved by the Ethics Committee of Peking University First Hospital in 2018 (Research No. 264).

Informed consent statement:

Patients who were enrolled in the work provided written informed consent.

Conflict-of-interest statement:

There are no conflicts of interest to report.

Data sharing statement: No

additional data are available.

STROBE statement: The authors

Tong-Tong Ji, Ning Tan, Hai-Ying Lu, Xiao-Yuan Xu, Yan-Yan Yu, Department of Infectious Diseases, Peking University First Hospital, Beijing 100034, China

Corresponding author: Hai-Ying Lu, PhD, Associate Professor, Department of Infectious Diseases, Peking University First Hospital, No. 8 Xishiku Road, Xicheng District, Beijing 100034, China. luhaiying00@126.com

Abstract**BACKGROUND**

Patients with chronic hepatitis B (CHB) with long-term nucleos(t)ide therapy may experience renal insufficiency. Traditional renal function indicators, such as urine protein, serum urea nitrogen (BUN), and serum creatinine, are normal when early mild lesions occur. Therefore, more sensitive renal function indicators are needed.

AIM

To investigate the significance of early renal injury indicators in evaluating renal injury in patients with CHB with long-term nucleos(t)ide therapy.

METHODS

We collected the clinical data of 69 outpatients with CHB at Peking University First Hospital from March 2018 to January 2020 who had been treated with long-term nucleos(t)ide therapy and analyzed the results of early renal injury indicators. Continuous normal distribution data were analyzed by the *t*-test to determine the difference between two groups. Continuous non-normally distributed data were analyzed by the Mann-Whitney U-test between two groups. The Kruskal-Wallis H test was used to determine the differences among multiple groups. Enumeration data were analyzed by the chi-square test. The related factors of early renal injury indicators were analyzed by logistic regression analysis.

RESULTS

The average treatment duration with nucleos(t)ide analogs of the 69 patients with CHB was 99.7 ± 28.7 mo. The cases of patients with elevated BUN and hypophosphatemia were 6 (8.7%) and 13 (18.8%), respectively; 31 (44.9%) patients had abnormal early renal injury indicators, including 9 patients with abnormal urine microalbumin, 7 patients with abnormal urine immunoglobulin, 6 patients with abnormal urine transferrin, and 19 patients with abnormal $\alpha 1$ microglobulin. There were no significant differences in the mean values of age, sex, BUN,

have read the STROBE Statement—checklist of items, and the manuscript was prepared and revised according to the STROBE Statement—checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Specialty type: Medicine, research and experimental

Country/Territory of origin: China

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

Received: September 23, 2020

Peer-review started: September 23, 2020

First decision: September 29, 2020

Revised: October 6, 2020

Accepted: November 2, 2020

Article in press: November 2, 2020

Published online: December 26, 2020

P-Reviewer: Kawaguchi T

S-Editor: Chen XF

L-Editor: Filipodia

P-Editor: Li JH



estimated glomerular filtration rate (eGFR), serum uric acid, serum calcium, or serum phosphorus between the two groups of patients with and without early renal injury indicators. However, the mean levels of serum creatinine and urine creatinine, N-acetyl- β -D-glucosidase enzyme, α 1 microglobulin, and urine immunoglobulin in the former group of patients were significantly higher than those in the latter group of patients ($P < 0.05$). The incidence of early renal injury in patients with eGFR ≥ 90 , 60-89, and 30-59 mL/(min \cdot 1.73 m²) was 36.4% (8/22), 47.6% (20/42), and 60% (3/5), respectively. Logistic regression analysis results showed that gamma-glutamyl transpeptidase [odds ratio (OR) = 1.05 (1.008-1.093), $P = 0.020$], direct bilirubin [OR = 1.548 (1.111-2.159), $P = 0.010$], serum creatinine [OR = 1.079 (1.022-1.139), $P = 0.006$], and age [OR = 0.981 (0.942-1.022), $P = 0.357$] were independent predictors of early renal injury.

CONCLUSION

Patients with CHB treated with long-term nucleos(t)ide analog therapy had a high probability of early renal injury, and early renal injury indicators were highly sensitive and could be used to monitor early renal impairment.

Key Words: Early renal injury; Chronic hepatitis B; Nucleos(t)ide analog; N-acetyl- β -D-glucosidase enzyme; α 1 microglobulin; Urine immunoglobulin

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Early renal injury indicators had higher sensitivity and could be used to screen early renal injury in patients with chronic hepatitis B with long-term nucleos(t)ide analog therapy.

Citation: Ji TT, Tan N, Lu HY, Xu XY, Yu YY. Early renal injury indicators can help evaluate renal injury in patients with chronic hepatitis B with long-term nucleos(t)ide therapy. *World J Clin Cases* 2020; 8(24): 6306-6314

URL: <https://www.wjgnet.com/2307-8960/full/v8/i24/6306.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v8.i24.6306>

INTRODUCTION

Nucleos(t)ide analogs (NAs), including lamivudine (LAM), adefovir dipivoxil (ADV), telbivudine (LdT), entecavir (ETV)^[1,2], tenofovir disoproxil fumarate (TDF)^[3,4], and tenofovir alafenamide fumarate (TAF)^[5-7], are important antiviral drugs for patients with chronic hepatitis B (CHB). Patients with CHB need to have their renal function monitored regularly during treatment because long-term antiviral therapy can cause renal injury. Early renal injury indicators can detect early mild renal lesions, which is helpful in the early detection, early diagnosis, and early treatment of renal dysfunction. In this article, we retrospectively analyzed the clinical data of patients with CHB who received long-term NA therapy in the outpatient Department of Liver Disease of Peking University First Hospital. We mainly observed and discussed the clinical value and significance of early renal injury indicators in the evaluation of renal injury in patients with CHB with long-term NA therapy.

MATERIALS AND METHODS

Subjects

In this article, we retrospectively analyzed the clinical data of 69 patients with CHB who had received long-term NAs in the outpatient Department of Liver Disease of Peking University First Hospital from March 2018 to January 2020, including 48 males (69.6%) with an average age of 54.1 \pm 12.0-years-old and 21 females (30.4%) with an average age of 57.6 \pm 9.2-years-old. The diagnostic criteria refer to the "Guidelines for the Prevention and Treatment of Chronic Hepatitis B"^[8] of China. The exclusion

criteria included antiviral treatment course < 2 years; patients with chronic liver disease caused by non-hepatitis B virus (HBV) or chronic HBV infection with other liver diseases; patients with other chronic kidney diseases; and patients with diabetes mellitus. The study protocol was approved by the Ethics Committee of Peking University First Hospital in 2018 (Research No. 264), and all patients signed informed consent forms.

Research methods

The clinical data of patients were collected, including age, sex, type of antiviral drugs, treatment time, liver function, HBV deoxyribonucleic acid (DNA), serum urea nitrogen (BUN, normal value: 1.8-7.1 $\mu\text{mol/L}$), creatinine (normal value: 44-133 $\mu\text{mol/L}$), uric acid (normal value: 150-420 $\mu\text{mol/L}$), serum calcium (normal value: 2.12-2.75 mmol/L), serum phosphorus (normal value: 0.96-1.62 mmol/L), serum magnesium (normal values: 0.8-1.2 mmol/L), estimated glomerular filtration rate [eGFR, mL/(min \cdot 1.73 m^2)], and early renal injury indicators such as urine creatinine (mmol/L), urine microalbumin (normal value: 0-19 mg/L), urine transferrin (normal value: 0-2 mg/L), urine α 1-microglobulin (normal value: 0-12 mg/L), urine immunoglobulin (normal value: < 8 mg/L), and N-acetyl- β -D-glucosidase (NAG) enzyme (normal value: 0.3-12 U/L). As long as one of the early renal injury indicators was abnormal, it was defined as early renal damage. Serum creatinine was measured by the picric acid rate method (Zhongsheng Beikong Company's kit; Hitachi 7600 automatic biochemical analyzer). The detection methods of early renal function indicators were as follows: Urinary creatinine was measured by the picric acid method (Zhongsheng Beikong Biotechnology, Beijing, China); urine microalbumin, urine transferrin, urine α 1-microglobulin, and urine immunoglobulin were determined by immune scatter turbidity (Beckman Coulter, Brea, CA, United States); and NAG enzyme was determined by the MNP-G1CNAc substrate method (Gcell Kit; Beijing Jiuqiang Biological Company, Beijing, China). HBV DNA was detected by the Da'an Real-Time Fluorescence polymerase chain reaction HBV DNA Quantitative Detection Kit of Sun Yat-sen University, and the lower limit of detection value was 100 IU/mL.

Statistical analysis

Statistical analyses were performed using SPSS 25.0 software (Armonk, NY, United States). The measurement data were expressed as (mean \pm standard deviation) or the median, and continuous normally distributed data were analyzed by *t*-test to determine the difference between two groups. Continuous non-normally distributed data were analyzed by the Mann-Whitney *U*-test between two groups. The Kruskal-Wallis H test was used to determine the differences among multiple groups. Enumeration data were analyzed by the chi-square test. The related factors of early renal injury indicators were analyzed by logistic regression analysis, and *P* < 0.05 was considered statistically significant.

RESULTS

Antiviral therapy efficacy

A total of 69 patients with CHB were enrolled in this study, including 48 males (69.6%) and 21 females (30.4%). Among them, one patient received LAM, 7 patients received ADV, 41 patients received ETV, 14 patients received ADV + ETV, one patient received LdT, 4 patients received TDF, and one patient received TAF. The average antiviral treatment duration was 99.7 \pm 28.7 mo. All patients with CHB had stable situations, and most of them had normal liver function with an undetectable HBV DNA level (specific values were omitted).

Influence of different antiviral drugs on renal function

The serum creatinine level of 69 patients was in the normal range. Six patients (8.7%) had an increase in BUN levels, and 13 patients had hypophosphatemia (18.8%). There was no significant abnormality in the routine urine test of all patients. Patients in the ADV/TDF treatment group had a higher percentage of increased BUN, early renal injury, and hypophosphatemia than the patients in the non-ADV/TDF treatment group, but the differences between the two groups were not statistically significant (*P* > 0.05). Specific data are shown in Table 1. Hypermagnesemia occurred in 4 patients and 6 patients in the ADV/TDF and non-ADV/TDF treatment groups, respectively, and hypercalcemia occurred in 1 patient and 2 patients, respectively. Elevated serum

Table 1 Renal function test results of patients with and without adefovir dipivoxil / tenofovir disoproxil fumarate treatment

Treatment group	Increased BUN	Early renal damage	eGFR \geq 90	eGFR 89-60	eGFR 59-30	Hypophosphatemia
ADV/TDF group, <i>n</i> = 25	3 (12%)	15 ¹ (60%)	6 (24%)	16 (64%)	3 (12%)	6 (24%)
Non-ADV/TDF group, <i>n</i> = 44	3 (6.8%)	16 (36.4%)	16 (36.4%)	26 (59.1%)	2 (4.5%)	7 (15.9%)

¹Compared with the non-ADV/TDF treatment group, $\chi^2 = 3.600$, $P = 0.057$. ADV: Adefovir dipivoxil; BUN: Serum urea nitrogen; eGFR: Estimated glomerular filtration rate; TDF: Tenofovir disoproxil fumarate.

creatinine levels and hypocalcemia were not found in all patients.

Impact of antiviral therapy on the eGFR

According to the eGFR value, 69 patients with CHB were divided into the following three groups: (1) EGFR \geq 90 mL/(min \cdot 1.73 m²) group; (2) EGFR = 60-89 mL/(min \cdot 1.73 m²) group; and (3) EGFR = 30-59 mL/(min \cdot 1.73 m²) group. The number of patients in the three groups was 22 (31.9%), 42 (60.9%), and 5 (7.2%), respectively. There was no significant difference in sex, the mean value of liver function, serum uric acid, serum calcium, or serum magnesium among the three groups ($P > 0.05$, specific data omitted). Differences in age, serum creatinine, and the duration of antiviral treatment among the three groups were statistically significant ($P < 0.05$). The serum levels of urea nitrogen and phosphorus in the third group of patients were significantly different from those in the first group and the second group of patients ($P < 0.05$). With the decrease in eGFR value, the percentage of patients with hypophosphatemia or early renal injury increased, but the difference among the three groups was not statistically significant ($P > 0.05$). The specific figures are shown in [Table 2](#).

Impact of antiviral therapy on early renal injury indicators

A total of 31 (44.9%) of 69 patients had abnormal early renal injury indicators, which were considered early renal damage, including 9 patients with abnormal urine microalbumin, 6 patients with abnormal urine transferrin, 14 patients with abnormal NAG enzyme, 19 patients with abnormal α 1 microglobulin, and 7 patients with abnormal urinary immunoglobulin. There were no significant differences in the mean values of age, sex, alanine aminotransferase, aspartate aminotransferase, microalbumin, alkaline phosphatase, total bilirubin, BUN, eGFR, serum uric acid, serum calcium, serum phosphorus, and serum magnesium in patients with or without early renal injury ($P > 0.05$). However, the differences in gamma-glutamyl transpeptidase (GGT), direct bilirubin (DBIL), and serum creatinine levels, as well as urine creatinine, NAG enzyme, α 1 microglobulin, and urine immunoglobulin levels were statistically significant between the two groups of patients ($P < 0.05$). Patients with early renal injury had a greater percentage of eGFR $<$ 70 mL/(min \cdot 1.73 m²) than patients without early renal injury, but the difference was not statistically significant ($P > 0.05$). The specific figures are shown in [Table 3](#).

Results of logistic regression analysis

The results of logistic regression analysis showed that GGT [odds ratio (OR) = 1.05 (1.008-1.093), $P = 0.020$], DBIL [OR = 1.548 (1.111-2.159), $P = 0.010$], blood creatinine [OR = 1.079 (1.022-1.139), $P = 0.006$], and age [OR = 0.981 (0.942-1.022), $P = 0.357$] were independent predictors of early renal injury for patients with CHB with NA treatment.

DISCUSSION

For most patients with CHB, NA therapy is safe and well tolerated, but some patients may experience adverse reactions, such as renal insufficiency, myositis, rhabdomyolysis, and lactic acidosis^[9,10]. Among the NA drugs, nephrotoxic side effects with ADV are common and mainly manifest as an increase in serum creatinine and a decrease in serum phosphorus, which can cause Fanconi syndrome, renal failure, osteomalacia, and fractures^[11,12]. The main reason for its nephrotoxicity is that the organic anion transporter 1 of the proximal tubule has a strong affinity for NAs and can promote the absorption of ADV, resulting in a higher concentration in the

Table 2 Examination results of patients in different estimated glomerular filtration rate value groups

eGFR value [mL/(min·1.73 m ²)]	≥ 90, n = 22	60-89, n = 42	30-59, n = 5
Age in yr	46.8 ± 11.4	58.3 ± 11.9	69.6 ± 16.10 ¹
Sex as M/F	18/4	27/15	3/2
Antiviral treatment time in mo	84.0 ± 30.9	111.1 ± 39.2	96.8 ± 38.1 ²
ALT in IU/L	33.36 ± 16.83	22.31 ± 10.35	21.80 ± 10.32
Creatinine in μmol/L	74.71 ± 9.98	86.20 ± 12.83	107.4 ± 12.54 ³
BUN in mmol/L	4.56 ± 1.17	5.38 ± 1.40	6.32 ± 1.49 ⁴
Cases of increased BUN	0	3 (7.1%)	3 (60%) ⁵
Blood phosphorus in mmol/L	0.94 ± 0.16	1.01 ± 0.15	0.86 ± 0.20 ⁶
Cases of decreased blood phosphorus	4 (18.2%)	7 (16.7%)	2 (40%)
Cases of early renal injury	8 (36.4%)	20 (47.6%)	3 (60%)
Cases of abnormal urine microalbumin	3 (13.6%)	4 (9.5%)	1 (20%)
Cases of abnormal urine transferrin	1 (4.5%)	5 (11.9%)	0
Cases of abnormal NAG enzyme	5 (22.7%)	8 (19.0%)	1 (20%)
Cases of abnormal α1 microglobulin	5 (22.7%)	11 (26.2%)	3 (60%)
Cases of abnormal urine immunoglobulin	1 (4.5%)	5 (11.9%)	1 (20%)

Comparison among the three groups.

¹Indicates K = 14.979, P = 0.001.

²Indicates K = 7.623, P = 0.022.

³Indicates K = 19.034, P = 0.000.

⁴Indicates comparison between the third group and the first group, t = 2.575, P = 0.012.

⁵Indicates Fisher's exact test without chi-square value, P = 0.011.

⁶Indicates comparison between the third group and the second group, t = 2.045, P = 0.047. ALT: Alanine aminotransferase; BUN: Serum urea nitrogen; eGFR: Estimated glomerular filtration rate; F: Female; M: Male; NAG: N-acetyl-β-D-glucosidase.

proximal tubule. This kind of kidney damage can be reversed after withdrawal of antiviral drugs, but a few patients still show persistent renal damage.

Gara *et al*^[12] carried out an average of 7.4 years of follow-up observations on 51 patients with CHB who received ADV or TDF antiviral therapy and found that 7 patients had proximal renal tubular damage. After the antiviral treatment was changed from ADV to ETV, kidney damage achieved improvement in 6 patients. Chinese scholars also reported that ADV could reduce the eGFR value, while LAM and ETV had no significant impact^[9]. Amarapurkar *et al*^[13] analyzed renal function in 292 patients with CHB with a 64-mo treatment of ADV combined with LAM and found that 9.6% of the patients had renal damage, and 27.1% of patients had hypophosphatemia, of which 14 developed Fanconi syndrome. In previous clinical observations, we also found that the renal side effect of ADV was the largest, ETV was the second largest, and LAM was the lowest^[14]. In this study, we retrospectively analyzed the clinical data of 69 patients with CHB with an average of 99.7 mo of antiviral treatment with NAs. The results showed that 6 patients (8.7%) had elevated levels of BUN, and 13 patients had hypophosphatemia (18.8%). With the decrease in the eGFR value, the mean values of BUN and creatinine increased significantly, the mean values of serum phosphorus decreased significantly (P < 0.05), and the percentage of patients with hypophosphatemia or early renal injury increased. Moreover, patients with ADV/TDF treatment had a greater percentage of high levels of BUN, early renal injury (60% vs 36.4%), hypophosphatemia, and eGFR value less than 70 mL/(min·1.73 m²) compared to patients without ADV/TDF treatment. This further illustrated that long-term treatment with NAs had a certain degree of impact on renal function, especially the potential renal toxicity of ADV, which was significant. Therefore, ADV should not be used in clinical antiviral therapy for patients with CHB.

Early renal mild lesions are reversible, but clinical symptoms are not obvious. Traditional renal function indicators such as urine protein, BUN, and serum creatinine are normal when early mild lesions occur. When the above renal function indicators present abnormalities, approximately 50% of the nephrons have been irreversibly

Table 3 Examination results of patients with or without early renal injury

Early renal injury groups	Normal group (n = 38)	Abnormal group (n = 31)
Age in yr	55.8 ± 15.0	55.0 ± 11.9
Sex as M/F	25/13	23/8
Antiviral treatment time in mo	99.3 ± 37.7	104.1 ± 39.4
GGT as IU/L	21.7 ± 11.3	37.2 ± 16.2 ¹
DBIL as μmol/L	2.5 ± 1.5	3.5 ± 1.4 ²
Creatinine as μmol/L	81.0 ± 14.0	87.9 ± 14.5 ³
BUN as mmol/L	5.0 ± 1.2	5.4 ± 1.6
Cases of increased serum BUN	2 (5.3%)	3 (9.1%)
eGFR as mL/(min 1.73 m ²)	86.5 ± 15.3	80.7 ± 15.5
eGFR ≥ 90, n = 22	14 (36.8%)	8 (25.8%)
eGFR: 70-89, n = 42	18 (47.4%)	13 (41.8%)
eGFR: 60-69, n = 42	4 (10.5%)	7 (22.6%)
eGFR: 30-59, n = 5	2 (5.3%)	3 (9.7%)
UA as μmol/L	314.2 ± 76.2	338.8 ± 88.4
Serum calcium as mmol/L	2.36 ± 0.10	2.35 ± 0.12
Serum phosphorus as mmol/L	0.99 ± 0.16	0.97 ± 0.17
Cases of decreased serum phosphorus	6 (15.8%)	7 (22.6%)
Magnesium as mmol/L	0.95 ± 0.07	0.97 ± 0.07
Urinary creatinine as mmol/L	6.9 (2.5-20.3)	12.4 (5.8-29.5) ⁴
Urine microalbumin as mg/L	4.0 (2-12.8)	11.2 (2-123) ⁵
NAG enzyme as U/L	4.1 (1-12)	10.4 (2-59) ⁶
α1 microglobulin as mg/L	4.9 (4-17.4)	15.5 (4.68-158) ⁷
Urinary immunoglobulin as mg/L	3 (3-4.1)	6.7 (3.14-29.3) ⁸

Comparison between the two groups.

¹Indicates $t = -2.508$, $P = 0.015$.

²Indicates $t = -2.208$, $P = 0.031$.

³Indicates $t = -2.023$, $P = 0.047$.

⁴Indicates $U = 306.5$, $P = 0.001$.

⁵Indicates $U = 184.5$, $P = 0.000$.

⁶Indicates $U = 202.0$, $P = 0.000$.

⁷Indicates $U = 91.0$, $P = 0.000$.

⁸Indicates $U = 42.5$, $P = 0.000$. BUN: Serum urea nitrogen; DBIL: Direct bilirubin; eGFR: Estimated glomerular filtration rate; F: Female; GGT: Gamma-glutamyl transpeptidase; M: Male; NAG: N-acetyl-β-D-glucosidase; UA: Uric acid.

damaged, and the optimal period of treatment has been missed. Therefore, early renal injury indicators have great significance for the early detection, early diagnosis, and early treatment of renal dysfunction. Early renal injury indicators are generally divided into two categories: (1) Glomerular markers, which are mainly used to detect damage to glomerular function, including urinary microalbumin, urinary immunoglobulin G, transferrin, *etc.*; and (2) Renal tubule markers, which reflect the status of tubular damage, including urine medium- and low-molecular-weight proteins (such as urine α1 microglobulin, urine β2 microglobulin, urine retinol-binding protein, *etc.*) and urine enzymes (such as NAG enzyme, lysozyme, *etc.*). In addition, some nonspecific indicators, such as alkaline phosphatase and GGT, can be used as supplementary markers for detecting proximal tubule injury.

Previous clinical research has mainly focused on reporting renal tubule damage caused by NAs. Of the 69 patients with CHB in our study, a total of 31 (44.9%) patients had abnormal early renal injury indicators, indicating the development of early renal damage, including 9 cases of abnormal urine microalbumin, 7 cases of abnormal urine

globulin, 6 cases of abnormal urinary transferrin, 14 cases of abnormal NAG enzyme, and 19 cases of abnormal α 1 microglobulin. This result suggested that long-term treatment with NAs had potential toxicity to tubule function as well as glomerulus function, but the probability of renal tubular injury was relatively high. There were no significant differences in the mean values of age, sex, BUN, eGFR, serum uric acid, serum calcium, serum phosphorus, and serum magnesium in patients with or without early renal damage; however, the levels of urinary creatinine, NAG enzyme, α 1 microglobulin, and urinary immunoglobulin of patients with early renal damage were significantly higher than those of patients without early renal damage ($P < 0.05$). While the serum creatinine values and routine urine test results were normal in all patients, early renal injury could be seen in the patients with eGFR ≥ 90 , 60-89, and 30-59 mL/(min 1.73 m^2), and the incidence of early renal damage was 36.4%, 47.6%, and 60%, respectively. This proved that, compared with traditional renal function indicators, early renal injury indicators could be more sensitive and used earlier to screen for mild renal injury in patients with CHB with NA treatment. Logistic regression analysis results showed that GGT, DBIL, serum creatinine, and age were independent predictors of early renal injury.

CONCLUSION

In summary, the results of this study showed that long-term treatment with NAs had potentially toxic effects on the glomerulus and tubules of the kidney, but renal tubule injury was more common. The incidence of early renal injury reached 44.9%, which should arouse sufficient attention by clinical doctors. Compared with traditional renal function indicators, early renal injury indicators had higher sensitivity and could be used to screen early renal injury in patients with CHB with long-term NA therapy.

ARTICLE HIGHLIGHTS

Research background

Patients with chronic hepatitis B (CHB) with long-term nucleos(t)ide (NA) therapy may experience renal insufficiency. Traditional renal function indicators, such as urine protein, serum urea nitrogen (BUN), and serum creatinine, are normal when early mild lesions occur. Therefore, more sensitive renal function indicators are needed.

Research motivation

To identify more sensitive renal function indicators.

Research objectives

To investigate the significance of early renal injury indicators in evaluating renal injury in patients with CHB with long-term NAs.

Research methods

We collected the clinical data of 69 outpatients with CHB at Peking University First Hospital from March 2018 to January 2020 who had been treated with long-term NA therapy and analyzed the results of early renal injury indicators. Continuous normal distribution data were analyzed by the *t*-test to determine the difference between two groups. Continuous nonnormally distributed data were analyzed by the Mann-Whitney *U*-test between two groups. The Kruskal-Wallis *H* test was used to determine the differences among multiple groups. Enumeration data were analyzed by the chi-square test. The related factors of early renal injury indicators were analyzed by logistic regression analysis.

Research results

The average treatment duration with NAs of the 69 patients with CHB was 99.7 ± 28.7 mo. The cases of patients with elevated BUN and hypophosphatemia were 6 (8.7%) and 13 (18.8%), respectively; 31 (44.9%) patients had abnormal early renal injury indicators, including 9 patients with abnormal urine microalbumin, 7 patients with abnormal urine immunoglobulin, 6 patients with abnormal urine transferrin, and 19 patients with abnormal α 1 microglobulin. There were no significant differences in the mean values of age, sex, BUN, estimated glomerular filtration rate (eGFR), serum uric

acid, serum calcium, or serum phosphorus between the two groups of patients with and without early renal injury indicators. However, the mean levels of serum creatinine and urine creatinine, NAG enzyme, α 1 microglobulin, and urine immunoglobulin in the former group of patients were significantly higher than those in the latter group of patients ($P < 0.05$). The incidence of early renal injury in patients with eGFR ≥ 90 , 60-89 and 30-59 mL/(min \cdot 1.73 m²) was 36.4% (8/22), 47.6% (20/42), and 60% (3/5), respectively. Logistic regression analysis results showed that gamma-glutamyl transpeptidase [odds ratio (OR) = 1.05 (1.008-1.093), $P = 0.020$], direct bilirubin [OR = 1.548 (1.111-2.159), $P = 0.010$], serum creatinine [OR = 1.079 (1.022-1.139), $P = 0.006$], and age [OR = 0.981 (0.942-1.022), $P = 0.357$] were independent predictors of early renal injury.

Research conclusions

Patients with CHB treated with long-term NA therapy had a high probability of early renal injury, and early renal injury indicators were highly sensitive and could be used to monitor early renal impairment.

Research perspectives

Retrospective analysis of CHB patients with deteriorating renal function undergoing NAs therapy can be performed to compare the predictive value of different early renal injury indicators for drug toxicity in renal.

ACKNOWLEDGEMENTS

The authors would like to sincerely thank Professor Lu HY (Peking University First Hospital) for his guidance with the study.

REFERENCES

- 1 **Chang TT**, Gish RG, de Man R, Gadano A, Sollano J, Chao YC, Lok AS, Han KH, Goodman Z, Zhu J, Cross A, DeHertogh D, Wilber R, Colonna R, Apelian D; BEHoLD A1463022 Study Group. A comparison of entecavir and lamivudine for HBeAg-positive chronic hepatitis B. *N Engl J Med* 2006; **354**: 1001-1010 [PMID: [16525137](#) DOI: [10.1056/NEJMoa051285](#)]
- 2 **Chang TT**, Lai CL, Kew Yoon S, Lee SS, Coelho HS, Carrilho FJ, Poordad F, Halota W, Horsmans Y, Tsai N, Zhang H, Tenney DJ, Tamez R, Iloeje U. Entecavir treatment for up to 5 years in patients with hepatitis B e antigen-positive chronic hepatitis B. *Hepatology* 2010; **51**: 422-430 [PMID: [20049753](#) DOI: [10.1002/hep.23327](#)]
- 3 **Hou JL**, Gao ZL, Xie Q, Zhang JM, Sheng JF, Cheng J, Chen CW, Mao Q, Zhao W, Ren H, Tan DM, Niu JQ, Chen SJ, Pan C, Tang H, Wang H, Mao YM, Jia JD, Ning Q, Xu M, Wu SM, Li J, Zhang XX, Ji Y, Dong J, Li J. Tenofovir disoproxil fumarate vs adefovir dipivoxil in Chinese patients with chronic hepatitis B after 48 weeks: a randomized controlled trial. *J Viral Hepat* 2015; **22**: 85-93 [PMID: [25243325](#) DOI: [10.1111/jvh.12313](#)]
- 4 **Liang X**, Gao Z, Xie Q, Zhang J, Sheng J, Cheng J, Chen C, Mao Q, Zhao W, Ren H, Tan D, Niu J, Chen S, Pan C, Tang H, Wang H, Mao Y, Jia J, Ning Q, Xu M, Wu S, Li J, Zhang X, Zhang W, Xiong C, Hou J. Long-term efficacy and safety of tenofovir disoproxil fumarate in Chinese patients with chronic hepatitis B: 5-year results. *Hepatol Int* 2019; **13**: 260-269 [PMID: [30977033](#) DOI: [10.1007/s12072-019-09943-6](#)]
- 5 **Chan HL**, Fung S, Seto WK, Chuang WL, Chen CY, Kim HJ, Hui AJ, Janssen HL, Chowdhury A, Tsang TY, Mehta R, Gane E, Flaherty JF, Massetto B, Gaggar A, Kitrinis KM, Lin L, Subramanian GM, McHutchison JG, Lim YS, Acharya SK, Agarwal K; GS-US-320-0110 Investigators. Tenofovir alafenamide versus tenofovir disoproxil fumarate for the treatment of HBeAg-positive chronic hepatitis B virus infection: a randomised, double-blind, phase 3, non-inferiority trial. *Lancet Gastroenterol Hepatol* 2016; **1**: 185-195 [PMID: [28404091](#) DOI: [10.1016/S2468-1253\(16\)30024-3](#)]
- 6 **Buti M**, Gane E, Seto WK, Chan HL, Chuang WL, Stepanova T, Hui AJ, Lim YS, Mehta R, Janssen HL, Acharya SK, Flaherty JF, Massetto B, Cathcart AL, Kim K, Gaggar A, Subramanian GM, McHutchison JG, Pan CQ, Brunetto M, Izumi N, Marcellin P; GS-US-320-0108 Investigators. Tenofovir alafenamide versus tenofovir disoproxil fumarate for the treatment of patients with HBeAg-negative chronic hepatitis B virus infection: a randomised, double-blind, phase 3, non-inferiority trial. *Lancet Gastroenterol Hepatol* 2016; **1**: 196-206 [PMID: [28404092](#) DOI: [10.1016/S2468-1253\(16\)30107-8](#)]
- 7 **Agarwal K**, Brunetto M, Seto WK, Lim YS, Fung S, Marcellin P, Ahn SH, Izumi N, Chuang WL, Bae H, Sharma M, Janssen HLA, Pan CQ, Çelen MK, Furusyo N, Shalimar D, Yoon KT, Trinh H, Flaherty JF, Gaggar A, Lau AH, Cathcart AL, Lin L, Bhardwaj N, Suri V, Mani Subramanian G, Gane EJ, Buti M, Chan HLY; GS-US-320-0110; GS-US-320-0108 Investigators. 96 weeks treatment

- of tenofovir alafenamide vs. tenofovir disoproxil fumarate for hepatitis B virus infection. *J Hepatol* 2018; **68**: 672-681 [PMID: 29756595 DOI: 10.1016/j.jhep.2017.11.039]
- 8 **Chinese Society of Hepatology**; Chinese Medical Association; Chinese Society of Infectious Diseases; Chinese Medical Association. The guideline of prevention and treatment for chronic hepatitis B (2015 version). *Shiyong Ganzangbing Zazhi* 2016; **19**: 389-400 [DOI: 10.3760/cma.j.issn.1007-3418.2015.12.034]
 - 9 **Qi X**, Wang JY, Mao RC, Zhang JM. Impact of nucleos(t)ide analogues on the estimated glomerular filtration rate in patients with chronic hepatitis B: a prospective cohort study in China. *J Viral Hepat* 2015; **22**: 46-54 [PMID: 25402626 DOI: 10.1111/jvh.12229]
 - 10 **Qi X**, Wang J, Chen L, Huang Y, Qin Y, Mao R, Zhang J. Impact of nucleos(t)ide analogue combination therapy on the estimated glomerular filtration rate in patients with chronic hepatitis B. *Medicine* **94**: e646 [PMID: 25881837 DOI: 10.1097/MD.0000000000000646]
 - 11 **Shimohata H**, Sakai S, Ogawa Y, Hirayama K, Kobayashi M. Osteomalacia due to Fanconi's syndrome and renal failure caused by long-term low-dose adefovir dipivoxil. *Clin Exp Nephrol* 2013; **17**: 147-148 [PMID: 23268286 DOI: 10.1007/s10157-012-0762-8]
 - 12 **Gara N**, Zhao X, Collins MT, Chong WH, Kleiner DE, Jake Liang T, Ghany MG, Hoofnagle JH. Renal tubular dysfunction during long-term adefovir or tenofovir therapy in chronic hepatitis B. *Aliment Pharmacol Ther* 2012; **35**: 1317-1325 [PMID: 22506503 DOI: 10.1111/j.1365-2036.2012.05093.x]
 - 13 **Amarapurkar DN**, Patel N. Increased eGFR with telbivudine in combination therapy of chronic hepatitis B infection. *Indian J Gastroenterol* 2014; **33**: 89-91 [PMID: 23512213 DOI: 10.1007/s12664-013-0325-2]
 - 14 **Huo N**, Lu HY, Wang GQ, Huang MJ, Wu CH, Lin XH. Effects of nucleos(t)ide analogues on renal function in patients with chronic hepatitis B. *Zhongguo Linchuang Yaolixue Yu Zhilixue* 2017; **22**: 1029-1034



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

