

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 34,900 条结果 (用时 0.96 秒)

Google 学术: Risk stratification for malignant progression in Barrett's esophagus: gender, age, duration and year of surveillance

Risk of malignant progression in patients with Barrett's ... - de Jonge - 被引用次数: 185

Barrett esophagus: risk factors for progression to ... - Öberg - 被引用次数: 125

... , surveillance and therapy of Barrett's esophagus - Wang - 被引用次数: 1067

Risk stratification and surveillance in Barrett oesophagus - NCBI

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: EC Gorospe - 2014 - 被引用次数: 1 - 相关文章

2013年12月10日 - EAC has been the fastest rising malignancy in Western countries over ... patient had frequent reflux symptoms, beginning at 10 years of age. ... After adjusting for age, gender and length of the Barrett mucosa, the persistence of nondysplastic Barrett oesophagus was associated with low risk of progression to ...

Predictors of Progression in Barrett's Esophagus: Current Knowledge ...

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: GA Prasad - 2010 - 被引用次数: 103 - 相关文章

2010年1月26日 - Clinical and demographic factors (age, male gender, length of BE ... that of any other malignancy in the United States over the last few decades (1). ... a tool to risk stratify patients with BE in terms of their risk of progression to EAC. ... the mean age at initiation of surveillance in progressors (77.1 years) was ...

Barrett Esophagus - NCBI - National Institutes of Health

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: S Öberg - 2005 - 被引用次数: 124 - 相关文章

Risk Factors for Progression to Dysplasia and Adenocarcinoma ... The risk for patients with Barrett esophagus to develop esophageal adenocarcinoma is low, and most patients undergoing surveillance will not develop malignancy. were age, gender, body mass index (BMI), history of tobacco use, duration of surveillance, ...

Risk Stratification of Barrett's Esophagus: Updated Analysis - Medscape

www.medscape.com/viewarticle/489429 ▾ [翻译此页](#)

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 28859

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Risk stratification for malignant progression in Barrett's esophagus: Gender, age, duration and year of surveillance

Gatenby P *et al.* Risk stratification for malignant progression in Barrett's esophagus

Piers Gatenby, Santanu Bhattacharjee, Christine Wall, Christine Caygill, Anthony Watson

Piers Gatenby, Santanu Bhattacharjee, Christine Wall, Christine Caygill, Anthony Watson, UK Barrett's Oesophagus Registry, UCL, NW32QG London, United Kingdom

Piers Gatenby, Regional Oesophagogastric Unit, Royal Surrey County Hospital,

Match Overview

1	Internet 137 words crawled on 19-Jul-2016 www.wjgnet.com	5%
2	Crossref 34 words Piers A.C. Gatenby. "Short segment columnar-lined oesophagus: an underestimated cancer risk? A large cohort study"	1%
3	Crossref 25 words Piers A. C. Gatenby. "Treatment modality and risk of development of dysplasia and adenocarcinoma in columnar-lined"	1%

学术搜索

找到约 1,500 条结果 (用时0.07秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...


按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☒ 包含引用 创建快讯小提示: 只搜索 **中文(简体)** 结果, 可在 学术搜索设置 指定搜索语言**Risk of malignant progression in patients with Barrett's oesophagus: a Dutch nationwide cohort study**[\[PDF\] researchgate.net](#)

PJF de Jonge, M Van Blankenstein, CWN Looman... - Gut, 2010 - gut.bmj.com

... This observation remained unchanged after stratifying patients by age. ... cancer risk in patients with BO with baseline LGD without affecting the overall cancer risk. ... overclassification of BO by community pathologists could have led to underestimation of reported cancer risks. ...

被引用次数: 198 相关文章 所有 14 个版本 引用 保存

Barrett esophagus: risk factors for progression to dysplasia and adenocarcinoma[\[PDF\] researchgate.net](#)

S Öberg, J Wenner, J Johansson, B Walther... - Annals of ..., 2005 - journals.lww.com

... Low-grade dysplasia is the only clinically useful risk factor that permits stratification of the ... To develop more rational surveillance programs, patients need to be stratified according to their individual ... antireflux surgery (1 in 294.4 patient-years) was lower than the risk for Barrett's ...

被引用次数: 127 相关文章 所有 12 个版本 引用 保存

Updated guidelines 2008 for the diagnosis, surveillance and therapy of Barrett's esophagus[\[PDF\] alabmed.com](#)

KK Wang, RE Sampliner - The American journal of ..., 2008 - search.proquest.com

... It is recommended that patients be advised of the benefits and risks of surveillance ... The ideal biomarker panel to risk stratify patients with BE would be noninvasive and relatively ... Non-steroidal anti-inflammatory drugs and risk of neoplastic progression in Barretts oesophagus: a ...

被引用次数: 1101 相关文章 所有 20 个版本 引用 保存

[引用] Leukocyte telomere length predicts cancer risk in Barrett's esophagus[Free from Publisher](#)

RA Risques, TL Vaughan, X Li, RD Odze... - ..., Biomarkers & Prevention, 2007 - AACR

被引用次数: 129 相关文章 所有 5 个版本 引用 保存

Predictors for neoplastic progression in patients with Barrett's esophagus: a prospective cohort study[\[HTML\] nature.com](#)

M Sikkema, CWN Looman, EW Steyerberg... - The American journal of ..., 2011 - nature.com

... to the development of esophageal adenocarcinoma (EAC) (3,4,5). This malignancy has a ... is able to identify patients with a high or a low risk of neoplastic progression. Risk stratification and

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▼](#)[搜索工具](#)

找到约 32,200 条结果 (用时 0.77 秒)

Google 学术 : Risk stratification for malignant progression in Barrett's esophagus: Gender, age, duration and year of surveillance

Risk of malignant progression in patients with Barrett's ... - de Jonge - 被引用次数 : 199

Barrett esophagus: risk factors for progression to ... - Öberg - 被引用次数 : 127

... , surveillance and therapy of Barrett's esophagus - Wang - 被引用次数 : 1102

Barrett Esophagus - NCBI - National Institutes of Health

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : S Öberg - 2005 - 被引用次数 : 127 - 相关文章

Risk Factors for Progression to Dysplasia and Adenocarcinoma ... are stratified according to their individual risk of progressing to dysplasia and invasive ... histologic surveillance in patients with Barrett esophagus to detect malignancy at an early ... were age, gender, body mass index (BMI), history of tobacco use, duration of ...

Risk stratification and surveillance in Barrett oesophagus - NCBI

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▼ 翻译此页

作者 : EC Gorospe - 2014 - 被引用次数 : 1 - 相关文章

2013年12月10日 - Risk stratification and surveillance in Barrett oesophagus ... during surveillance is associated with low risk of malignant progression, ... After adjusting for age, gender and length of the Barrett mucosa, the persistence of ...

Diagnosis and Management of Barrett's Esophagus

www.ncbi.nlm.nih.gov > Journal List > HHS Author Manuscripts - 翻译此页

作者 : EM Nelsen - 2012 - 被引用次数 : 11 - 相关文章

2012年8月20日 - The absolute annual risk of progression to EAC remains a subject of debate. ... have shown comparable life expectancy (to age-matched and gender-matched ... to cancer development replaces the stratified squamous epithelium that Screening and surveillance for Barrett esophagus in high-risk groups: ...