



24056-Review

Quotes Excluded
Bibliography Excluded17%
SIMILARName of Journal: *World Journal of Hepatology*

ESPS Manuscript NO: 24056

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

3 Risk factors for deterioration of long-term liver function after radiofrequency ablation therapy

2 Koichi Honda, Masataka Seike, Junya Oribe, Mizuki Endo, Mie Arakawa, Hiroki Syo, Masao Iwao, Masanori Tokoro, Junko Nishimura, Tetsu Mori, Tsutomu Yamashita, Satoshi Fukuchi, Toyokichi Muro, Kazunari

Match Overview



| | | |
|----------|---|----|
| 1 | CrossCheck 95 words "AASLD Abstracts", Hepatology, 2013. | 3% |
| 2 | Internet 54 words crawled on 08-Nov-2014 www.aasld.org | 2% |
| 3 | CrossCheck 46 words Hidekatsu Kuroda. "Changes in liver function parameters after percutaneous radiofrequency ablation therapy in pat ... | 1% |
| 4 | CrossCheck 44 words "AASLD Abstracts", Hepatology, 2012. | 1% |
| 5 | CrossCheck 35 words "Liver Cancer", Hepatology, 2013. | 1% |
| 6 | Internet 34 words crawled on 08-Mar-2016 www.wjgnet.com | 1% |

[全部](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 52,300 条结果 (用时 0.88 秒)

Antiviral therapies for hepatitis B virus-related hepatocellular ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

作者: YQ Zhang - 2015 - 被引用次数: 2 - 相关文章

2015年4月7日 - Chronic hepatitis B virus (HBV) infection is a critical risk factor for the carcinogenesis ... survival of HBV-related HCC patients after curative and palliative therapies. ... Hepatocellular carcinoma (HCC) is the most common form of liver cancer. The prolonged fibrogenic response is accompanied by regional ...

Thrombocytopenia represents a risk for deterioration of liver ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: HS Lee - 2012 - 被引用次数: 6 - 相关文章

2012年9月25日 - Liver function had deteriorated 12 months after RFA in 29 patients (28.4%). ... A pre-RFA thrombocytopenia represents a major risk factor for the deterioration ... Anti-viral therapies in patients with chronic hepatitis B were permitted. tumor ablation for hepatocellular carcinoma: long-term follow up for 227 ...

Hepatitis B viral load affects prognosis of hepatocellular ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: SJ Yu - 2014 - 被引用次数: 8 - 相关文章

2014年9月14日 - Hepatitis B virus (HBV) is an established major risk factor of HCC ... patients due to cancer progression, it is expected that long-term ... HCC after treatment

[全部](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 57,000 条结果 (用时 1.10 秒)

The influence on liver function after transcatheter arterial ...

www.sciencedirect.com/science/article/.../S0929664612000770 ▾ [翻译此页](#)

作者: JX Li - 2012 - 被引用次数: 19 - 相关文章

Hepatocellular carcinoma (HCC) ranks fifth among the most common malignant tumors ... Biochemical changes of **liver function** occur in **patients after** resection, **RFA** or TACE. ... a major **risk factor** for the aggravation of **liver function after RFA therapy**. ... The **ablation time** was **related** to tumor size and proximity to vasculature.

Thrombocytopenia represents a risk for deterioration of liver ...

www.ncbi.nlm.nih.gov > ... > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: HS Lee - 2012 - 被引用次数: 6 - 相关文章

2012年9月25日 - **Liver function** had **deteriorated** 12 months **after RFA** in 29 **patients** (28.4%). ... A pre-**RFA** thrombocytopenia represents a major **risk factor** for the **deterioration** ... Anti-viral **therapies** in **patients** with chronic **hepatitis B** were permitted. tumor **ablation** for **hepatocellular carcinoma**; long-term follow up for 227 ...

Antiviral therapies for hepatitis B virus-related hepatocellular ...

www.ncbi.nlm.nih.gov > ... > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: YQ Zhang - 2015 - 被引用次数: 2 - 相关文章

2015年4月7日 - Chronic **hepatitis B virus** (HBV) infection is a critical **risk factor** for the carcinogenesis ... survival of HBV-**related HCC patients after** curative and palliative

[全部](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 48,300 条结果 (用时 0.72 秒)

Google 学术: Risk factors for deterioration of long-term liver function after radiofrequency ablation therapy

... Percutaneous Image-guided Radiofrequency Ablation ... - Lencioni - 被引用次数: 687

Long-term survival and pattern of recurrence after ... - Poon - 被引用次数: 623

... for hepatocellular carcinoma: risk factors, treatment, ... - Shah - 被引用次数: 226

The influence on liver function after transcatheter arterial ...

www.sciencedirect.com/science/article/.../S0929664612000770 ▾ 翻译此页

作者: JX Li - 2012 - 被引用次数: 19 - 相关文章

Therefore, locoregional therapies such as percutaneous ethanol injection (PEI), ... unresectable HCC, and these treatments are often associated with a satisfactory long-term prognosis. ... The ablation time was related to tumor size and proximity to vasculature. Risk factors for deterioration of liver function after TACE+RFA.

Thrombocytopenia represents a risk for deterioration of liver ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: HS Lee - 2012 - 被引用次数: 6 - 相关文章

2012年9月25日 - Liver function had deteriorated 12 months after RFA in 29 patients (28.4%). ... function parameters after RFA therapy and identified risk factors for aggravation of Significant long-term survival after radiofrequency ablation of ...

Branched Chain Amino Acids in Clinical Nutrition

<https://books.google.com.hk/books?isbn=1493919148> - 翻译此页

Rajkumar Rajendram, Victor R. Preedy, Vinood B. Patel - 2014 - Medical

BCAA-enriched amino acid infusion solution is effective in the treatment of HE In the