

**Name of Journal:** *World Journal of Hepatology*

**ESPS Manuscript NO:** 30550

**Manuscript Type:** Minireviews

**Limitations and opportunities of non-invasive liver stiffness measurement in children**

**Guido Engelmann, Jasmin Quader, Ulrike Teufel, Jens Peter Schenk**

### Abstract

Changes in liver structure are an important issue in chronic hepatopathies. Until the end of the 20<sup>th</sup> century, these changes could only be determined by histological analyses of a liver specimen obtained *via* biopsy. The well-known limitations of this

### Match Overview



1

**Crossref** 157 wordsGuido Engelmann, "Feasibility study and control values of transient elastography in healthy children", *European Journal of*

4%

2

**Crossref** 92 words

Schenk, Jens-Peter, Buket Selmi, Christa Flechtenmacher, S. araa El Sakka, Ulrike Teufel, and Guido Engelmann, "Real-time

3%

3

**Internet** 10 words

crawled on 04-May-2016

[hepatistresearchandnewsupdates.blogspot.co.nz](http://hepatistresearchandnewsupdates.blogspot.co.nz)

&lt;1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 1,380,000 条结果 (用时 0.70 秒)

## Google 学术: Limitations and opportunities of non-invasive liver stiffness measurement in children

Non-invasive evaluation of liver fibrosis using transient ... - Castera - 被引用次数: 789

Liver stiffness measurement in children using ... - de Lédinghen - 被引用次数: 175

Diagnosis of cirrhosis by transient elastography ( ... - Foucher - 被引用次数: 1068

## Non-invasive assessment of liver fibrosis

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: V Papastergiou - 2012 - 被引用次数: 46 - 相关文章

In parallel with this, noninvasive assessment of liver fibrosis has experienced ... Due to these limitations, consideration of liver biopsy as the "gold standard" has ..... Importantly, in a more recent report, repeated liver stiffness measurements ... or in Child C cirrhosis) allowing to avoid about 33% of upper GI endoscopies [167].

## Application and limitations of transient liver elastography in children.

<https://www.ncbi.nlm.nih.gov/pubmed/23539048> - 翻译此页

作者: I Goldschmidt - 2013 - 被引用次数: 37 - 相关文章

Application and limitations of transient liver elastography in children. ... has gained popularity recently for the noninvasive diagnosis of hepatic fibrosis. ... such as probe choice and site of measurement on FibroScan results is not clear. ... General anesthesia significantly increased liver stiffness in healthy children (5.4 [3-9.5] ...

## Application And Limitations of Transient Liver Elastography In Children.

<https://www.researchgate.net/.../236090466> Application And Limitations o ... - 翻译此页

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 1,390,000 条结果 (用时 0.72 秒)

## Google 学术: Limitations and opportunities of non-invasive liver stiffness measurement in children

Non-invasive evaluation of liver fibrosis using transient ... - Castera - 被引用次数: 794

Liver stiffness measurement in children using ... - de Lédinghen - 被引用次数: 175

Diagnosis of cirrhosis by transient elastography ( ... - Foucher - 被引用次数: 1068

## Non-invasive assessment of liver fibrosis

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: V Papastergiou - 2012 - 被引用次数: 46 - 相关文章

In parallel with this, noninvasive assessment of liver fibrosis has experienced ... Due to these limitations, consideration of liver biopsy as the "gold standard" has .... Importantly, in a more recent report, repeated liver stiffness measurements ... or in Child C cirrhosis) allowing to avoid about 33% of upper GI endoscopies [167].

## Application and limitations of transient liver elastography in children.

<https://www.ncbi.nlm.nih.gov/pubmed/23539048> - 翻译此页

作者: I Goldschmidt - 2013 - 被引用次数: 37 - 相关文章

Application and limitations of transient liver elastography in children. ... has gained popularity recently for the noninvasive diagnosis of hepatic fibrosis. ... such as probe choice and site of measurement on FibroScan results is not clear. ... General anesthesia significantly increased liver stiffness in healthy children (5.4 [3-9.5] ...