

World Journal of *Diabetes*

World J Diabetes 2021 September 15; 12(9): 1363-1586



REVIEW

- 1363 Conundrum of vitamin D on glucose and fuel homeostasis
Chang Villacreses MM, Karnchanasorn R, Panjawatanan P, Ou HY, Chiu KC
- 1386 Review of the management of sight-threatening diabetic retinopathy during pregnancy
Choo PP, Md Din N, Azmi N, Bastion MLC
- 1401 Epidemiology of type 2 diabetes in the Middle East and North Africa: Challenges and call for action
El-Kebbi IM, Bidikian NH, Hneiny L, Nasrallah MP
- 1426 Dipeptidyl peptidase-4 inhibitor-induced autoimmune diseases: Current evidence
Roy A, Sahoo J, Narayanan N, Merugu C, Kamalanathan S, Naik D
- 1442 Cellular targets in diabetic retinopathy therapy
Rodríguez ML, Millán I, Ortega ÁL
- 1463 Holistic perspective of the role of gut microbes in diabetes mellitus and its management
Alagiakrishnan K, Halverson T
- 1479 Non-alcoholic fatty liver disease in diabetes: When to refer to the hepatologist?
Khandelwal R, Dassanayake AS, Conjeevaram HS, Singh SP

MINIREVIEWS

- 1494 Improving nutrition for the prevention of gestational diabetes: Current status and perspectives
Popova PV, Pustozarov EA, Tkachuk AS, Grineva EN
- 1507 Salivary resistin level and its association with insulin resistance in obese individuals
Abdalla MMI
- 1518 Psychosocial factors affecting the etiology and management of type 1 diabetes mellitus: A narrative review
Turin A, Drobnič Radobuljac M
- 1530 Sugar intake from sweetened beverages and diabetes: A narrative review
Tseng TS, Lin WT, Gonzalez GV, Kao YH, Chen LS, Lin HY
- 1539 Role of an acidic environment in the treatment of diabetic foot infections: A review
Nagoba B, Gavkare A, Rayate A, Mumbre S, Rao A, Warad B, Nanaware N, Jamadar N
- 1550 Diabetes and COVID-19: Role of insulin resistance as a risk factor for COVID-19 severity
Gangadharan C, Ahluwalia R, Sigamani A

- 1563** Obesity and bariatric surgery in kidney transplantation: A clinical review

Veroux M, Mattone E, Cavallo M, Gioco R, Corona D, Volpicelli A, Veroux P

META-ANALYSIS

- 1576** Effectiveness of drug interventions in nonalcoholic fatty liver disease: A network meta-analysis

Huang YZ, Yang GY, Wang C, Chen XY, Zhang LL

ABOUT COVER

Editorial Board Member of *World Journal of Diabetes*, Jayaprakash Sahoo, MBBS, MD, Associate Professor, Department of Endocrinology, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Puducherry 605006, India. jppgi@yahoo.com

AIMS AND SCOPE

The primary aim of *World Journal of Diabetes* (WJD, *World J Diabetes*) is to provide scholars and readers from various fields of diabetes with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJD mainly publishes articles reporting research results and findings obtained in the field of diabetes and covering a wide range of topics including risk factors for diabetes, diabetes complications, experimental diabetes mellitus, type 1 diabetes mellitus, type 2 diabetes mellitus, gestational diabetes, diabetic angiopathies, diabetic cardiomyopathies, diabetic coma, diabetic ketoacidosis, diabetic nephropathies, diabetic neuropathies, Donohue syndrome, fetal macrosomia, and prediabetic state.

INDEXING/ABSTRACTING

The WJD is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, and PubMed Central. The 2021 Edition of Journal Citation Reports® cites the 2020 impact factor (IF) for WJD as 3.763; IF without journal self cites: 3.684; 5-year IF: 7.348; Journal Citation Indicator: 0.64□Ranking: 80 among 145 journals in endocrinology and metabolism; and Quartile category: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Lin-YuTong Wang; Production Department Director: Yu-Jie Ma; Editorial Office Director: Jia-Ping Yan.

NAME OF JOURNAL

World Journal of Diabetes

ISSN

ISSN 1948-9358 (online)

LAUNCH DATE

June 15, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Lu Cai, Md. Shahidul Islam, Jian-Bo Xiao, Manfredi Rizzo

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/1948-9358/editorialboard.htm>

PUBLICATION DATE

September 15, 2021

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Holistic perspective of the role of gut microbes in diabetes mellitus and its management

Kannayiram Alagiakrishnan, Tyler Halverson

ORCID number: Kannayiram Alagiakrishnan 0000-0003-1444-2451; Tyler Halverson 0000-0001-5841-7659.

Author contributions:

Alagiakrishnan K and Halverson T contributed equally by reviewing the literature and drafting the manuscript; both the authors have read and approved the final version.

Conflict-of-interest statement: The authors declare no potential conflict of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Specialty type: Endocrinology and metabolism

Kannayiram Alagiakrishnan, Tyler Halverson, Department of Medicine, University of Alberta, Edmonton T6G 2G3, Alberta, Canada

Corresponding author: Kannayiram Alagiakrishnan, FRCPC, MD, Full Professor, Medicine, University of Alberta, B139-152 University Hospital Clinical Sciences Building, Edmonton T6G 2G3, Alberta, Canada. kalagiakri@aol.com

Abstract

The gut microbiota (GM) plays a role in the development and progression of type 1 and type 2 diabetes mellitus (DM) and its complications. Gut dysbiosis contributes to the pathogenesis of DM. The GM has been shown to influence the efficacy of different antidiabetic medications. Intake of gut biotics, like prebiotics, probiotics and synbiotics, can improve the glucose control as well as the metabolic profile associated with DM. There is some preliminary evidence that it might even help with the cardiovascular, ophthalmic, nervous, and renal complications of DM and even contribute to the prevention of DM. More large-scale research studies are needed before wide spread use of gut biotics in clinical practice as an adjuvant therapy to the current management of DM.

Key Words: Probiotics; Prebiotics; Synbiotics; Diabetes mellitus; Microbial dysbiosis; Antidiabetic drugs

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The emerging role of the gut microbiome on diabetes development, progression as well as prevention has been discussed in this manuscript. The significance of gut dysbiosis in the aetiopathogenesis of diabetes mellitus and its complications has been reviewed. A bidirectional relationship exists between the antidiabetic drugs and the gut microbiome. Faecal transplantation, and bariatric surgery, typically used to treat morbid obesity, have also been shown to improve commensal gut microbiota changes. Diabetic outcomes and management can improve with better understanding of the drug-gut microbiome interactions. There is emerging evidence pointing out that gut biotics can be an add-on therapy with the antidiabetic management. To our knowledge, there is no evidence about the role of gut microbes of diabetic patients who had pancreatic cell transplantation, as well as the role of gut

Country/Territory of origin: Canada**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

Received: February 26, 2021**Peer-review started:** February 26, 2021**First decision:** April 20, 2021**Revised:** April 24, 2021**Accepted:** August 13, 2021**Article in press:** August 13, 2021**Published online:** September 15, 2021**P-Reviewer:** Park SC**S-Editor:** Wang JL**L-Editor:** A**P-Editor:** Xing YX

biotics influencing the management in this group.

Citation: Alagiakrishnan K, Halverson T. Holistic perspective of the role of gut microbes in diabetes mellitus and its management. *World J Diabetes* 2021; 12(9): 1463-1478**URL:** <https://www.wjnet.com/1948-9358/full/v12/i9/1463.htm>**DOI:** <https://dx.doi.org/10.4239/wjd.v12.i9.1463>

INTRODUCTION

Globally diabetes mellitus (DM) is a common medical disorder and is seen in pandemic proportions[1] with the global prevalence in adult subjects is roughly 10% [2]. The International Diabetes Federation projected by 2035, there will be 592 million cases of diabetes in the world[3]. DM type 1 is secondary to auto-immune-mediated loss of beta-cell function and is seen in 5% of the diabetic population. DM type 2 is mainly due to insulin resistance and is seen in 95% of diabetic subjects[4]. The 2016 US National Health Interview Survey data showed roughly 8.58% of the population had type 2 DM and 0.55% had type 1 DM[5].

Various research has been done in the last decade since the study of the human microbiome in 2012[6,7]. Microbes contribute to 2% of human body weight and the bacterial genomes exceeds human genes by a factor of 150[8,9]. Gut microbiota (GM) varies with age, diet, geographical location, life style, and the use of xenobiotics[10-12]. In the recent years there have been more focus on the GM in the development, progression, and distant organ complications due to DM[13]. Many studies have shown the role of the gut microbiome in DM[14-17].

The gut microbiome starts to develop with the mode of birth and it is influenced by environmental factors, diet, as well as certain medications, including antibiotics[18]. There are differences between the gut microbes seen between non-diabetic and diabetic subjects[20] (Table 1). Gut dysbiosis plays a role in numerous diseases including DM. Both altered GM and endocrine disrupters can influence the development of DM[21]. In this literature review, we analyzed the evidence for the role of GM in the development, pathogenesis, complications, management, and prevention of DM.

LITERATURE SEARCH

A literature search was performed using the electronic databases MEDLINE (1966–February 2021), EMBASE and SCOPUS (1965–February 2021), and DARE (1966–February 2021). The main search items were gut bacteria, GM, intestinal flora, gut dysbiosis, type 1 DM, type 2 DM, diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, probiotics, prebiotics, synbiotics, bariatric surgery, and faecal transplantation. Non-English articles were excluded.

GM IN TYPE 1 DM

Studies have shown that Firmicutes/Bacteroides ratio is altered in type 1 DM[22]. In the study by Huang *et al*[23] (2018) negative association was seen with gut microbe *Faecalibacterium* and *Ruminococcaceae* and hemoglobin A1c (HbA1c), whereas in the study by Fassatoui *et al*[24]. (2019) a negative association was seen between HbA1c and *Akkermansia muciniphila*. A systematic review of studies done in Hispanic populations showed that patients with newly diagnosed type 1 DM have high levels of *Bacteroides* with a reduced proportion of *Prevotella*, *Megamonas*, and *Acidaminococcus*. With the initiation of insulin treatment these subjects showed an increase of *Prevotella* levels. Prior to the development of type 1 DM, inverse relationship of Firmicutes/Bacteroidetes ratio has been reported[25].

Table 1 Changes in the microbiome in type 1 and type 2 diabetes mellitus

Location	Change in microbiome	Ref.
Type 1 diabetes		
Gastrointestinal tract	(1) Decreased: <i>Prevotella</i> ; <i>Megamonas</i> ; <i>Acidaminococcus</i> ; and (2) Increased: <i>Bacteriodes</i>	Elena <i>et al</i> [25], 2019
Gastrointestinal tract	(1) Decreased: <i>Bifidobacterium adolescentis</i> ; <i>Bifidobacteria</i> ; and (2) Increased: <i>Clostridium perfringens</i> ; <i>Bacteroides</i>	De Goffau <i>et al</i> [122], 2013
Gastrointestinal tract	Increased: <i>Leptotrichia goodfellowii</i> ; <i>Bacillus cerus</i> ; <i>Enterobacter mori</i> LMG 25706	Tai <i>et al</i> [123], 2016
Gastrointestinal tract	Increased: <i>Bacteroidetes/Firmicutes</i>	Giongo <i>et al</i> [124], 2011
Gastrointestinal tract	(1) Decreased: <i>Faecalibacterium prausnitzii</i> ; and (2) Increased: <i>Bacteroides dorei</i> ; <i>Bacteroides vulgatus</i>	De Goffa <i>et al</i> [125], 2014
Gastrointestinal tract	(1) Decreased: <i>Prevotella</i> ; <i>Akkermansia</i> ; <i>Bifidobacterium adolescentis</i> ; <i>Roseburia faecis</i> ; <i>Faecalibacterium prausnitzii</i> ; and (2) Increased: <i>Dialister invisus</i> ; <i>Gemella sanguinis</i> ; <i>Difidobacterium longum</i>	Brown <i>et al</i> [126], 2011
Type 2 diabetes		
Gastrointestinal tract	(1) Decreased: <i>Clostridium coccoides</i> ; <i>Clostridium leptum</i> ; and (2) Increased: <i>Lactobacillus</i>	Chen <i>et al</i> [28], 2019
Gastrointestinal tract	(1) Decreased: <i>Bifidobacterium</i> ; <i>Bacteroides</i> ; <i>Faecalibacterium</i> ; <i>Akkermansia</i> ; <i>Roseburia</i> ; and (2) Increased: <i>Ruminococcus</i> ; <i>Fusobacterium</i> ; <i>Blautia</i>	Gurung <i>et al</i> [30], 2020
Gastrointestinal tract	(1) Decreased: <i>Bifidobacterium</i> ; <i>Akkermansia</i> ; and (2) Increased: <i>Dorea</i>	Li <i>et al</i> [127], 2020
Gastrointestinal tract	(1) Decreased: <i>Bifidobacterium</i> ; and (2) Increased: <i>Lactobacillus</i>	Sedighi <i>et al</i> [31], 2017
Blood	(1) Decreased: <i>Aquabacterium</i> ; <i>Xanthomonas</i> ; <i>Pseudonocardia</i> ; and (2) Increased: <i>Actinotalea</i> ; <i>Alishewanella</i> ; <i>Seiminibacterium</i> ; <i>Pseudoclavibacter</i>	Qiu <i>et al</i> [38], 2019

GM IN TYPE 2 DM

The type of gut microbes and the changes seen with them influence the development of DM. The prominent GM seen in the intestine are the gram-positive Firmicutes and gram-negative Bacteroidetes and it is influenced by dietary changes[26]. A change in the ratio of Bacteroidetes to Firmicutes is associated with DM[27]. A case-control study done by Chen *et al*[28] (2019) in newly diagnosed type 2 DM subjects, *Lactobacillus* faecal count was significantly higher whereas *Clostridium coccoides* and *Clostridium leptum* was lower, and these changes in the microbes was positively correlated with glycated hemoglobin with higher *Lactobacillus* count subjects, and negatively correlated with decreased *Clostridium* count subjects when compared with healthy controls. Another study found that patients with DM showed an affiliation with the following phyla of bacteria: Firmicutes, Bacteroidetes, Proteobacteria and Actinobacteria[29]. Alterations in the gut microbe population may be related to DM, and gut microbes *Ruminococcus* and *Fusobacterium* has been shown with the development of type 2 DM, when compared to healthy adults[30]. A study by Sedighi *et al*[31] (2017) found that patients with type 2 DM has increased levels of *Lactobacillus*, while healthy controls showed increased *Bifidobacterium*. With respect to the *Lactobacillus* genus, there are various mixed results suggesting its association with type 2 DM. Certain strain such as *L. acidophilus*, *L. gasseri*, and *L. salivarius* have been increased where as *L. amyloyorus* has been decreased[30]. However, many species from this genus, such as *L. plantarum*, *L. casei*, and *L. rhamnosus* are often involved in probiotic preparation and have shown to be beneficial in diabetic mice models[30]. Overall, it looks that there may be a strain-specific association with DM.

Further changes in the microbiome in patients with DM are listed in Table 1. Nutrient imbalance by affecting the GM can influence the development of type 2 DM. With newly diagnosed type 2 DM different measurement parameters like age, blood lipids, body-mass index, blood pressure, and dietary nutrient intake was related to the gut microbiome composition[32].

RELATIONSHIP BETWEEN GUT AND BLOOD MICROBIOME AND ITS ASSOCIATION WITH TYPE 2 DM

Cani *et al*[33] (2008) in their animal study showed lipopolysaccharide produced by gram negative intestinal bacteria can translocate into systemic circulation through a leaky gut and can result in endotoxemia causing metabolic dysfunction and obesity. Recent evidence points out in addition to gut microbiome, the blood microbiome plays a role in DM. Blood is usually considered to be sterile, but the research suggests the presence of a microbe or microbial component in healthy humans is known as a blood microbiome. The evidence for blood human microbiome is slowly growing[34-36].

In a study by Sato *et al*[37] (2014) with Japanese type 2 DM subjects, blood microbiome translocation from gut microbiome was detected at a higher rate (28%) in type 2 diabetic subjects when compared with healthy controls (4%) ($P < 0.01$). A recent nested case control study by Qiu *et al*[38] (2019) showed the blood microbe *Sediminibacterium* is associated with increased risk of type 2 DM [Odd ratio (OR) = 14.098, 95%CI: 1.358-146.330] whereas the microbe *Bacteroides* in blood have a reduced risk for type 2 DM (OR = 0.367, 95%CI: 0.151- 0.894).

GM AS A COMPLEX ENDOCRINE ORGAN

The regulation of the GI system is done by short-chain fatty acids (SCFA) derived from the metabolism of carbohydrates, and GM plays a role in this function. In addition, the gut microbes produce hormone like chemicals that can act at distant targets. Neuroactive compounds like tryptophan and neurotransmitters like serotonin, dopamine, noradrenaline, GABA, and hormones like leptin, ghrelin and glucagon-like peptide 1 (GLP-1) are indirectly regulated by SCFAs *via* enteroendocrine cells. Overall, the gut microbes produce several substances of a hormonal nature into the circulation which act as distant sites. Because of the GM's ability to influence distant organs and systems as mentioned above it is considered as an endocrine organ. Overall GM functions as an autonomous endocrine organ and plays a role in bodily endocrine actions including neuroendocrine and immunoendocrine regulations[39-42].

DM AND GUT DYSBIOSIS

Gut dysbiosis, is a state of increased or altered prevalence of gut bacteria which might in turn result in many disorders such as gastrointestinal, obesity, DM, immunological, and neurobehavioral diseases[43]. Shifts in the GM's composition with more pathogenic species and phyla can contribute to the above-mentioned diseases. Hyperglycemia was associated with changes of microbiota composition, preferring the non-commensal ones, on the detriment of beneficial phyla such as Bacteroidetes, Proteobacteria, and Actinobacteria. The ratio of Firmicutes/Bacteroidetes has been found to be correlated with plasma glucose concentration. Microbiota are capable to ferment undigested carbohydrates, fiber, and other dietary and xenobiotic compounds to produce SCFAs, which through their ubiquitous receptor play an important role in host glucose metabolism[37,44,45]. The Human Microbiome plays a role in gut permeability, modification of bile acids, glucose breakdown and in the absorption of nutrients[46,47].

Normal commensal bacteria are helpful in maintaining the gut wall integrity, innate immunity, insulin sensitivity, metabolism, and in communication with the brain functions, as well as help to prevent the penetration of harmful microorganisms in the bowel. Bidirectional relationship exists between the GM and the brain. This chain of communication depends on the interaction of gut microbe through immune and neuroendocrine system with the central nervous system. Short-chain fatty acids, such as butyrates, acetates and propionates, produced by the GM are beneficial to different metabolic processes. The imbalance between the microbiome and host organism lead to dysbiosis. Gut microbiome dysbiosis through inflammation and metabolic dysregulation increases insulin resistance and influence the development of type 2 DM[48] (Figure 1).

Microbial dysbiosis can also be the result of nutritional imbalance which can lead to a low-grade inflammatory state, obesity, and other metabolic disorders[49]. Gut microbes affect gut permeability, glucose and lipid metabolism, energy homeostasis, and insulin sensitivity. Like any other medical conditions, gut microbes play a role in




Figure 1 The role of gut dysbiosis in diabetes mellitus. The ingestion of a diet rich in carbohydrates and fats along with certain xenobiotics can lead to a disruption of the gut microbiome (dysbiosis). Under normal conditions, the gut bacteria produce metabolic products such as short chain fatty acids (SCFA) (Hexagons) that act locally and have a positive benefit on metabolism. Under conditions of dysbiosis there can be a disruption to the enteroendocrine cells and lead to gut permeability. This can lead to an increase in these metabolic products as well as bacterial translocation to the bloodstream, leading to endotoxemia resulting in metabolic dysfunction and insulin resistance contributing to type 2 diabetes. Gut dysbiosis also results in altered production of SCFA and release of lipopolysaccharides (LPS) (Triangles) and an increase production of other metabolites such as imidazole propionate and bacteria derived amino acids. These metabolites can act directly to affect insulin resistance. Excess SCFA and LPS can act on hepatic, skeletal, adipose, and pancreatic cells leading to metabolic dysfunction, altered inflammation and immune response which can influence insulin resistance. These factors can contribute to the development of type 1 and type 2 diabetes. SCFA: Short chain fatty acids; EEC: Enteroendocrine cells; LPS: Lipopolysaccharides; BCAA: Bacteria derived amino acids; IMP: Imidazole propionate; GM: Gut microbiota.

inflammation and immunity[50]. A diet rich in fat and sugar may lead to an abundance of lipopolysaccharide (LPS) release from GM and this LPS, by entering into systemic circulation, can affect β -cells, leading to decreased insulin release, and thereby altering systemic insulin sensitivity, resulting in insulin resistance, and potentially leading to DM[51].

Diets rich in carbohydrates and fat as well as xenobiotics (medications affecting the gut microbes) can cause gut dysbiosis. Normally GM produces metabolic products like SCFA, acetate, butyrate and propionate which acts locally leading to beneficial effects on different metabolic process. When there is gut dysbiosis, it can affect the enteroendocrine L-type cells in the intestinal epithelium and increase the gut permeability (leaky gut) causing these metabolic products to enter into the systemic circulation, as well as translocation of the gut microbiome into the circulation leading to the formation of the blood microbiome. This blood microbiome can cause endotoxemia and affect both metabolic dysfunction and insulin resistance. Gut dysbiosis results in excessive production of SCFA and LPS, as well as additional GM metabolites like imidazole propionate (IMP), derived from histidine, and bacteria derived amino acids. Excessive SCFA and LPS by acting on hepatic, adipose, skeletal and pancreatic cells causes metabolic dysfunction, inflammation and altered immune response. When there is a metabolic dysfunction due to gut dysbiosis combined with inflammatory and altered immune response it can cause type 1 DM, and when combined with insulin resistance due to gut dysbiosis as well as the effect of blood microbiome it can lead to the development of type 2 DM (Figure 1).

GUT MICROBES AND METABOLIC NETWORKS

The human gut contains a wide variety of microbial communities that carry out a wide range of biochemical functions that can influence the human body through metabolite production, physiological regulation, and interacting with the host's cellular response and immunity[52]. It has also been found that the host's own genetics can influence the composition of their gut microbiome, making each host a unique ecosystem[53]. Dynamic changes in the gut microbiome have been seen within individuals often in various disease states, such as obesity, and DM[19,54-56]. The GM has been found to cause enhanced transcriptional changes in the intestinal cells and protein biosynthesis in the crypts within the intestine[57].

SCFAs produced by GM can serve as signaling molecules that can influence the host's lipid and glucose levels, liver, skeletal muscle, and even immunity[52]. When there is a disruption of the gut microbiome, the altered mixture of SCFA may influence obesity, insulin sensitivity, weight gain and other comorbidities[58,59]. Obese individuals with type 2 DM have shown changes in the GM that are distinct, from non-diabetic subjects. It was found that individuals with type 2 DM showed an increase level of *Proteobacteria* and *Bacteroides* with a decreased level of *Firmicutes*[19].

The GM has been found to influence the host's metabolism and show great adaptability to the changing environment within the intestines based on diet, genetics, and various physiological cues from the host[52]. The human gut microbiome can modulate absorption as well as nutrient availability within the host. This can be achieved through gene expression changes, alteration of hormones and immunity[52].

ASSOCIATION BETWEEN MICROBIOME, OBESITY AND DM

Microbial diversity and the production of SCFA as well as products such as butyrate, propionate, and acetate have been found to have a protective role against obesity and insulin resistance[60,61]. SCFAs are able to act as signaling molecule that can activate a variety of pathways that are involved in cholesterol, lipid, and glucose metabolism [58]. Modifications of the microbiome can influence metabolic parameters, in particular when there is a higher abundance of Firmicutes leading to a higher Firmicutes/Bacteroidetes ratio that may be linked to obesity[62]. This may in part be due to the fact that Firmicutes are more efficient at promoting the nutrient absorption leading to subsequent weight gain compared to Bacteroidetes[63].

A study showed the GM composition is different in obese subjects with and without type 2 DM[20]. A recent study also showed for the first time in subjects with type 2 DM the relationship between body composition and GM[64]. Faecal microbiota of obese subjects without DM had increased numbers of SCFA producing microbes, whereas obese subjects with type 2 DM had less beneficial SCFA butyrate producing microbes[65].

ROLE OF GUT MICROBES IN THE PROGRESSION OF DM

The progression of DM is seen as macrovascular[66] and microvascular complications like retinopathy, nephropathy, and neuropathy[67]. Gut microbes seem to play a role in the progression of DM and also shown to play a role in these complications. Diet induced diabetic animal models helps to study these complications[68]. Studies have shown that subjects with DM and eye complications have higher bacterial conjunctival flora when compared to subjects without DM[69-72]. Beli *et al*[73] (2018) in their animal study showed the association between the GM and diabetic retinopathy (Table 2). More research is needed to understand the mechanism how GM causes diabetic retinopathy[74].

Diabetic neuropathy is seen as autonomic neuropathy as well as distal sensory and motor neuropathy and correlate with diabetic control, and GM also seems to play a role[75]. In a human study with early diabetic nephropathy, Barrios *et al*[76] (2015) showed an increase in colonic GM, whereas with end-stage renal disease patients microbes producing urease, uricase, p-cresol and indole-forming enzymes were seen [77]. The proposed mechanisms for progression of kidney disease could be due to GM imbalance, metabolic shifts, immunosuppression, inflammation, as well as accumulation of uremic toxins[78].

Table 2 Selected animal studies showing the effect of various interventions on the gut microbiome and the role of gut microbiota in diabetes mellitus management

Intervention	Organism	Health benefit	Change in microbiome	Ref.
Intermittent fasting	Mice	Protection from diabetic retinopathy by increasing Tauroursodeoxycholate (a neuroprotective bile acid) producing microbes	Increased Firmicutes and decreased Bacteroidetes and Verrucomicrobia in diabetic mice undergoing intermittent fasting	Beli <i>et al</i> [73], 2018
Antibiotic treatment (ampicillin, metronidazole, neomycin, vancomycin, or their cocktail)	Mice	Reduction in fasting glucose. Change in glucose tolerance (seen with ampicillin, vancomycin, or cocktail)	Alterations in the α - and β - diversity. An association with <i>Akkermansia muciniphila</i> with decrease fasting glucose. The effect is mediated through systemic changes in glucose metabolism	Rodrigues <i>et al</i> [94], 2017
Prebiotic: Acorn and sago	Mice	Mice fed acorn and sago derived prebiotics had an amelioration of the glucose intolerance and insulin resistance induced by a high-fat diet feeding. Intake of both novel prebiotics as well as inulin increases SCFAs levels in the mouse gut		Ahmadi <i>et al</i> [103], 2019
Combination of a functional fibre [PolyGlycopleX (PGX) with metformin (MET) or sitagliptin and metformin (S/MET)]	Mice	PGX + MET and PGX + S/MET showed reduced glycemia compared to controls and single treatment ($P = 0.001$). HbA1c was lower in PGX + S/MET compared to all other treatments ($P = 0.001$)		Reimer <i>et al</i> [93], 2014
Artificial sweetener (Neotame)	Mice	Decreased butyrate synthetic genes in Neotame group. Higher concentrations of cholesterol ($P < 0.05$) and fatty acids ($P < 0.05$) in Neotame treated mice feces	Reduction in α -diversity and altered β -diversity. Reduced Firmicutes ($P < 0.01$) and increased Bacteroides ($P < 0.01$)	Chi <i>et al</i> [85], 2018
Combination of metformin and a prebiotic [konjac mannan-oligosaccharides (MOS)]	Mice	Combination of metformin and MOS help ameliorate insulin resistance and improved glycemic control ($P < 0.05$) and repair islet and hepatic histology	Metformin and MOS change the microbiome ($P < 0.0001$) with: Decreased: Rikenellaceae and Clostridiales; Increased: <i>Akkermansia muciniphila</i> and <i>Bifidobacterium pseudolongum</i>	Zheng <i>et al</i> [96], 2018

MANAGEMENT

In DM, normal GM can be restored using diet, gut biotics, faecal transplantation, and bariatric surgery, which may help with the proper management of DM.

Faecal transplant, bariatric surgery

There is some evidence from human studies, that both faecal transplant and bariatric surgery improved the glucose and metabolic parameters by altering the GM [48]. A meta-analysis done by Magouliotis *et al* [79] (2017) showed some discrepancy between the human studies and the benefits witnessed from bariatric surgery. Another study looking at obese insulin resistant subjects who received allogenic faecal transplants from a lean insulin sensitive donor show improved insulin sensitivity for a short period of 6 weeks, however the benefit was not seen past 12 weeks [80] (Table 3).

Nutritional therapy

Diet can modulate the GM and play a role in the management of DM by preventing gut dysbiosis [81] (Table 2). Fruits and vegetables contain polyphenols which can increase beneficial GM like *A. muciniphila*, *Lactobacilli* and *Bifidobacteria* [82]. Unbalanced dietary intake can affect the structure and abundance of GM which can play a role in the development of DM [83].

Artificial sweeteners

Artificial sweeteners are no-calorie sugar substitutes, may induce glucose intolerance by affecting the gut microbes. In an animal study with saccharin-fed mice showed an increase in Bacteroides and a reduction in *Lactobacillus reuteri* leading to GM dysbiosis and glucose intolerance [84]. Similar effects were seen in another study by Chi *et al* [85] (2018) using the artificial sweetener, Neotame. In a cross-sectional human study by Frankenfeld *et al* [86] (2015), showed sweeteners like aspartame or acesulfame-K found no effect on gut bacterial abundance. A recent randomized-blinded crossover study in healthy participants did not demonstrate measurable changes in the GM or in SCFAs after 14 d daily intake of aspartame and sucralose [87]. These preliminary observations

Table 3 Selected human studies showing the effect of diet, gut biotics, faecal transplantation and bariatric surgery on gut microbiome and the role of gut microbiota in diabetes mellitus management

Intervention	Organism	Health benefit	Change in microbiome	Ref.
Probiotics	Human	Decreased fasting blood glucose and HbA1c levels. Increased HDL levels, however no significant effect on BMI and LDL levels were found		Kocsis <i>et al</i> [112], 2020
Artificial sweeteners (aspartame and acesulfame-K)	Human		Compared to controls, aspartame and acesulfame-K had different bacterial diversity ($P < 0.01$, $P = 0.03$ respectively), compared to controls	Frankenfeld <i>et al</i> [86], 2015
Probiotics, Prebiotics, or synbiotics	Human (meta-analysis)	The use of probiotics, prebiotics, or synbiotics showed a decrease in FBG ($P < 0.01$), total cholesterol ($P = 0.02$), triacylglycerols ($P = 0.01$) and insulinaemia ($P < 0.01$), as well as increased HDL-cholesterol levels ($P < 0.01$). Even though HbA1c reduction is seen it is not statistically significant. No effect on LDL-cholesterol was seen		Bock <i>et al</i> [115], 2020
Laparoscopic sleeve gastrectomy	Human	Decreased weight and BMI. Restored insulin tolerance and type 2 DM remission	Increased: <i>Bacteroidetes/Firmicutes</i> ratio at 1- and 3-months post surgery. <i>Lactobacillales</i>	Kikuchi <i>et al</i> [128], 2018; Li <i>et al</i> [129], 2013
Roux-en-Y gastric bypass	Human	Type 2 DM remission and improved BMI and weight loss. Improved gastric emptying and bile acid metabolism	Decreased: <i>Bacteroidetes/Firmicutes</i> ratio. Improved probiotic supplementation effects due to lowered pH environment	Selber-Hnatiw <i>et al</i> [52], 2020; Li <i>et al</i> [129], 2013

BMI: Body mass index; HbA1c: Hemoglobin A1c; FBG: Fasting blood glucose; HDL: High-density lipoprotein; LDL: Low-density lipoprotein.

needed to be established in future human research studies.

ALTERATION OF GM BY ANTIDIABETIC DRUGS AND ITS ROLE IN DM MANAGEMENT

Antidiabetic drugs can influence the gut microbiome by affecting the drug microbiome interface, whereas the gut microbiome also influences the metabolism and play a role in the efficacy of antidiabetic drugs. The interactions of antidiabetic drugs and microbiota is getting more attention as it may play a role in the management of DM [88]. Antidiabetic agents cause alteration of the specific gut microbes. Metformin increases the population of *Akkermansia muciniphila* by 18-fold, enhancing the digestion of mucin and increasing SCFA [89]. Metformin, in addition to *Akkermasia*, causes increase in *Lactobacillus* and *Bifidobacterium*, whereas insulin increase *Fusobacterium* [90]. This first line antidiabetic agent in type 2 DM modifies the GM, alter the bile acid circulation and thereby a possibility that primary site of action may be gut and the GM [91].

Understanding the pharmacokinetics, pharmacodynamics and pharmacomicrobiomics of antidiabetic medications and gut microbes can help to understand drug-gut microbiome and its potential benefit with antidiabetic drugs. Overall, it may help to better manage the DM management [92].

Antidiabetic drugs have been shown to affect the different gut microbes and their metabolic effects through the medication-microbiome-metabolism axis. GM can influence the pharmacokinetics of various antidiabetic drugs such as drug absorption, drug metabolism which can affect the potency of these medications. Overall, there is a bidirectional relationship exist between antidiabetic drugs and gut microbes [88].

Different combinations of antidiabetic drugs are used to better control DM. The commonly used combination is metformin with sulphonylureas, thiazolidinediones, DPP-4 inhibitors and insulin. One animal study showed some delay in the progression of DM when sitagliptin/metformin combination given with a prebiotic fibre [93]. Currently, there is a need for more research of different combination therapies on GM.

GUT BIOTICS AND DM

Animal studies

Several animal studies have showed that gut biotics, like prebiotics and probiotics, can improve the efficacy of antidiabetic drugs. Treatment with individual or a cocktail of antibiotics reduced dysbiosis and decrease fasting glucose but did not affect body weight, as well as antibiotic treatment also changed gene expression in the ileum and liver, and shifted the alpha (α) and beta (β) diversities of GM[94]. In an animal study with mice, combining probiotics and/or prebiotics with antidiabetic medications showed an improvement in glycemic control and insulin sensitivity[95]. A study by Reimer *et al*[93] (2014) found that using a combination of sitagliptin and metformin with a functional fiber can delay DM progression. In an animal study usage of mannan-oligosaccharides by altering the GM increased the hypoglycemic effects of metformin[96]. Yang *et al*[97] (2020) found that Genistein found in soybeans and soy derived foods (prebiotic) helped to improve glucose and lipid metabolism by altering GM composition[97]. In another animal study, certain GM like *Bacteroides fragilis*, *A. muciniphila*, *L. plantarum*, *L. casei* can induce interleukin 10 (IL-10), which has been shown to improve both insulin resistance and glucose metabolism[98] (Table 2).

Human studies

Many gut microbes have been shown to have antidiabetic effect in humans by different mechanisms including effect on insulin sensitivity[99]. *Roseburia intestinalis* can improve insulin sensitivity by increasing IL-22 production[100]. Some strains of *Lactobacilli* act like acarbose and have been shown to inhibit alpha glucosidase[101]. Prebiotics can feed the gut microbiome and increase the population of L-cells in the intestine and thereby increase the amount of GLP-1[102] and prevent high fat diet induced insulin resistance[103]. In the recent PREMOT randomized control trial (RTC) study, probiotics showed antidiabetic effect by altering metabolic homeostasis [104]. Thus, GM may be useful in the management of DM[105]. Jafarnejad *et al*[106] (2015) and Asemi *et al*[107] (2014), in their two studies showed multi-probiotic supplement as well as synbiotic (*L. sporogenes* plus inulin) product helps to reduce glucose and other metabolic parameters. Tonucci *et al*[108] (2015) in their double-blind RCT study comparing fermented milk containing *L. acidophilus* (LA-5) plus *B. animalis* (*Lactis BB-12*) with plain fermented milk in 45 type 2 DM subjects showed decreased in HbA1c as well as low-density lipoproteins cholesterol and inflammatory cytokines. Multiple RTCs and the meta-analysis of these RCT's with different gut microbes demonstrated antidiabetic effect as well as effect on different metabolic parameters [109-111] (Table 3).

A recent meta-analysis of 14 RCTs showed significant decrease in HbA1c in the probiotic group compared to placebo controls, weighted mean difference (WMD) is -0.33%, 95%CI -0.53 to -0.13, $P = 0.001$. In this meta-analysis, probiotics significantly reduced fasting blood glucose, insulin, lipid profile and inflammatory marker in addition to blood pressure levels[112]. Another meta-analysis showed similar result with reduction in HbA1c% (WMD = -0.24, 95%CI: -0.44 to -0.04, $P = 0.02$), fasting blood glucose (WMD = -0.44 mmol/L, 95%CI: -0.74 to -0.15, $P = 0.003$)[113,114]. A meta-analysis study done in 2021 with probiotics, prebiotics or synbiotics on type 2 DM also showed significant improvement in glucose and other metabolic parameters [115]. Prebiotic inulin improves glycemic control in young adults with type 1 DM [116]. Certain specific species of probiotic microbes as well as certain prebiotics by altering the GM was shown to improve the auto-immune condition, which plays a major role in the pathogenesis of type 1 DM[117].

A study by Didari *et al*[118] (2014) looked at the safety of probiotics and synbiotics and found that certain populations, such as patients who are immunocompromised, with cardiac valvular disease, having a central venous catheter, or those with short-bowel syndrome may have an increased risk for systemic infections. Thus, caution may be warranted when using these products in diabetic patients and a risk-benefit analysis should be considered.

GUT MICROBES AND THE PREVENTION OF DM

Some preliminary evidence in animal studies indicates altering GM may help to prevent DM[119,120]. A recent study by Gurung *et al*[30] (2020) showed with certain gut microbes like *Bifidobacterium*, *Bacteroides*, *Faecalibacterium*, *Akkermansia* and *Roseburia* have a negative association with DM and appears to be protective. In 42

healthy adults, GM *Lactobacillus johnsonii* seems to reduce the risk of type 1 DM[121].

CONCLUSION

Gut dysbiosis plays a role in the development and progression of DM. The current evidence also points out that the GM can play a role in DM related complications. Modulation of the gut bacteria or dysbiosis can be corrected by fibre, diet, antidiabetic medications, and by using gut biotics like prebiotics, probiotics, and synbiotics as well as by bariatric surgery and faecal transplantation. The interaction between gut microbes and antidiabetic agents is a promising field that may change the landscape of DM management in the future. There is some preliminary evidence to show that GM may play a role in the prevention of DM. More research is needed on a large scale to confirm these findings.

REFERENCES

- 1 **Toniolo A**, Cassani G, Puggioni A, Rossi A, Colombo A, Onodera T, Ferrannini E. The diabetes pandemic and associated infections: suggestions for clinical microbiology. *Rev Med Microbiol* 2019; **30**: 1-17 [PMID: 30662163 DOI: 10.1097/MRM.0000000000000155]
- 2 **NCD Risk Factor Collaboration (NCD-RisC)**. Worldwide trends in diabetes since 1980: a pooled analysis of 751 population-based studies with 4.4 million participants. *Lancet* 2016; **387**: 1513-1530 [PMID: 27061677 DOI: 10.1016/S0140-6736(16)00618-8]
- 3 **Ayadurai S**, Hattigh HL, Tee LB, Md Said SN. A Narrative Review of Diabetes Intervention Studies to Explore Diabetes Care Opportunities for Pharmacists. *J Diabetes Res* 2016; **2016**: 5897452 [PMID: 27247949 DOI: 10.1155/2016/5897452]
- 4 **American Diabetes Association**. 2. Classification and Diagnosis of Diabetes: *Standards of Medical Care in Diabetes-2018*. *Diabetes Care* 2018; **41**: S13-S27 [PMID: 29222373 DOI: 10.2337/dc18-S002]
- 5 **Bullard KM**, Cowie CC, Lessem SE, Saydah SH, Menke A, Geiss LS, Orchard TJ, Rolka DB, Imperatore G. Prevalence of Diagnosed Diabetes in Adults by Diabetes Type - United States, 2016. *MMWR Morb Mortal Wkly Rep* 2018; **67**: 359-361 [PMID: 29596402 DOI: 10.15585/mmwr.mm6712a2]
- 6 **Human Microbiome Project Consortium**. Structure, function and diversity of the healthy human microbiome. *Nature* 2012; **486**: 207-214 [PMID: 22699609 DOI: 10.1038/nature11234]
- 7 **Chattopadhyay A**, Mythili S. The journey of gut microbiome – an introduction and its influence on metabolic disorders. *Front Biol* 2018; **5**: 327-341 [DOI: 10.1007/s11515-018-1490-6]
- 8 **Molina DK**, DiMaio VJ. Normal organ weights in men: part I-the heart. *Am J Forensic Med Pathol* 2012; **33**: 362-367 [PMID: 22182983 DOI: 10.1097/PAF.0b013e31823d298b]
- 9 **Qin J**, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, Nielsen T, Pons N, Levenez F, Yamada T, Mende DR, Li J, Xu J, Li S, Li D, Cao J, Wang B, Liang H, Zheng H, Xie Y, Tap J, Lepage P, Bertalan M, Batto JM, Hansen T, Le Paslier D, Linneberg A, Nielsen HB, Pelletier E, Renault P, Sicheritz-Ponten T, Turner K, Zhu H, Yu C, Jian M, Zhou Y, Li Y, Zhang X, Qin N, Yang H, Wang J, Brunak S, Doré J, Guarner F, Kristiansen K, Pedersen O, Parkhill J, Weissenbach J, MetaHIT Consortium, Bork P, Ehrlich SD, Wang J. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* 2010; **464**: 59-65 [PMID: 20203603 DOI: 10.1038/nature08821]
- 10 **Conlon MA**, Bird AR. The impact of diet and lifestyle on gut microbiota and human health. *Nutrients* 2014; **7**: 17-44 [PMID: 25545101 DOI: 10.3390/nu7010017]
- 11 **He Y**, Wu W, Zheng HM, Li P, McDonald D, Sheng HF, Chen MX, Chen ZH, Ji GY, Zheng ZD, Mujagond P, Chen XJ, Rong ZH, Chen P, Lyu LY, Wang X, Wu CB, Yu N, Xu YJ, Yin J, Raes J, Knight R, Ma WJ, Zhou HW. Regional variation limits applications of healthy gut microbiome reference ranges and disease models. *Nat Med* 2018; **24**: 1532-1535 [PMID: 30150716 DOI: 10.1038/s41591-018-0164-x]
- 12 **Maier L**, Pruteanu M, Kuhn M, Zeller G, Telzerow A, Anderson EE, Brochado AR, Fernandez KC, Dose H, Mori H, Patil KR, Bork P, Typas A. Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature* 2018; **555**: 623-628 [PMID: 29555994 DOI: 10.1038/nature25979]
- 13 **Schroeder BO**, Bäckhed F. Signals from the gut microbiota to distant organs in physiology and disease. *Nat Med* 2016; **22**: 1079-1089 [PMID: 27711063 DOI: 10.1038/nm.4185]
- 14 **Barlow GM**, Yu A, Mathur R. Role of the Gut Microbiome in Obesity and Diabetes Mellitus. *Nutr Clin Pract* 2015; **30**: 787-797 [PMID: 26452391 DOI: 10.1177/0884533615609896]
- 15 **Bouter KE**, van Raalte DH, Groen AK, Nieuwdorp M. Role of the Gut Microbiome in the Pathogenesis of Obesity and Obesity-Related Metabolic Dysfunction. *Gastroenterology* 2017; **152**: 1671-1678 [PMID: 28192102 DOI: 10.1053/j.gastro.2016.12.048]
- 16 **Chen X**, Devaraj S. Gut Microbiome in Obesity, Metabolic Syndrome, and Diabetes. *Curr Diab Rep* 2018; **18**: 129 [PMID: 30338410 DOI: 10.1007/s11892-018-1104-3]

- 17 **Vallianou NG**, Stratigou T, Tsarakis S. Microbiome and diabetes: Where are we now? *Diabetes Res Clin Pract* 2018; **146**: 111-118 [PMID: [30342053](#) DOI: [10.1016/j.diabres.2018.10.008](#)]
- 18 **Purkait D**, Hameed S, Fatima Z. Chapter 15 - gut microbiome: Current development, challenges, and perspectives. In: Rastegari AA, Yadav AN, Yadav N. New and future developments in microbial biotechnology and bioengineering: Elsevier, 2020: 227-241
- 19 **Larsen N**, Vogensen FK, van den Berg FW, Nielsen DS, Andreasen AS, Pedersen BK, Al-Soud WA, Sørensen SJ, Hansen LH, Jakobsen M. Gut microbiota in human adults with type 2 diabetes differs from non-diabetic adults. *PLoS One* 2010; **5**: e9085 [PMID: [20140211](#) DOI: [10.1371/journal.pone.0009085](#)]
- 20 **Thingholm LB**, Rühlemann MC, Koch M, Fuqua B, Laucke G, Boehm R, Bang C, Franzosa EA, Hübenthal M, Rahnard A, Frost F, Lloyd-Price J, Schirmer M, Lusis AJ, Vulpe CD, Lerch MM, Homuth G, Kacprowski T, Schmidt CO, Nöthlings U, Karlsen TH, Lieb W, Laudes M, Franke A, Huttenhower C. Obese Individuals with and without Type 2 Diabetes Show Different Gut Microbial Functional Capacity and Composition. *Cell Host Microbe* 2019; **26**: 252-264.e10 [PMID: [31399369](#) DOI: [10.1016/j.chom.2019.07.004](#)]
- 21 **Nowak K**, Jabłońska E, Ratajczak-Wrona W. Immunomodulatory effects of synthetic endocrine disrupting chemicals on the development and functions of human immune cells. *Environ Int* 2019; **125**: 350-364 [PMID: [30743143](#) DOI: [10.1016/j.envint.2019.01.078](#)]
- 22 **Mrozinska S**, Kapusta P, Gosiewski T, Sroka-Oleksiak A, Ludwig-Słomczyńska AH, Matejko B, Kiec-Wilk B, Bulanda M, Malecki MT, Wolkow PP, Klupa T. The Gut Microbiota Profile According to Glycemic Control in Type 1 Diabetes Patients Treated with Personal Insulin Pumps. *Microorganisms* 2021; **9** [PMID: [33445500](#) DOI: [10.3390/microorganisms9010155](#)]
- 23 **Huang Y**, Li SC, Hu J, Ruan HB, Guo HM, Zhang HH, Wang X, Pei YF, Pan Y, Fang C. Gut microbiota profiling in Han Chinese with type 1 diabetes. *Diabetes Res Clin Pract* 2018; **141**: 256-263 [PMID: [29733871](#) DOI: [10.1016/j.diabres.2018.04.032](#)]
- 24 **Fassatoui M**, Lopez-Siles M, Diaz-Rizzolo DA, Jmel H, Naouali C, Abdesslem G, Chikhaoui A, Nadal B, Jamoussi H, Abid A, Gomis R, Abdelhak S, Martinez-Medina M, Kefi R. Gut microbiota imbalances in Tunisian participants with type 1 and type 2 diabetes mellitus. *Biosci Rep* 2019; **39** [PMID: [31147456](#) DOI: [10.1042/BSR20182348](#)]
- 25 **Elena RM**, Gabriela GD, Arnulfo GC, Enrique CA. Studying the Gut Microbiome of Latin America and Hispanic/Latino Populations. Insight into Obesity and Diabetes: Systematic Review. *Curr Diabetes Rev* 2019; **15**: 294-301 [PMID: [30058496](#) DOI: [10.2174/1573399814666180730124817](#)]
- 26 **Walker AW**, Ince J, Duncan SH, Webster LM, Holtrop G, Ze X, Brown D, Stares MD, Scott P, Bergerat A, Louis P, McIntosh F, Johnstone AM, Lobley GE, Parkhill J, Flint HJ. Dominant and diet-responsive groups of bacteria within the human colonic microbiota. *ISME J* 2011; **5**: 220-230 [PMID: [20686513](#) DOI: [10.1038/ismej.2010.118](#)]
- 27 **Blandino G**, Inturri R, Lazzara F, Di Rosa M, Malaguarnera L. Impact of gut microbiota on diabetes mellitus. *Diabetes Metab* 2016; **42**: 303-315 [PMID: [27179626](#) DOI: [10.1016/j.diabet.2016.04.004](#)]
- 28 **Chen PC**, Chien YW, Yang SC. The alteration of gut microbiota in newly diagnosed type 2 diabetic patients. *Nutrition* 2019; **63-64**: 51-56 [PMID: [30933725](#) DOI: [10.1016/j.nut.2018.11.019](#)]
- 29 **Das T**, Jayasudha R, Chakravarthy S, Prashanthi GS, Bhargava A, Tyagi M, Rani PK, Pappuru RR, Sharma S, Shivaji S. Alterations in the gut bacterial microbiome in people with type 2 diabetes mellitus and diabetic retinopathy. *Sci Rep* 2021; **11**: 2738 [PMID: [33531650](#) DOI: [10.1038/s41598-021-82538-0](#)]
- 30 **Gurung M**, Li Z, You H, Rodrigues R, Jump DB, Morgun A, Shulzhenko N. Role of gut microbiota in type 2 diabetes pathophysiology. *EBioMedicine* 2020; **51**: 102590 [PMID: [31901868](#) DOI: [10.1016/j.ebiom.2019.11.051](#)]
- 31 **Sedighi M**, Razavi S, Navab-Moghadam F, Khamseh ME, Alaei-Shahmiri F, Mehrtash A, Amirmozafari N. Comparison of gut microbiota in adult patients with type 2 diabetes and healthy individuals. *Microb Pathog* 2017; **111**: 362-369 [PMID: [28912092](#) DOI: [10.1016/j.micpath.2017.08.038](#)]
- 32 **Nuli R**, Cai J, Kadeer A, Zhang Y, Mohemaiti P. Integrative Analysis Toward Different Glucose Tolerance-Related Gut Microbiota and Diet. *Front Endocrinol (Lausanne)* 2019; **10**: 295 [PMID: [31191448](#) DOI: [10.3389/fendo.2019.00295](#)]
- 33 **Canli PD**, Bibiloni R, Knauf C, Waget A, Neyrinck AM, Delzenne NM, Burcelin R. Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes* 2008; **57**: 1470-1481 [PMID: [18305141](#) DOI: [10.2337/db07-1403](#)]
- 34 **Païssé S**, Valle C, Servant F, Courtney M, Burcelin R, Amar J, Lelouvier B. Comprehensive description of blood microbiome from healthy donors assessed by 16S targeted metagenomic sequencing. *Transfusion* 2016; **56**: 1138-1147 [PMID: [26865079](#) DOI: [10.1111/trf.13477](#)]
- 35 **Olde Loohuis LM**, Mangul S, Ori APS, Jospin G, Koslicki D, Yang HT, Wu T, Boks MP, Lomen-Hoerth C, Wiedau-Pazos M, Cantor RM, de Vos WM, Kahn RS, Eskin E, Ophoff RA. Transcriptome analysis in whole blood reveals increased microbial diversity in schizophrenia. *Transl Psychiatry* 2018; **8**: 96 [PMID: [29743478](#) DOI: [10.1038/s41398-018-0107-9](#)]
- 36 **Whittle E**, Leonard MO, Harrison R, Gant TW, Tonge DP. Multi-Method Characterization of the Human Circulating Microbiome. *Front Microbiol* 2018; **9**: 3266 [PMID: [30705670](#) DOI: [10.3389/fmicb.2018.03266](#)]
- 37 **Sato J**, Kanazawa A, Ikeda F, Yoshihara T, Goto H, Abe H, Komiya K, Kawaguchi M, Shimizu T, Ogihara T, Tamura Y, Sakurai Y, Yamamoto R, Mita T, Fujitani Y, Fukuda H, Nomoto K,

- Takahashi T, Asahara T, Hirose T, Nagata S, Yamashiro Y, Watada H. Gut dysbiosis and detection of "live gut bacteria" in blood of Japanese patients with type 2 diabetes. *Diabetes Care* 2014; **37**: 2343-2350 [PMID: [24824547](#) DOI: [10.2337/dc13-2817](#)]
- 38 **Qiu J**, Zhou H, Jing Y, Dong C. Association between blood microbiome and type 2 diabetes mellitus: A nested case-control study. *J Clin Lab Anal* 2019; **33**: e22842 [PMID: [30714640](#) DOI: [10.1002/jcla.22842](#)]
- 39 **Evans JM**, Morris LS, Marchesi JR. The gut microbiome: the role of a virtual organ in the endocrinology of the host. *J Endocrinol* 2013; **218**: R37-R47 [PMID: [23833275](#) DOI: [10.1530/JOE-13-0131](#)]
- 40 **Lyte M**. Probiotics function mechanistically as delivery vehicles for neuroactive compounds: Microbial endocrinology in the design and use of probiotics. *Bioessays* 2011; **33**: 574-581 [PMID: [21732396](#) DOI: [10.1002/bies.201100024](#)]
- 41 **Hampel R**, Stárka L. Endocrine disruptors and gut microbiome interactions. *Physiol Res* 2020; **69**: S211-S223 [PMID: [33094620](#) DOI: [10.33549/physiolres.934513](#)]
- 42 **Clarke G**, Stilling RM, Kennedy PJ, Stanton C, Cryan JF, Dinan TG. Minireview: Gut microbiota: the neglected endocrine organ. *Mol Endocrinol* 2014; **28**: 1221-1238 [PMID: [24892638](#) DOI: [10.1210/me.2014-1108](#)]
- 43 **Rosenfeld CS**. Gut Dysbiosis in Animals Due to Environmental Chemical Exposures. *Front Cell Infect Microbiol* 2017; **7**: 396 [PMID: [28936425](#) DOI: [10.3389/fcimb.2017.00396](#)]
- 44 **Velmurugan G**, Ramprasad T, Gilles M, Swaminathan K, Ramasamy S. Gut Microbiota, Endocrine-Disrupting Chemicals, and the Diabetes Epidemic. *Trends Endocrinol Metab* 2017; **28**: 612-625 [PMID: [28571659](#) DOI: [10.1016/j.tem.2017.05.001](#)]
- 45 **Needell JC**, Zipris D. The Role of the Intestinal Microbiome in Type 1 Diabetes Pathogenesis. *Curr Diab Rep* 2016; **16**: 89 [PMID: [27523648](#) DOI: [10.1007/s11892-016-0781-z](#)]
- 46 **Singer-Englar T**, Barlow G, Mathur R. Obesity, diabetes, and the gut microbiome: an updated review. *Expert Rev Gastroenterol Hepatol* 2019; **13**: 3-15 [PMID: [30791839](#) DOI: [10.1080/17474124.2019.1543023](#)]
- 47 **Yu F**, Han W, Zhan G, Li S, Jiang X, Wang L, Xiang S, Zhu B, Yang L, Luo A, Hua F, Yang C. Abnormal gut microbiota composition contributes to the development of type 2 diabetes mellitus in db/db mice. *Aging (Albany NY)* 2019; **11**: 10454-10467 [PMID: [31760385](#) DOI: [10.18632/aging.102469](#)]
- 48 **Halmos T**, Suba I. [Physiological patterns of intestinal microbiota. The role of dysbacteriosis in obesity, insulin resistance, diabetes and metabolic syndrome]. *Orv Hetil* 2016; **157**: 13-22 [PMID: [26708682](#) DOI: [10.1556/650.2015.30296](#)]
- 49 **Fallucca F**, Porrata C, Fallucca S, Pianesi M. Influence of diet on gut microbiota, inflammation and type 2 diabetes mellitus. First experience with macrobiotic Ma-Pi 2 diet. *Diabetes Metab Res Rev* 2014; **30** Suppl 1: 48-54 [PMID: [24532292](#) DOI: [10.1002/dmrr.2518](#)]
- 50 **Aw W**, Fukuda S. Understanding the role of the gut ecosystem in diabetes mellitus. *J Diabetes Investig* 2018; **9**: 5-12 [PMID: [28390093](#) DOI: [10.1111/jdi.12673](#)]
- 51 **Acosta-Montañó P**, Rodríguez-Velázquez E, Ibarra-López E, Frayde-Gómez H, Mas-Oliva J, Delgado-Coello B, Rivero IA, Alatorre-Meda M, Aguilera J, Guevara-Olaya L, García-González V. Fatty Acid and Lipopolysaccharide Effect on Beta Cells Proteostasis and its Impact on Insulin Secretion. *Cells* 2019; **8** [PMID: [31412623](#) DOI: [10.3390/cells8080884](#)]
- 52 **Selber-Hnatiw S**, Sultana T, Tse W, Abdollahi N, Abdullah S, Al Rahbani J, Alazar D, Alrumhein NJ, Aprikian S, Arshad R, Azuelos JD, Bernadotte D, Beswick N, Chazbey H, Church K, Ciubotaru E, D'Amato L, Del Corpo T, Deng J, Di Giulio BL, Diveeva D, Elahie E, Frank JGM, Furze E, Garner R, Gibbs V, Goldberg-Hall R, Goldman CJ, Goltsios FF, Gorjipour K, Grant T, Greco B, Guliyev N, Habrich A, Hyland H, Ibrahim N, Iozzo T, Jawaheer-Fenaoui A, Jaworski JJ, Jhaji MK, Jones J, Joyette R, Kaudeer S, Kelley S, Kiani S, Koayes M, Kpata AJAL, Maingot S, Martin S, Mathers K, McCulloch S, McNamara K, Mendonca J, Mohammad K, Momtaz SA, Navaratnarajah T, Nguyen-Duong K, Omran M, Ortiz A, Patel A, Paul-Cole K, Plaisir PA, Porras Marroquin JA, Prevost A, Quach A, Rafal AJ, Ramsarun R, Rhnima S, Rili L, Safir N, Samson E, Sandiford RR, Secondi S, Shahid S, Shahrooz M, Sidibé F, Smith M, Sreng Flores AM, Suarez Ybarra A, Sénéchal R, Taifour T, Tang L, Trapid A, Tremblay Potvin M, Wainberg J, Wang DN, Weissenberg M, White A, Wilkinson G, Williams B, Wilson JR, Zoppi J, Zouboulakis K, Gamberi C. Metabolic networks of the human gut microbiota. *Microbiology (Reading)* 2020; **166**: 96-119 [PMID: [31799915](#) DOI: [10.1099/mic.0.000853](#)]
- 53 **Gordon JI**, Hooper LV, McNevin MS, Wong M, Bry L. Epithelial cell growth and differentiation. III. Promoting diversity in the intestine: conversations between the microflora, epithelium, and diffuse GALT. *Am J Physiol* 1997; **273**: G565-G570 [PMID: [9316460](#) DOI: [10.1152/ajpgi.1997.273.3.G565](#)]
- 54 **Karlsson F**, Tremaroli V, Nielsen J, Bäckhed F. Assessing the human gut microbiota in metabolic diseases. *Diabetes* 2013; **62**: 3341-3349 [PMID: [24065795](#) DOI: [10.2337/db13-0844](#)]
- 55 **Tremaroli V**, Bäckhed F. Functional interactions between the gut microbiota and host metabolism. *Nature* 2012; **489**: 242-249 [PMID: [22972297](#) DOI: [10.1038/nature11552](#)]
- 56 **Ussar S**, Griffin NW, Bezy O, Fujisaka S, Vienberg S, Softic S, Deng L, Bry L, Gordon JI, Kahn CR. Interactions between Gut Microbiota, Host Genetics and Diet Modulate the Predisposition to Obesity and Metabolic Syndrome. *Cell Metab* 2015; **22**: 516-530 [PMID: [26299453](#) DOI: [10.1016/j.cmet.2015.07.007](#)]

- 57 **El Aidy S**, Hooiveld G, Tremaroli V, Bäckhed F, Kleerebezem M. The gut microbiota and mucosal homeostasis: colonized at birth or at adulthood, does it matter? *Gut Microbes* 2013; **4**: 118-124 [PMID: [23333858](#) DOI: [10.4161/gmic.23362](#)]
- 58 **den Besten G**, van Eunen K, Groen AK, Venema K, Reijngoud DJ, Bakker BM. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *J Lipid Res* 2013; **54**: 2325-2340 [PMID: [23821742](#) DOI: [10.1194/jlr.R036012](#)]
- 59 **Canfora EE**, Jocken JW, Blaak EE. Short-chain fatty acids in control of body weight and insulin sensitivity. *Nat Rev Endocrinol* 2015; **11**: 577-591 [PMID: [26260141](#) DOI: [10.1038/nrendo.2015.128](#)]
- 60 **Lin HV**, Frassetto A, Kowalik EJ Jr, Nawrocki AR, Lu MM, Kosinski JR, Hubert JA, Szeto D, Yao X, Forrest G, Marsh DJ. Butyrate and propionate protect against diet-induced obesity and regulate gut hormones *via* free fatty acid receptor 3-independent mechanisms. *PLoS One* 2012; **7**: e35240 [PMID: [22506074](#) DOI: [10.1371/journal.pone.0035240](#)]
- 61 **Gao Z**, Yin J, Zhang J, Ward RE, Martin RJ, Lefevre M, Cefalu WT, Ye J. Butyrate improves insulin sensitivity and increases energy expenditure in mice. *Diabetes* 2009; **58**: 1509-1517 [PMID: [19366864](#) DOI: [10.2337/db08-1637](#)]
- 62 **Rácz B**, Dušková M, Stárka L, Hainer V, Kunešová M. Links between the circadian rhythm, obesity and the microbiome. *Physiol Res* 2018; **67**: S409-S420 [PMID: [30484668](#) DOI: [10.33549/physiolres.934020](#)]
- 63 **Koliada A**, Syzenko G, Moseiko V, Budovska L, Puchkov K, Perederiy V, Gavalko Y, Dorofeyev A, Romanenko M, Tkach S, Sineok L, Lushchak O, Vaiserman A. Association between body mass index and Firmicutes/Bacteroidetes ratio in an adult Ukrainian population. *BMC Microbiol* 2017; **17**: 120 [PMID: [28532414](#) DOI: [10.1186/s12866-017-1027-1](#)]
- 64 **Hung WC**, Hung WW, Tsai HJ, Chang CC, Chiu YW, Hwang SJ, Kuo MC, Chen SC, Dai CY, Tsai YC. The Association of Targeted Gut Microbiota with Body Composition in Type 2 Diabetes Mellitus. *Int J Med Sci* 2021; **18**: 511-519 [PMID: [33390820](#) DOI: [10.7150/ijms.51164](#)]
- 65 **Mejnikman AS**, Gerdes VE, Nieuwdorp M, Herrema H. Evaluating Causality of Gut Microbiota in Obesity and Diabetes in Humans. *Endocr Rev* 2018; **39**: 133-153 [PMID: [29309555](#) DOI: [10.1210/er.2017-00192](#)]
- 66 **Leustean AM**, Ciocoiu M, Sava A, Costea CF, Floria M, Tarniceriu CC, Tanase DM. Implications of the Intestinal Microbiota in Diagnosing the Progression of Diabetes and the Presence of Cardiovascular Complications. *J Diabetes Res* 2018; **2018**: 5205126 [PMID: [30539026](#) DOI: [10.1155/2018/5205126](#)]
- 67 **Zhao S**, Jang S, Lee YK, Kim DG, Jin Z, Koh HJ. Genetic Basis of Tiller Dynamics of Rice Revealed by Genome-Wide Association Studies. *Plants (Basel)* 2020; **9** [PMID: [33276582](#) DOI: [10.3390/nu12123719](#)]
- 68 **Preguiça I**, Alves A, Nunes S, Gomes P, Fernandes R, Viana SD, Reis F. Diet-Induced Rodent Models of Diabetic Peripheral Neuropathy, Retinopathy and Nephropathy. *Nutrients* 2020; **12** [PMID: [31963709](#) DOI: [10.3390/nu12010250](#)]
- 69 **Ham B**, Hwang HB, Jung SH, Chang S, Kang KD, Kwon MJ. Distribution and Diversity of Ocular Microbial Communities in Diabetic Patients Compared with Healthy Subjects. *Curr Eye Res* 2018; **43**: 314-324 [PMID: [29172724](#) DOI: [10.1080/02713683.2017.1406528](#)]
- 70 **Rowan S**, Taylor A. The Role of Microbiota in Retinal Disease. *Adv Exp Med Biol* 2018; **1074**: 429-435 [PMID: [29721973](#) DOI: [10.1007/978-3-319-75402-4_53](#)]
- 71 **Baim AD**, Movahedan A, Farooq AV, Skondra D. The microbiome and ophthalmic disease. *Exp Biol Med (Maywood)* 2019; **244**: 419-429 [PMID: [30463439](#) DOI: [10.1177/1535370218813616](#)]
- 72 **Floyd JL**, Grant MB. The Gut-Eye Axis: Lessons Learned from Murine Models. *Ophthalmol Ther* 2020; **9**: 499-513 [PMID: [32617914](#) DOI: [10.1007/s40123-020-00278-2](#)]
- 73 **Beli E**, Yan Y, Moldovan L, Vieira CP, Gao R, Duan Y, Prasad R, Bhatwadekar A, White FA, Townsend SD, Chan L, Ryan CN, Morton D, Moldovan EG, Chu FI, Oudit GY, Derendorf H, Adorini L, Wang XX, Evans-Molina C, Mirmira RG, Boulton ME, Yoder MC, Li Q, Levi M, Busik JV, Grant MB. Restructuring of the Gut Microbiome by Intermittent Fasting Prevents Retinopathy and Prolongs Survival in *db/db* Mice. *Diabetes* 2018; **67**: 1867-1879 [PMID: [29712667](#) DOI: [10.2337/db18-0158](#)]
- 74 **St Leger AJ**, Caspi RR. Visions of Eye Commensals: The Known and the Unknown About How the Microbiome Affects Eye Disease. *Bioessays* 2018; **40**: e1800046 [PMID: [30289987](#) DOI: [10.1002/bies.201800046](#)]
- 75 **Grasset E**, Burcelin R. The gut microbiota to the brain axis in the metabolic control. *Rev Endocr Metab Disord* 2019; **20**: 427-438 [PMID: [31656993](#) DOI: [10.1007/s11154-019-09511-1](#)]
- 76 **Barrios C**, Beaumont M, Pallister T, Villar J, Goodrich JK, Clark A, Pascual J, Ley RE, Spector TD, Bell JT, Menni C. Gut-Microbiota-Metabolite Axis in Early Renal Function Decline. *PLoS One* 2015; **10**: e0134311 [PMID: [26241311](#) DOI: [10.1371/journal.pone.0134311](#)]
- 77 **Wong J**, Piceno YM, DeSantis TZ, Pahl M, Andersen GL, Vaziri ND. Expansion of urease- and uricase-containing, indole- and p-cresol-forming and contraction of short-chain fatty acid-producing intestinal microbiota in ESRD. *Am J Nephrol* 2014; **39**: 230-237 [PMID: [24643131](#) DOI: [10.1159/000360010](#)]
- 78 **Stavropoulou E**, Kantartzi K, Tsigalou C, Konstantinidis T, Romanidou G, Voudarou C, Bezirtzoglou E. Focus on the Gut-Kidney Axis in Health and Disease. *Front Med (Lausanne)* 2020; **7**: 620102 [PMID: [33553216](#) DOI: [10.3389/fmed.2020.620102](#)]

- 79 **Magouliotis DE**, Tasiopoulou VS, Sioka E, Chatedaki C, Zacharoulis D. Impact of Bariatric Surgery on Metabolic and Gut Microbiota Profile: a Systematic Review and Meta-analysis. *Obes Surg* 2017; **27**: 1345-1357 [PMID: [28265960](#) DOI: [10.1007/s11695-017-2595-8](#)]
- 80 **Vrieze A**, Van Nood E, Holleman F, Salojärvi J, Kootte RS, Bartelsman JF, Dallinga-Thie GM, Ackermans MT, Serlie MJ, Oozeer R, Derrien M, Druesne A, Van Hylckama Vlieg JE, Bloks VW, Groen AK, Heilig HG, Zoetendal EG, Stroes ES, de Vos WM, Hoekstra JB, Nieuwdorp M. Transfer of intestinal microbiota from lean donors increases insulin sensitivity in individuals with metabolic syndrome. *Gastroenterology* 2012; **143**: 913-6.e7 [PMID: [22728514](#) DOI: [10.1053/j.gastro.2012.06.031](#)]
- 81 **Candela M**, Biagi E, Soverini M, Consolandi C, Quercia S, Severgnini M, Peano C, Turroni S, Rampelli S, Pozzilli P, Pianesi M, Fallucca F, Brigidi P. Modulation of gut microbiota dysbioses in type 2 diabetic patients by macrobiotic Ma-Pi 2 diet. *Br J Nutr* 2016; **116**: 80-93 [PMID: [27151248](#) DOI: [10.1017/S0007114516001045](#)]
- 82 **Quddus MR**, Lawrence WD. A Seminal Observation! *Int J Surg Pathol* 2016; **24**: 47 [PMID: [26338717](#) DOI: [10.1177/1066896915604025](#)]
- 83 **Krezalek MA**, Yeh A, Alverdy JC, Morowitz M. Influence of nutrition therapy on the intestinal microbiome. *Curr Opin Clin Nutr Metab Care* 2017; **20**: 131-137 [PMID: [27997410](#) DOI: [10.1097/MCO.0000000000000348](#)]
- 84 **Suez J**, Korem T, Zeevi D, Zilberman-Schapira G, Thaiss CA, Maza O, Israeli D, Zmora N, Gilad S, Weinberger A, Kuperman Y, Harmelin A, Kolodkin-Gal I, Shapiro H, Halpern Z, Segal E, Elinav E. Artificial sweeteners induce glucose intolerance by altering the gut microbiota. *Nature* 2014; **514**: 181-186 [PMID: [25231862](#) DOI: [10.1038/nature13793](#)]
- 85 **Chi L**, Bian X, Gao B, Tu P, Lai Y, Ru H, Lu K. Effects of the Artificial Sweetener Neotame on the Gut Microbiome and Fecal Metabolites in Mice. *Molecules* 2018; **23** [PMID: [29425148](#) DOI: [10.3390/molecules23020367](#)]
- 86 **Frankenfeld CL**, Sikaroodi M, Lamb E, Shoemaker S, Gillevet PM. High-intensity sweetener consumption and gut microbiome content and predicted gene function in a cross-sectional study of adults in the United States. *Ann Epidemiol* 2015; **25**: 736-42.e4 [PMID: [26272781](#) DOI: [10.1016/j.annepidem.2015.06.083](#)]
- 87 **Ahmad SY**, Friel J, Mackay D. The Effects of Non-Nutritive Artificial Sweeteners, Aspartame and Sucralose, on the Gut Microbiome in Healthy Adults: Secondary Outcomes of a Randomized Double-Blinded Crossover Clinical Trial. *Nutrients* 2020; **12** [PMID: [33171964](#) DOI: [10.3390/nu12113408](#)]
- 88 **Whang A**, Nagpal R, Yadav H. Bi-directional drug-microbiome interactions of anti-diabetics. *EBioMedicine* 2019; **39**: 591-602 [PMID: [30553752](#) DOI: [10.1016/j.ebiom.2018.11.046](#)]
- 89 **Wu H**, Esteve E, Tremaroli V, Khan MT, Caesar R, Mannerås-Holm L, Ståhlman M, Olsson LM, Serino M, Planas-Félix M, Xifra G, Mercader JM, Torrents D, Burcelin R, Ricart W, Perkins R, Fernández-Real JM, Bäckhed F. Metformin alters the gut microbiome of individuals with treatment-naïve type 2 diabetes, contributing to the therapeutic effects of the drug. *Nat Med* 2017; **23**: 850-858 [PMID: [28530702](#) DOI: [10.1038/nm.4345](#)]
- 90 **Zhang F**, Wang M, Yang J, Xu Q, Liang C, Chen B, Zhang J, Yang Y, Wang H, Shang Y, Wang Y, Mu X, Zhu D, Zhang C, Yao M, Zhang L. Response of gut microbiota in type 2 diabetes to hypoglycemic agents. *Endocrine* 2019; **66**: 485-493 [PMID: [31410749](#) DOI: [10.1007/s12020-019-02041-5](#)]
- 91 **Buse JB**, DeFronzo RA, Rosenstock J, Kim T, Burns C, Skare S, Baron A, Fineman M. The Primary Glucose-Lowering Effect of Metformin Resides in the Gut, Not the Circulation: Results From Short-term Pharmacokinetic and 12-Week Dose-Ranging Studies. *Diabetes Care* 2016; **39**: 198-205 [PMID: [26285584](#) DOI: [10.2337/dc15-0488](#)]
- 92 **Rizkallah MR**, Saad R, Aziz RK. The human microbiome project, personalized medicine and the birth of pharmacomicrobiomics. *Curr Pharmacogenomics Person Med* 2010; **8**: 182-193 [DOI: [10.2174/187569210792246326](#)]
- 93 **Reimer RA**, Grover GJ, Koetzner L, Gahler RJ, Lyon MR, Wood S. Combining sitagliptin/metformin with a functional fiber delays diabetes progression in Zucker rats. *J Endocrinol* 2014; **220**: 361-373 [PMID: [24389593](#) DOI: [10.1530/JOE-13-0484](#)]
- 94 **Rodrigues RR**, Greer RL, Dong X, DSouza KN, Gurung M, Wu JY, Morgun A, Shulzhenko N. Antibiotic-Induced Alterations in Gut Microbiota Are Associated with Changes in Glucose Metabolism in Healthy Mice. *Front Microbiol* 2017; **8**: 2306 [PMID: [29213261](#) DOI: [10.3389/fmicb.2017.02306](#)]
- 95 **Stenman LK**, Waget A, Garret C, Briand F, Burcelin R, Sulpice T, Lahtinen S. Probiotic B420 and prebiotic polydextrose improve efficacy of antidiabetic drugs in mice. *Diabetol Metab Syndr* 2015; **7**: 75 [PMID: [26366205](#) DOI: [10.1186/s13098-015-0075-7](#)]
- 96 **Zheng J**, Li H, Zhang X, Jiang M, Luo C, Lu Z, Xu Z, Shi J. Prebiotic Mannan-Oligosaccharides Augment the Hypoglycemic Effects of Metformin in Correlation with Modulating Gut Microbiota. *J Agric Food Chem* 2018; **66**: 5821-5831 [PMID: [29701959](#) DOI: [10.1021/acs.jafc.8b00829](#)]
- 97 **Yang R**, Jia Q, Mehmood S, Ma S, Liu X. Genistein ameliorates inflammation and insulin resistance through mediation of gut microbiota composition in type 2 diabetic mice. *Eur J Nutr* 2021; **60**: 2155-2168 [PMID: [33068158](#) DOI: [10.1007/s00394-020-02403-0](#)]
- 98 **Dagdeviren S**, Jung DY, Friedline RH, Noh HL, Kim JH, Patel PR, Tsitsilianos N, Inashima K, Tran DA, Hu X, Loubato MM, Craige SM, Kwon JY, Lee KW, Kim JK. IL-10 prevents aging-

- associated inflammation and insulin resistance in skeletal muscle. *FASEB J* 2017; **31**: 701-710 [PMID: 27811060 DOI: 10.1096/fj.201600832R]
- 99 **Andreassen AS**, Larsen N, Pedersen-Skovsgaard T, Berg RM, Møller K, Svendsen KD, Jakobsen M, Pedersen BK. Effects of *Lactobacillus acidophilus* NCFM on insulin sensitivity and the systemic inflammatory response in human subjects. *Br J Nutr* 2010; **104**: 1831-1838 [PMID: 20815975 DOI: 10.1017/S0007114510002874]
 - 100 **Wang X**, Ota N, Manzanillo P, Kates L, Zavala-Solorio J, Eidenschenk C, Zhang J, Lesch J, Lee WP, Ross J, Diehl L, van Bruggen N, Kolumam G, Ouyang W. Interleukin-22 alleviates metabolic disorders and restores mucosal immunity in diabetes. *Nature* 2014; **514**: 237-241 [PMID: 25119041 DOI: 10.1038/nature13564]
 - 101 **Panwar H**, Calderwood D, Grant IR, Grover S, Green BD. *Lactobacillus* strains isolated from infant faeces possess potent inhibitory activity against intestinal alpha- and beta-glucosidases suggesting anti-diabetic potential. *Eur J Nutr* 2014; **53**: 1465-1474 [PMID: 24414142 DOI: 10.1007/s00394-013-0649-9]
 - 102 **Napolitano A**, Miller S, Nicholls AW, Baker D, Van Horn S, Thomas E, Rajpal D, Spivak A, Brown JR, Nunez DJ. Novel gut-based pharmacology of metformin in patients with type 2 diabetes mellitus. *PLoS One* 2014; **9**: e100778 [PMID: 24988476 DOI: 10.1371/journal.pone.0100778]
 - 103 **Ahmadi S**, Nagpal R, Wang S, Gagliano J, Kitzman DW, Soleimanian-Zad S, Sheikh-Zeinoddin M, Read R, Yadav H. Prebiotics from acorn and sago prevent high-fat-diet-induced insulin resistance via microbiome-gut-brain axis modulation. *J Nutr Biochem* 2019; **67**: 1-13 [PMID: 30831458 DOI: 10.1016/j.jnutbio.2019.01.011]
 - 104 **Zhang Y**, Gu Y, Ren H, Wang S, Zhong H, Zhao X, Ma J, Gu X, Xue Y, Huang S, Yang J, Chen L, Chen G, Qu S, Liang J, Qin L, Huang Q, Peng Y, Li Q, Wang X, Kong P, Hou G, Gao M, Shi Z, Li X, Qiu Y, Zou Y, Yang H, Wang J, Xu G, Lai S, Li J, Ning G, Wang W. Gut microbiome-related effects of berberine and probiotics on type 2 diabetes (the PREMOT study). *Nat Commun* 2020; **11**: 5015 [PMID: 33024120 DOI: 10.1038/s41467-020-18414-8]
 - 105 **Adeshirlarijaney A**, Gewirtz AT. Considering gut microbiota in treatment of type 2 diabetes mellitus. *Gut Microbes* 2020; **11**: 253-264 [PMID: 32005089 DOI: 10.1080/19490976.2020.1717719]
 - 106 **Jafarnejad S**, Saremi S, Jafarnejad F, Arab A. Effects of a Multispecies Probiotic Mixture on Glycemic Control and Inflammatory Status in Women with Gestational Diabetes: A Randomized Controlled Clinical Trial. *J Nutr Metab* 2016; **2016**: 5190846 [PMID: 27429803 DOI: 10.1155/2016/5190846]
 - 107 **Asemi Z**, Khorrami-Rad A, Alizadeh SA, Shakeri H, Esmailzadeh A. Effects of synbiotic food consumption on metabolic status of diabetic patients: a double-blind randomized cross-over controlled clinical trial. *Clin Nutr* 2014; **33**: 198-203 [PMID: 23786900 DOI: 10.1016/j.clnu.2013.05.015]
 - 108 **Tonucci LB**, Olbrich Dos Santos KM, Licursi de Oliveira L, Rocha Ribeiro SM, Duarte Martino HS. Clinical application of probiotics in type 2 diabetes mellitus: A randomized, double-blind, placebo-controlled study. *Clin Nutr* 2017; **36**: 85-92 [PMID: 26732026 DOI: 10.1016/j.clnu.2015.11.011]
 - 109 **Zhang Q**, Wu Y, Fei X. Effect of probiotics on glucose metabolism in patients with type 2 diabetes mellitus: A meta-analysis of randomized controlled trials. *Medicina (Kaunas)* 2016; **52**: 28-34 [PMID: 26987497 DOI: 10.1016/j.medic.2015.11.008]
 - 110 **Yao K**, Zeng L, He Q, Wang W, Lei J, Zou X. Effect of Probiotics on Glucose and Lipid Metabolism in Type 2 Diabetes Mellitus: A Meta-Analysis of 12 Randomized Controlled Trials. *Med Sci Monit* 2017; **23**: 3044-3053 [PMID: 28638006 DOI: 10.12659/msm.902600]
 - 111 **Kasińska MA**, Drzewoski J. Effectiveness of probiotics in type 2 diabetes: a meta-analysis. *Pol Arch Med Wewn* 2015; **125**: 803-813 [PMID: 26431318 DOI: 10.20452/pamw.3156]
 - 112 **Kocsis T**, Molnár B, Németh D, Hegyi P, Szakács Z, Bálint A, Garami A, Soós A, Márta K, Solymár M. Probiotics have beneficial metabolic effects in patients with type 2 diabetes mellitus: a meta-analysis of randomized clinical trials. *Sci Rep* 2020; **10**: 11787 [PMID: 32678128 DOI: 10.1038/s41598-020-68440-1]
 - 113 **Tao YW**, Gu YL, Mao XQ, Zhang L, Pei YF. Correction to: Effects of probiotics on type II diabetes mellitus: a meta-analysis. *J Transl Med* 2020; **18**: 105 [PMID: 32111256 DOI: 10.1186/s12967-020-02274-3]
 - 114 **Tao YW**, Gu YL, Mao XQ, Zhang L, Pei YF. Effects of probiotics on type II diabetes mellitus: a meta-analysis. *J Transl Med* 2020; **18**: 30 [PMID: 31952517 DOI: 10.1186/s12967-020-02213-2]
 - 115 **Bock PM**, Telo GH, Ramalho R, Sbaraini M, Leivas G, Martins AF, Schaan BD. The effect of probiotics, prebiotics or synbiotics on metabolic outcomes in individuals with diabetes: a systematic review and meta-analysis. *Diabetologia* 2021; **64**: 26-41 [PMID: 33047170 DOI: 10.1007/s00125-020-05295-1]
 - 116 **Ho J**, Reimer RA, Doulla M, Huang C. Effect of prebiotic intake on gut microbiota, intestinal permeability and glycemic control in children with type 1 diabetes: study protocol for a randomized controlled trial. *Trials* 2016; **17**: 347 [PMID: 27456494 DOI: 10.1186/s13063-016-1486-y]
 - 117 **Mishra SP**, Wang S, Nagpal R, Miller B, Singh R, Taraphder S, Yadav H. Probiotics and Prebiotics for the Amelioration of Type 1 Diabetes: Present and Future Perspectives. *Microorganisms* 2019; **7** [PMID: 30832381 DOI: 10.3390/microorganisms7030067]
 - 118 **Didari T**, Solki S, Mozaffari S, Nikfar S, Abdollahi M. A systematic review of the safety of

- probiotics. *Expert Opin Drug Saf* 2014; **13**: 227-239 [PMID: [24405164](#) DOI: [10.1517/14740338.2014.872627](#)]
- 119 **Matsuzaki T**, Nagata Y, Kado S, Uchida K, Kato I, Hashimoto S, Yokokura T. Prevention of onset in an insulin-dependent diabetes mellitus model, NOD mice, by oral feeding of *Lactobacillus casei*. *APMIS* 1997; **105**: 643-649 [PMID: [9298104](#) DOI: [10.1111/j.1699-0463.1997.tb05066.x](#)]
 - 120 **Xie Y**, Xiao M, Ni Y, Jiang S, Feng G, Sang S, Du G. *Alpinia oxyphylla* Miq. Extract Prevents Diabetes in Mice by Modulating Gut Microbiota. *J Diabetes Res* 2018; **2018**: 4230590 [PMID: [29967794](#) DOI: [10.1155/2018/4230590](#)]
 - 121 **Marcial GE**, Ford AL, Haller MJ, Gezan SA, Harrison NA, Cai D, Meyer JL, Perry DJ, Atkinson MA, Wasserfall CH, Garrett T, Gonzalez CF, Brusko TM, Dahl WJ, Lorca GL. *Lactobacillus johnsonii* N6.2 Modulates the Host Immune Responses: A Double-Blind, Randomized Trial in Healthy Adults. *Front Immunol* 2017; **8**: 655 [PMID: [28659913](#) DOI: [10.3389/fimmu.2017.00655](#)]
 - 122 **de Goffau MC**, Luopajarvi K, Knip M, Ilonen J, Ruotula T, Härkönen T, Orivuori L, Hakala S, Welling GW, Harmsen HJ, Vaarala O. Fecal microbiota composition differs between children with β -cell autoimmunity and those without. *Diabetes* 2013; **62**: 1238-1244 [PMID: [23274889](#) DOI: [10.2337/db12-0526](#)]
 - 123 **Tai N**, Peng J, Liu F, Gulden E, Hu Y, Zhang X, Chen L, Wong FS, Wen L. Microbial antigen mimics activate diabetogenic CD8 T cells in NOD mice. *J Exp Med* 2016; **213**: 2129-2146 [PMID: [27621416](#) DOI: [10.1084/jem.20160526](#)]
 - 124 **Giongo A**, Gano KA, Crabb DB, Mukherjee N, Novelo LL, Casella G, Drew JC, Ilonen J, Knip M, Hyöty H, Veijola R, Simell T, Simell O, Neu J, Wasserfall CH, Schatz D, Atkinson MA, Triplett EW. Toward defining the autoimmune microbiome for type 1 diabetes. *ISME J* 2011; **5**: 82-91 [PMID: [20613793](#) DOI: [10.1038/ismej.2010.92](#)]
 - 125 **de Goffau MC**, Fuentes S, van den Bogert B, Honkanen H, de Vos WM, Welling GW, Hyöty H, Harmsen HJ. Aberrant gut microbiota composition at the onset of type 1 diabetes in young children. *Diabetologia* 2014; **57**: 1569-1577 [PMID: [24930037](#) DOI: [10.1007/s00125-014-3274-0](#)]
 - 126 **Brown CT**, Davis-Richardson AG, Giongo A, Gano KA, Crabb DB, Mukherjee N, Casella G, Drew JC, Ilonen J, Knip M, Hyöty H, Veijola R, Simell T, Simell O, Neu J, Wasserfall CH, Schatz D, Atkinson MA, Triplett EW. Gut microbiome metagenomics analysis suggests a functional model for the development of autoimmunity for type 1 diabetes. *PLoS One* 2011; **6**: e25792 [PMID: [22043294](#) DOI: [10.1371/journal.pone.0025792](#)]
 - 127 **Li Q**, Chang Y, Zhang K, Chen H, Tao S, Zhang Z. Implication of the gut microbiome composition of type 2 diabetic patients from northern China. *Sci Rep* 2020; **10**: 5450 [PMID: [32214153](#) DOI: [10.1038/s41598-020-62224-3](#)]
 - 128 **Kikuchi R**, Irie J, Yamada-Goto N, Kikkawa E, Seki Y, Kasama K, Itoh H. The Impact of Laparoscopic Sleeve Gastrectomy with Duodenojejunal Bypass on Intestinal Microbiota Differs from that of Laparoscopic Sleeve Gastrectomy in Japanese Patients with Obesity. *Clin Drug Investig* 2018; **38**: 545-552 [PMID: [29508311](#) DOI: [10.1007/s40261-018-0638-0](#)]
 - 129 **Li JF**, Lai DD, Ni B, Sun KX. Comparison of laparoscopic Roux-en-Y gastric bypass with laparoscopic sleeve gastrectomy for morbid obesity or type 2 diabetes mellitus: a meta-analysis of randomized controlled trials. *Can J Surg* 2013; **56**: E158-E164 [PMID: [24284156](#) DOI: [10.1503/cjs.026912](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

