



Name of Journal: *World Journal of Clinical Urology*

ESPS Manuscript NO: 29404

Manuscript Type: Original Article

Observational Study

Urine chemokine levels correlate with treatment response to

Phosphodiesterase 4 inhibitor in prostatitis

Pradeep Tyagi, Kim Killinger, Gregory McLennan, Nirmal Jayabalan, Michael

Chancellor, Kenneth Peters

Match Overview

1	Crossref 139 words "Society for Urodynamics and Female Urology, 2012 Winter Meeting, February 28-March 3, 2012, The Roosevelt ...	3%
2	Internet 55 words crawled on 10-May-2015 www.wjgnet.com	1%
3	Crossref 46 words ABBAS KHADRA, "Interleukin-8 levels in seminal plasma in chronic prostatitis/chronic pelvic pain syndrome and	1%
4	Internet 24 words crawled on 29-Jul-2010 www.faqs.org	1%
5	Crossref 24 words "Non-discussed Poster Presentations, ICS/IUGA 2010", International Urogynecology Journal, 06/2011	1%
6	Internet 19 words crawled on 25-May-2012 tuberculosis.newsbeet.com	<1%

学术搜索

找到约 210 条结果 (用时0.06秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[HTML] Treatment of rheumatoid arthritis with a recombinant human tumor necrosis factor receptor (p75)-Fc fusion protein

LW Moreland, SW Baumgartner... - ... England Journal of ..., 1997 - Mass Medical Soc

... aminotransferase, total protein, and albumin), and analysis of a clean-catch urine specimen with ... Measures of disease activity moved toward base-line levels after the cessation of TNFR ... of TNFR:Fc in vivo is probably minimal because of the high concentration of immunoglobulin ...

被引用次数: 1683 相关文章 所有 8 个版本 引用 保存

New developments in the diagnosis and treatment of chronic prostatitis/chronic pelvic pain syndrome

M Pontari, L Giusto - Current opinion in urology, 2013 - journals.lww.com

... In 20 patients with CPPS who were treated, the levels of B7-H3 were higher ... T-cell homing to the prostate was dependent on expression of the chemokine receptor CXCR3 ... Other options include PDE5 inhibitor tadalafil [24], which can treat lower urinary tract symptoms, erectile ...

被引用次数: 30 相关文章 所有 4 个版本 引用 保存

Immunomodulatory effects of quinolones

A Dalhoff, I Shalit - The Lancet infectious diseases, 2003 - Elsevier

... an immunomodulatory effect, drug concentrations far beyond clinically relevant levels were examined ... On the other hand, urine concentrations of, for example, ciprofloxacin amount to 600 ... concentrations at various infection sites far exceed the cytotoxic threshold concentration. ...

被引用次数: 238 相关文章 所有 7 个版本 引用 保存

Method for diagnosing and treating chronic pelvic pain syndrome

RB Alexander, S Ponniah - US Patent 6,180,355, 2001 - freepatentsonline.com

... leukocytosis in expressed prostatic secretions or sediment in a post-massage urine sample. ... patent database, by searching, eg.using a boolean search such as "chemokine /2 (antagonist ... 1998

学术搜索

找到约 234 条结果 (用时0.07秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☐ 包含引用

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[HTML] **Treatment of rheumatoid arthritis with a recombinant human tumor necrosis factor receptor (p75)-Fc fusion protein**

LW Moreland, SW Baumgartner... - ... England Journal of ..., 1997 - Mass Medical Soc
... total protein, and albumin), and analysis of a clean-catch **urine** specimen with ... by objective laboratory measures (erythrocyte sedimentation rate and C-reactive protein **level**) (Table 2 ... Measures of disease activity moved toward base-line **levels** after the cessation of TNFR:Fc ...

被引用次数: 1696 相关文章 所有 8 个版本 引用 保存

New developments in the diagnosis and treatment of chronic prostatitis/chronic pelvic pain syndrome

M Pontari, L Giusto - Current opinion in urology, 2013 - journals.lww.com
... In 20 patients with CPPS who were **treated**, the **levels** of B7-H3 were higher ... T-cell homing to the prostate was dependent on expression of the **chemokine** receptor CXCR3 ... Other options include PDE5 **inhibitor** tadalafil [24] , which can **treat** lower **urinary** tract symptoms, erectile ...

被引用次数: 31 相关文章 所有 4 个版本 引用 保存

Mechanism of action of phosphodiesterase type 5 inhibition in metabolic syndrome - associated prostate alterations: An experimental study in the rabbit

A Morelli, P Comeglio, S Filippi, E Sarchielli... - The ..., 2013 - Wiley Online Library
... erectile dysfunction (ED), and benign prostate hyperplasia (BPH)-associated lower **urinary** tract symptoms ... 2b) and normalized to RD **levels** by testosterone supplementation (Fig. ... Correlations coefficients (r) and **level** of significance (P-value) are derived from univariate analysis. ...

被引用次数: 39 相关文章 所有 5 个版本 引用 保存

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 1,150 条结果 (用时 0.80 秒)

Google 学术: Urine chemokine levels correlate with treatment response to Phosphodiesterase 4 inhibitor in prostatitis

Treatment of rheumatoid arthritis with a recombinant ... - Moreland - 被引用次数: 1696

... in the diagnosis and treatment of chronic prostatitis/ ... - Pontari - 被引用次数: 31

... of action of phosphodiesterase type 5 inhibition in ... - Morelli - 被引用次数: 39

New treatments for chronic prostatitis/chronic pelvic pain syndrome

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2837110/> - 翻译此页

作者: AC Strauss - 2010 - 被引用次数: 38 - 相关文章

2010年2月9日 - New treatments for chronic prostatitis/chronic pelvic pain syndrome Allodynia is a painful response to a stimulus that is not normally painful. ... In an RCT of the COX2 inhibitor rofecoxib, the NIH-CPSI total and Postprostatic massage urine and seminal plasma NGF levels have been shown to correlate ...

缺少字词: phosphodiesterase

Promising Molecular Targets and Biomarkers for Male BPH and LUTS

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3851290/> - 翻译此页

作者: M Gharaee-Kermani - 2013 - 被引用次数: 10 - 相关文章

跳到 Targeted Therapeutic Strategies for the Treatment of BPH/LUTS - I. 5 α -Reductase Inhibitors ... for the treatment of BPH/LUTS for ... high levels plasma and urine ... of phosphodiesterase 5 inhibitors ... prostate in the context of prostatitis ... levels of CXC-type chemokine ... Upregulation of TGF β 1 correlates ...

Inflammation, a Key Event in Cancer Development | Molecular Cancer