

Curcumin Improved Regulatory T Cells in Gut-Associated Lymphoid Tissue in Mic



全部 图片 视频 新闻 购物 地图 图书

找到约 64,800 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Google 学术: Curcumin Improved Regulatory T Cells in Gut-Associated Lymphoid Tissue in Mice with Experimental Colitis](#)

[Intestinal CD103+ dendritic cells: master regulators of ...](#) - Scott - 被引用次数: 138

[... induced HSP are targets of regulatory T cells: a role for ...](#) - Wieten - 被引用次数: 59

[Dietary polyphenols can modulate the intestinal ...](#) - Romier - 被引用次数: 103

[Essential role for CD103 in the T cell-mediated regulation of ...](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC2213206/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2213206/) ▾

Essential role for CD103 in the T cell-mediated regulation of experimental colitis ... However, wild-type CD4<sup>+</sup>CD25<sup>+</sup> T cells were unable to prevent colitis in ... Non-T cell expression of CD103 is restricted primarily to CD11c<sup>high</sup>MHC class II<sup>high</sup> ... frequency in the gut-associated lymphoid tissue and appears to mediate a ...

[Harnessing Regulatory T Cells for the Treatment of Inflammatory ...](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC4437827/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4437827/)

19 Mar 2015 ... The original description of Treg cells preventing colitis in mice by ... of Treg cells in peripheral tissues may lead to chronic intestinal inflammation. ... the survival of Foxp3<sup>+</sup> Treg cells may be improved with IL-2 and FA supplementation. ... Role of the gut-associated and secondary lymphoid tissue in the ...

[Animal models of ulcerative colitis and their application in drug ...](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC3829622/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3829622/) ▾

12 Nov 2013 ... Experimental colitis can be induced using chemical irritant or bacterial infection. ... Mouse models of inflammatory bowel disease (IBD). ... Natural killer T cells and their associated cytokine, mainly IL-13, are also intimately ... intestinal tract and its associated lymphoid tissue and is recognized by its receptor ...

[Cell stress induced HSP are targets of regulatory T cells: A role for ...](#)

Name of journal: World Journal of Gastroenterology

ESPS Manuscript NO: 25188

Manuscript Type: ORIGINAL ARTICLE

*Basic Study*

**Curcumin improved regulatory T cells in gut-associated lymphoid tissue in colitis mice**

Hai-Mei Zhao, Rong Xu, Xiao-Ying Huang, Shao-Min Cheng, Min-Fang

Huang, Hai-Yang Yue, Xin Wang, Yong Zou, Ai-Ping Lu, Duan-Yong Liu

**Abstract**

### Match Overview

|   |   |     |
|---|---|-----|
| 1 | Internet 61 words<br>crawled on 10-Dec-2009<br><a href="http://www.jleukbio.org">www.jleukbio.org</a>   | 2%  |
| 2 | CrossCheck 25 words<br><a href="#">Immunology of the Skin, 2016.</a>  | 1%  |
| 3 | Internet 24 words<br>crawled on 23-Mar-2014<br><a href="http://trinity.engr.uconn.edu">trinity.engr.uconn.edu</a>                                 | 1%  |
| 4 | Internet 20 words<br>crawled on 02-Feb-2016<br><a href="http://arizona.openrepository.com">arizona.openrepository.com</a>                         | 1%  |
| 5 | Internet 11 words<br>crawled on 06-Nov-2015<br><a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>                                     | <1% |
| 6 | CrossCheck 10 words<br>Bayry, J.. "Rescuing CD4+CD25+ regulatory T-cell funcio ...<br>s in rheumatoid arthritis by cytokine-targeted monoclonal a | <1% |

Curcumin improved regulatory T cells in gut-associated lymphoid tissue in colitis |



全部 图片 视频 新闻 购物 地图 图书

找到约 73,000 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Google 学术: Curcumin improved regulatory T cells in gut-associated lymphoid tissue in colitis mice](#)

[Intestinal CD103+ dendritic cells: master regulators of ...](#) - Scott - 被引用次数: 142

[... induced HSP are targets of regulatory T cells: a role for ...](#) - Wieten - 被引用次数: 59

[Dietary polyphenols can modulate the intestinal ...](#) - Romier - 被引用次数: 103

[Cell stress induced HSP are targets of regulatory T cells: A role for ...](#)

[www.sciencedirect.com/science/article/pii/S0014579307004942](http://www.sciencedirect.com/science/article/pii/S0014579307004942)

T cell responses to heat shock proteins (HSP) have disease suppressive ... GALT, gut associated lymphoid tissue;; GGA, geranylgeranyl acetone;; PG, .... In this model up-regulation of HSP70 and HSP110 was found in multiple mouse tissues .... colitis [46] and it induced improvements in other forms of inflammatory bowel ...

[Animal models of ulcerative colitis and their application in drug ...](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC3829622/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3829622/)

12 Nov 2013 ... Mouse models of inflammatory bowel disease (IBD). ... Natural killer T cells and their associated cytokine, mainly IL-13, are also intimately ..... only in the intestinal tract and its associated lymphoid tissue and is recognized by its ..., barrier regulation and gut microbiota to improve intestinal inflammation.96.

[Harnessing Regulatory T Cells for the Treatment of Inflammatory ...](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC4437827/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4437827/)

19 Mar 2015 ... The original description of Treg cells preventing colitis in mice by ... of Treg cells in peripheral tissues may lead to chronic intestinal inflammation. .... the survival of Foxp3+ Treg cells may be improved with IL-2 and FA supplementation. .... Role of the gut-associated and secondary lymphoid tissue in the ...

[Changes of CD8+ T cells in dextran sulfate sodium-induced colitis ...](#)



全部 图片 新闻 视频 购物 更多 ▾ 搜索工具

找到约 69,200 条结果 (用时 0.77 秒)

## Google 学术 : Curcumin improved regulatory T cells in gut-associated lymphoid tissue in colitis mice

Intestinal CD103+ dendritic cells: master regulators of ... - Scott - 被引用次数 : 144

... induced HSP are targets of regulatory T cells; a role for ... - Wieten - 被引用次数 : 59

Dietary polyphenols can modulate the intestinal ... - Romier - 被引用次数 : 103

## Cell stress induced HSP are targets of regulatory T cells: A ...

[www.sciencedirect.com/science/article/.../S0014579307004942](http://www.sciencedirect.com/science/article/.../S0014579307004942) ▾ 翻译此页

作者 : L Wieten - 2007 - 被引用次数 : 59 - 相关文章

T cell responses to heat shock proteins (HSP) have disease suppressive ... GALT, gut associated lymphoid tissue;; GGA, geranylgeranyl acetone;; PG, .... In this model up-regulation of HSP70 and HSP110 was found in multiple mouse tissues .... colitis [46] and it induced improvements in other forms of inflammatory bowel ...

## Animal models of ulcerative colitis and their application in ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > PubMed Central (PMC) ▾ 翻译此页

作者 : D Low - 2013 - 被引用次数 : 30 - 相关文章

2013年11月12日 - Mouse models of inflammatory bowel disease (IBD). .... Natural killer T cells and their associated cytokine, mainly IL-13, are also intimately .... only in the intestinal tract and its associated lymphoid tissue and is recognized by its .... barrier regulation and gut microbiota to improve intestinal inflammation,96.

## Changes of CD8<sup>+</sup> T cells in dextran sulfate sodium-induced ...

[www.ncbi.nlm.nih.gov/pubmed/22884149](http://www.ncbi.nlm.nih.gov/pubmed/22884149) ▾ 翻译此页

作者 : YF Ye - 2012 - 相关文章

Changes of CD8<sup>+</sup> T cells in dextran sulfate sodium-induced colitis mice ... sodium