



**Name of Journal:** *World Journal of Gastrointestinal Surgery*

**ESPS Manuscript NO:** 29900

**Manuscript Type:** Original Article

*Retrospective Study*

**Critical analysis of feeding jejunostomy following resection of upper gastrointestinal malignancies**

Andrew M. Blakely, MD, Saad Ajmal, MBBS, Rachel E. Sargent, BA, Thomas T. Ng, MD, FACS, Thomas J. Miner, MD, FACS

## Match Overview

1

Internet 33 words  
crawled on 26-Feb-2015  
[expo.jspargo.com](http://expo.jspargo.com)

19

2

Crossref 15 words  
Patel, Sameer H., David A. Kooby, Charles A. Staley, and S  
hishir K. Maithe, "An assessment of feeding jejunostomy ...

&lt;19

3

Crossref 11 words  
Respiratory Medicine, 2014.

&lt;19

4

Crossref 10 words  
"Jejunal Feeding Tube Complications", Gastrointestinal Su  
rgery, 2015.

&lt;19

5

Crossref 10 words  
Blakely, Andrew M., Daithi S. Heffernan, Jane McPhillips, Wi  
lliam G. Cioffi, and Thomas J. Miner. "Elevated C-reactive p

&lt;19

学术搜索

找到约 4,430 条结果 (用时0.12秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

### A prospective, randomized trial of early enteral feeding after resection of upper gastrointestinal malignancy.

MJ Heslin, L Latkany, D Leung, AD Brooks... - Annals of ..., 1997 - ncbi.nlm.nih.gov

... In the trauma and critical care literature, enteral nutrition has been shown to decrease the risk of ... had abdominal cramps or Figure 2 shows the daily total caloric intake as a percent of caloric intake from oral, intravenous, or tube ing, there was an increase in ...

被引用次数: 424 相关文章 所有 11 个版本 引用 保存

### Response to neoadjuvant chemotherapy best predicts survival after curative resection of gastric cancer.

AM Lowy, PF Mansfield, SD Leach, R Pazdur... - Annals of ..., 1999 - ncbi.nlm.nih.gov

... Although this may be a result of patient numbers, response to neoadjuvant chemotherapy remained predictive of survival on multivariate analysis. Further, patients who did respond to neoadjuvant therapy had an actuarial 5-year survival of 83%, despite a 44% incidence of ...

被引用次数: 249 相关文章 所有 5 个版本 引用 保存

### Early enteral nutrition in acutely ill patients: a systematic review

PE Marik, GP Zaloga - Critical care medicine, 2001 - journals.lww.com

... demonstrated at the cellular and tissue level and in animal studies of critical illness, the ... of available studies that compared early with delayed enteral nutrition to provide an estimate of ... When the tests' statistics were not available, given a p value, we computed the corresponding ...

被引用次数: 847 相关文章 所有 8 个版本 引用 保存

### A randomized controlled trial of preoperative oral supplementation with a specialized diet in patients with gastrointestinal cancer

L Gianotti, M Braga, L Nespoli, G Radaelli, A Beneduce... - Gastroenterology, 2002 - Elsevier

The perioperative administration of an enteral diet enriched with arginine, ω 3 fatty acids, and



学术搜索

找到约 3,920 条结果 (用时0.09秒)

文章

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

我的图书馆

### A prospective, randomized trial of early enteral feeding after resection of upper gastrointestinal malignancy.

MJ Heslin, L Latkany, D Leung, AD Brooks... - Annals of ..., 1997 - ncbi.nlm.nih.gov

... In the trauma and critical care literature, enteral nutrition has been shown to decrease the risk of septic complications ... Multivariate analyses of the primary endpoints were carried out using step-wise logistic regressions. ... control group by intent-to-treat or by subgroup analysis. ...

被引用次数: 425 相关文章 所有 11 个版本 引用 保存

时间不限

2016以来

2015以来

2012以来

自定义范围...

### Response to neoadjuvant chemotherapy best predicts survival after curative resection of gastric cancer.

AM Lowy, PF Mansfield, SD Leach, R Pazdur... - Annals of ..., 1999 - ncbi.nlm.nih.gov

... For all analyses,  $p < 0.05$  was considered statistically significant ... chemotherapy plus resection to resection alone, we thought it of interest to analyze survival as ... of patient numbers, response to neoadjuvant chemotherapy remained predictive of survival on multivariate analysis. ...

被引用次数: 253 相关文章 所有 5 个版本 引用 保存

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

### Early enteral nutrition in acutely ill patients: a systematic review

PE Marik, GP Zaloga - Critical care medicine, 2001 - journals.lww.com

... may reduce septic and nonseptic complications and improve the outcome of the critically ill and ... been demonstrated at the cellular and tissue level and in animal studies of critical illness, the ... 12 were excluded; the remaining 15 trials (31-45) were included in this meta-analysis. ...

被引用次数: 857 相关文章 所有 8 个版本 引用 保存

☒ 包括专利☐ 包含引用

[全部](#)[图片](#)[新闻](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 34,400 条结果 (用时 0.65 秒)

## Google 学术: Critical analysis of feeding jejunostomy following resection of upper gastrointestinal malignancies

... after resection of upper gastrointestinal malignancy. - Heslin - 被引用次数: 427

... survival after curative resection of gastric cancer. - Lowy - 被引用次数: 255

... nutrition in acutely ill patients: a systematic review - Marik - 被引用次数: 861

### Resection of Upper Gastrointestinal Malignancy - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1191079/> - 翻译此页

作者: MJ Heslin - 1997 - 被引用次数: 428 - 相关文章

... after major upper GI surgery for malignancy with early enteral feeding are limited. Methods ... underwent resection and were randomized to IEF via jejunostomy tube or control (CNTL). Tube ... In the trauma and critical care literature, enteral nutrition has ... before operation, and the study was conducted following approval ...

### A prospective, randomized trial of early enteral feeding after resection ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1191079/> - 翻译此页

作者: MJ Heslin - 1997 - 被引用次数: 426 - 相关文章

... trial of early enteral feeding after resection of upper gastrointestinal malignancy. .... to jejunostomy feeding after major abdominal trauma—a critical analysis.

### Evaluation of supporting role of early enteral feeding via tube ... - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3587889/> - 翻译此页

作者: A Ghafouri - 2012 - 被引用次数: 7 - 相关文章

Evaluation of supporting role of early enteral feeding via tube jejunostomy following resection of upper gastrointestinal tract. ... One of the important side effects in upper GI tract malignancies is malnutrition.