



Autotransplantation of a premolar to the maxillary anterior region in youn



网页

图片

新闻

视频

更多

搜索工具

找到约 84,900 条结果 (用时 0.73 秒)

[PDF] Case Report

www.aapd.org/assets/1/25/160-6.pdf 翻译此页

Case Report: Autotransplantation for a Missing Permanent Maxillary Incisor. Jahnavi Rao, DDS ... patients with missing permanent anterior teeth, they should be,

Tooth auto-transplantation as an alternative treatment option ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) 翻译此页

作者: T Nimčenko - 2013 - 被引用次数: 4 - 相关文章

Therefore, patient's young age is a restriction for implantation and a particular ... Thus, tooth auto-transplantation can be a good choice for treatment. ... In cases of severe ectopic position of maxillary canines (ectopic canines present about Yu Z, Bayer T. A long-term study of 370 autotransplanted premolars. Part IV. Root ...

[PDF] Planning esthetic treatment after avulsion of maxillary inci...

www.moderndentistrymedia.com/mar.../zachrisson.pdf 翻译此页

作者: BU Zachrisson - 被引用次数: 28 - 相关文章

occurs in children from 7 to 9 years of age, when the ... This case report describes an innovative solution that combined ... should be treated with root canal therapy. The case report included here ... to the anterior region. ... There was a small apical perforation ... autotransplantation of premolars to the maxillary anterior.

Tooth Autotransplantation as a Treatment Option: A Review ...

www.researchgate.net/.../50590815_Tooth_Autotransplantation... 翻译此页

2015年6月4日 - ABSTRACT Autotransplantation can provide patients with all the ... but is seldom considered as a treatment option, in large part because of a lack Maxillary anterior region within postoperative 14 days, and anterior/premolar donors. ... of

Name of Journal: World Journal of Stomatology

ESPS Manuscript NO: 20684

Manuscript type: CASE REPORT

12

Autotransplantation of a premolar to the maxillary anterior region in young children- how long should the donor root be? A case report

Uri Zilberman, Amiqam Zaguri

Abstract

Autotransplantation of premolars to anterior region after incisor loss due to trauma is accepted as the best restoration procedure with very long follow-ups. There are two main protocols: premolars with only two thirds of the root or premolars with complete root development. Premolars with two third of the root completed remain vital and show complete pulp obliteration while premolars with closed apex require root canal treatment. The problem arises when the child is very young and the root of the donor premolar is developed

Match Overview

1	Internet 47 words crawled on 09-Apr-2010 cda-adc.ca	2%
2	CrossCheck 28 words Denys, D., M. Shahbazian, R. Jacobs, A. Laenen, J. Wyatt, F. Vinckier, and G. Willems. "Importance of root developme	1%
3	Internet 16 words crawled on 03-Apr-2012 www.moderndentistrymedia.com	1%
4	Internet 15 words crawled on 24-Jun-2013 bimetra.be	1%
5	CrossCheck 10 words Satoshi Imazato. "Potential efficacy of GTR and autogenous bone graft for autotransplantation to recipient sites with	<1%
6	CrossCheck 9 words A. Mendoza-Mendoza. "Retrospective long-term evaluation of autotransplantation of premolars to the central incisor re	<1%
7	CrossCheck 8 words Winfried Harzer. "Autotransplantation of first premolar to replace a maxillary incisor - 3D-volume tomography for eva	<1%
8	Internet 8 words crawled on 10-Feb-2015 www.sciencepub.net	<1%
	CrossCheck 8 words	<1%



Autotransplantation of a premolar to the maxillary anterior region in young ch



网页

图片

视频

新闻

更多

搜索工具

找到约 67,900 条结果 (用时 0.85 秒)

[PDF] Case Report

www.aapd.org/assets/1/25/160-6.pdf 翻译此页

Case Report: Autotransplantation for a Missing Permanent Maxillary Incisor. Jahnavi Rao, DDS ... patients with missing permanent anterior teeth, they should be.

Tooth auto-transplantation as an alternative treatment option ...

[www.ncbi.nlm.nih.gov/.../PubMed Central \(PMC\)](http://www.ncbi.nlm.nih.gov/.../PubMed Central (PMC)) 翻译此页

作者: T Nimčenko - 2013 - 被引用次数: 4 - 相关文章

Thus, tooth auto-transplantation can be a good choice for treatment. ... In cases of severe ectopic position of maxillary canines (ectopic canines present about ... and 80-87% over 10 years with transplanted anterior teeth with closed apices. Yu Z, Bayer T. A long-term study of 370 autotransplanted premolars. Part IV. Root ...

Tooth Autotransplantation as a Treatment Option: A Review ...

www.researchgate.net/.../50590815_Tooth_Autotransplantation... - 翻译此页

2015年9月26日 - ABSTRACT Autotransplantation can provide patients with all the ... is seldom considered as a treatment option, in large part because of a lack of ... Match case T. A long-term study of 370 autotransplanted premolars. Part IV. Root [A clinical study on orthodontic treatment after tooth autotransplantation].

[PDF] Planning esthetic treatment after avulsion of maxillary inci...

www.moderndentistrymedia.com/mar.../zachrisson.pdf 翻译此页

作者: BU Zachrisson - 被引用次数: 30 - 相关文章

Most traumatic dental injuries occur in childhood and ... This case report describes an innovative solution that combined ... after avulsion of incisors, clinicians working in cooperation can optimize the ... to the anterior region. ... autotransplantation of

[网页](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 62,500 条结果 (用时 0.94 秒)

[PDF] Case Report

www.aapd.org/assets/1/25/160-6.pdf ▾ [翻译此页](#)

Case Report: Autotransplantation for a Missing Permanent Maxillary Incisor. Jahnavi Rao, DDS ... patients with missing permanent anterior teeth, they should be.

Tooth Autotransplantation as a Treatment Option: A Review ...

www.researchgate.net/.../50590815_Tooth_Autotransplantation... - [翻译此页](#)

7 天前 - Autotransplantation can provide patients with all the benefits of a natural tooth, but is seldom considered as a treatment option, in large part ...

Tooth auto-transplantation as an alternative treatment option ...

[www.ncbi.nlm.nih.gov/.../PubMed Central \(PMC\)](http://www.ncbi.nlm.nih.gov/.../PubMed Central (PMC)) ▾ [翻译此页](#)

作者: T Nimčenko - 2013 - 被引用次数: 5 - 相关文章

Thus, tooth auto-transplantation can be a good choice for treatment. ... method in case of lost teeth (because of its predictable and long-term results), ... third molars, followed by mandibular second premolars, maxillary lateral incisors. and complete root formation transplanted into the upper incisor region over 4 years.

[PDF] Planning esthetic treatment after avulsion of maxillary incis...

www.moderndentistrymedia.com/mar.../zachrisson.pdf ▾ [翻译此页](#)

作者: BU Zachrisson - 被引用次数: 30 - 相关文章