

1 Name of journal: World Journal of Nephrology

2 ESPS Manuscript NO: 15198

3 Columns: Original Article

4

5 *Randomized Controlled Trial*

6

7 Changes in ³urinary excretion of water and sodium transporters during amiloride and
8 bendroflumethiazide treatment

9

10 Janni Majgaard ²Jensen, Frank Holden Mose, Anna-Ewa Oczachowska-Kulik, Jesper
11 Noergaard Bech, Robert A Fenton, Erling Bjerregaard Pedersen

12

13 Abstract

14 AIM: To quantify changes in urinary excretion of ¹aquaporin2 water channels (u-AQP2),
15 the sodium-potassium-chloride co-transporter (u-NKCC2) and the epithelial sodium
16 channels (u-ENaC) during treatment with bendroflumethiazide (BETZ), amiloride and

Match Overview

- 1 **CrossCheck** 15 words <1%
Kurnatowska, I., P. Grzelak, A. Masajtis-Zagajewska, M. Kacz
marska, L. Stefa czyk, C. Vermeer, K. Maresz, M. Nowicki, L.
- 2 **Internet** 13 words <1%
crawled on 01-Apr-2015
paperity.org
- 3 **CrossCheck** 8 words <1%
Yang, A., J. J. Low, A. Leon, H. S. Rasmussen, S. A. Therwan
i, M. Bruschi, L. Santucci, M. Bonsano, G. Candiano, G. M. G

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 23,400 条结果 (用时 0.68 秒)

Google 学术: Changes in urinary excretion of water and sodium transporters during amiloride and bendroflumethiazide treatment

... -2, Na-Cl co-transporter, and epithelial sodium channel - Kim - 被引用次数: 96

... treatment inhibits amiloride-sensitive sodium transport ... - Thomsen - 被引用次数: 25

Diuretic therapy - Wood - 被引用次数: 553

acute angiotensin ii infusions elicit pressure natriuresis in ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: D Zhao - 2008 - 被引用次数: 15 - 相关文章

2008年5月26日 - In the first group, we measured changes in urinary sodium excretion (UNaV) in ... sodium transporters with amiloride (5 mg/kg BW) plus bendroflumethiazide (12 mg/kg BW). ... Normal salt diet along with tap water was provided. but not below 100 mm Hg during treatment with AM plus BFTZ in dogs (9).

Enhanced Distal Nephron Sodium Reabsorption in Chronic ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: D Zhao - 2009 - 被引用次数: 43 - 相关文章

2009年6月1日 - Ang II-infused mice had lower urinary sodium excretion (0.16 ± 0.04 ... Urinary Ang II concentrations, measured during distal blockade, were ... filtered sodium, amiloride, bendroflumethiazide, renal plasma flow, glomerular filtration rate ... II stimulates

[网页](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 73,400 条结果 (用时 0.78 秒)

Google 学术: Changes in urinary excretion of water and sodium transporters during amiloride and bendroflumethiazide treatment

... -2, Na-Cl co-transporter, and epithelial sodium channel - Kim - 被引用次数: 98

... treatment inhibits amiloride-sensitive sodium transport ... - Thomsen - 被引用次数:

25

Diuretic therapy - Wood - 被引用次数: 556

acute angiotensin ii infusions elicit pressure natriuresis in ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: D Zhao - 2008 - 被引用次数: 16 - 相关文章

2008年5月26日 - In the first group, we measured changes in urinary sodium excretion (UNaV) in ... sodium transporters with amiloride (5 mg/kg BW) plus bendroflumethiazide (12 mg/kg BW). ... Normal salt diet along with tap water was provided. but not below 100 mm Hg during treatment with AM plus BFTZ in dogs (9).

Enhanced Distal Nephron Sodium Reabsorption in Chronic ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: D Zhao - 2009 - 被引用次数: 43 - 相关文章

2009年6月1日 - Ang II-infused mice had lower urinary sodium excretion (0.16 ± 0.04 ... Urinary Ang II concentrations, measured during distal blockade, were ... filtered sodium, amiloride, bendroflumethiazide, renal plasma flow, glomerular filtration rate ... II stimulates