

91100_Auto_Edited -check.docx

Prevalence and outcome of sarcopenia in non-alcoholic fatty liver disease

Abstract

BACKGROUND

Nonalcoholic fatty liver disease (NAFLD) includes a spectrum of conditions, progressing from mild steatosis to advanced fibrosis. Sarcopenia, characterized by decreased muscle strength and mass, shares common pathophysiological traits with NAFLD. An association exists between sarcopenia and increased NAFLD prevalence. However, data on the prevalence of sarcopenia in NAFLD and its impact on the outcomes of NAFLD remain inconsistent.

AIM

To analyze the prevalence and outcomes of sarcopenia in patients with NAFLD.

METHODS

We conducted a comprehensive search for relevant studies in MEDLINE, Embase, and Scopus from their inception to June 2023. We included studies that focused on patients with NAFLD, reported the prevalence of sarcopenia as the primary outcome, and examined secondary outcomes, such as liver fibrosis and other adverse events. We also used the Newcastle-Ottawa scale for quality assessment.

RESULTS

Of the 29 studies included, the prevalence of sarcopenia in NAFLD varied widely (1.6% to 63.0%), with 20 studies reporting a prevalence of more than 10.0%. Substantial heterogeneity was noted in the measurement modalities for sarcopenia. Sarcopenia was associated with a higher risk of advanced fibrosis (odd ratio: 1.97, 95% confidence interval: 1.44-2.70). Increased odds were consistently observed in fibrosis assessment through biopsy, NAFLD fibrosis score/body mass index, aspartate aminotransferase to alanine aminotransferase ratio, diabetes (BARD) score, and transient elastography, whereas the fibrosis-4 score showed no such association.

Sarcopenia in NAFLD was associated with a higher risk of steatohepatitis, insulin resistance, cardiovascular risks, and mortality.

CONCLUSION

This systematic review highlights the critical need for standardized diagnostic criteria and measurement methods for sarcopenia in NAFLD patients. The variability in study designs and assessment methods for sarcopenia and liver fibrosis may account for the inconsistent findings. This review demonstrates the multidimensional impact of sarcopenia on NAFLD, indicating its importance beyond liver-related events to include cardiovascular risks, mortality, and metabolic complications.

Key Words: Non-alcoholic fatty liver disease; Sarcopenia; Hepatic fibrosis; Low muscle mass; Hand grip strength; Bioelectric impedance analysis; Dual X-ray absorptiometry

Giri S, Anirvan P, Angadi S, Singh A, Lavekar A. Prevalence and outcome of sarcopenia in non-alcoholic fatty liver disease. *World J Gastrointest Pathophysiol* 2024; In press

Core Tip: The prevalence of sarcopenia in nonalcoholic fatty liver disease (NAFLD) varies widely. Sarcopenia in NAFLD is consistently associated with a higher risk of advanced fibrosis. In addition to liver-related events, sarcopenia in NAFLD is associated with adverse outcomes, including an increased risk of nonalcoholic steatohepatitis, mortality, cardiovascular risks, and metabolic complications. The heterogeneity in prevalence and associations highlights the importance of accurately defining measurement modalities and cutoff criteria. Establishing consensus guidelines is crucial for advancing research and enhancing clinical management in the complex relationship between sarcopenia and NAFLD.

INTRODUCTION

Nonalcoholic fatty liver disease (NAFLD) includes a spectrum of liver conditions, beginning with mild steatosis and potentially advancing through steatohepatitis, fibrosis, and cirrhosis^[1]. Sarcopenia, prevalent in aging populations, is defined by a reduction in muscle strength and/or function, which is often evidenced by a decrease in muscle mass observed in cross-sectional imaging^[2]. Sarcopenic obesity is characterized by the concurrent presence of sarcopenia and increased fat mass, typically measured by body mass index (BMI) or waist circumference^[3]. Factors such as hyperammonemia, endotoxemia, and endocrine disturbances, including insulin resistance and decreased testosterone levels, contribute to the increased prevalence of sarcopenia in individuals with liver cirrhosis^[4]. Several pathophysiological similarities exist between NAFLD and sarcopenia, including insulin resistance, myostatin and adiponectin dysregulation, hormonal imbalances, chronic inflammation, impaired glucose uptake, and myosteatosis^[5,6].

Studies have reported an association of sarcopenia with a higher prevalence of NAFLD and more severe liver damage in individuals with NAFLD. An increased fat mass in patients with NAFLD is associated with a higher incidence of sarcopenic obesity. A meta-analysis of five cross-sectional studies involving 27804 patients identified an increased risk of NAFLD in individuals with sarcopenia^[7]. However, data regarding the prevalence of sarcopenia among patients with NAFLD are inconsistent. Moreover, the effects of sarcopenia on the outcomes of patients with NAFLD remain unclear. Thus, this systematic review aimed to analyze the prevalence and impact of sarcopenia in individuals with NAFLD.

MATERIALS AND METHODS

The current meta-analysis was conducted in accordance with the updated Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines^[8].

Database search

We searched for relevant studies in MEDLINE, Embase, and Scopus from the inception of these databases until June 31, 2023 by using the following keywords: (NAFLD OR Fatty liver OR Steatotic liver disease OR MAFLD OR NASH) AND

¹ (Sarcopenia OR sarcopenic OR Muscle wasting). The titles and abstracts of the retrieved studies were screened by two independent reviewers, who then assessed the full texts for eligibility before inclusion. Furthermore, the bibliographies of the included studies were reviewed to identify additional for relevant studies. Any disagreements between the two independent reviewers were resolved by a third reviewer.

¹ **Study inclusion**

Both prospective and retrospective studies that met the following criteria were included in this systematic review: (1) Studies including patients with NAFLD as determined by serology, ultrasonography (USG), transient elastography (TE), or magnetic resonance imaging (MRI); (2) studies examining the prevalence of sarcopenia as the primary outcome; and (3) studies evaluating the effect of sarcopenia on the risk of liver fibrosis or other adverse outcomes as secondary outcomes. Editorials, correspondences, case reports, case series, and review articles were excluded. Moreover, studies with insufficient or irrelevant clinical data were excluded.

Data extraction and quality assessment

Two reviewers independently extracted the data, and a third reviewer resolved any disagreements. The extracted information for each study included the title, first author, year of publication, country, number of patients, age and sex distribution, BMI, prevalence of metabolic syndrome parameters, diagnostic method used for fatty liver diagnosis, assessment method for sarcopenia, prevalence of sarcopenia, and the effect of sarcopenia on the outcomes of patients with NAFLD. The quality of the included studies was assessed by two independent reviewers by using the Newcastle-Ottawa Scale for cohort studies^[9,10]. In case of a disagreement, a third reviewer was consulted.

RESULTS

Study characteristics and quality assessment

A total of 2134 records were identified using the predefined search strategy, with 29 studies ultimately included in the systematic review. Figure 1 illustrates the PRISMA flowchart detailing the study selection and inclusion process. Table 1 presents the baseline characteristics and outcomes of the included studies. The majority of the studies were from Asia, followed by North America. The mean age of participants in the included studies ranged from 41.9 to 67.8 years, and the proportion of male participants varied from 19.5% to 89.8%. Only three studies included biopsy-proven NAFLD cases^[11-13], whereas the remaining studies used noninvasive methods for NAFLD diagnosis. Among the studies using noninvasive modalities, four used serological tests^[10,14-16], 16 used USG^[17-32], two used MRI^[33,34], and four used TE with controlled attenuation parameters^[35-38]. Bioimpedance analysis was the most common modality used for the assessment of sarcopenia (16 studies)^[11,12,17,18,20-23,28,30-37], followed by dual X-ray absorptiometry (DEXA; 7 studies)^[10,19,14,25,26,28,29], computed tomography (1 study)^[13], hand-grip strength (1 study)^[15], and MRI (1 study)^[33]. Both the studies by Wijampreecha *et al*^[18] and Kim *et al*^[23] analyzed the data from the third National Health and Nutrition Examination Survey conducted from 1988 to 1994 but analyzed different outcomes. Six studies were of good quality^[1-13,30,35,37], 20 were of fair quality^[10,14,15,17-23,24-29,31-34,38], and three were of poor quality^[16,24,35].

Prevalence of sarcopenia in NAFLD

A total of 24 studies reported the prevalence of sarcopenia in NAFLD. The overall prevalence varied significantly, from 1.6% when determined using MRI^[33] to 63.0% when assessed using DEXA^[24]. Four studies reported a prevalence of less than 10.0%^[17,20,33,34], 14 reported a prevalence of 10.0%-30.0%^[10,11,13,14,19,22,26-31,35,38], four reported a prevalence of 30.0%-50.0%^[12,14,33,38], and two reported a prevalence of more than 50.0%^[24,36]. In studies using DEXA, the prevalence of sarcopenia ranged from 12.2% when using an appendicular skeletal muscle mass (ASM)/BMI (cutoff of 0.789 in men and 0.521 in women)^[10] to 63.0% when using ASM/weight (cutoff of 29.0 in men and 22.9 in women^[25]). In studies using BIA, the prevalence of sarcopenia ranged from 4.4% by using a combination of ASM/weight, ASM/height²,

and ASM/BMI^[34] to 54.8% using ASM/height² (cutoff of 7.0 in men and 5.7 in women)^[32].

Risk of advanced fibrosis in NAFLD with sarcopenia

Ten studies examined the correlation between advanced fibrosis and sarcopenia in NAFLD^[11-13,18,22,29,35,37]. The combined analysis of these studies revealed that sarcopenia was associated with a higher risk of advanced fibrosis, with an odds ratio (OR) of 1.97 [95% confidence interval (95%CI): 1.44-2.70; $I^2 = 79.8\%$]. When considering individual modalities for the assessment of fibrosis, including biopsy, NAFLD fibrosis scores/BMI, the ratio of aspartate aminotransferase to alanine aminotransferase, diabetes (NFS/BARD) scores, and TE, sarcopenia was consistently associated with an increased risk of advanced fibrosis with ORs of 1.98 (95%CI: 1.39-2.82; $I^2 = 0.0\%$), 2.09 (95%CI: 1.55-2.81; $I^2 = 0.0\%$), and 3.71 (95%CI: 2.62-5.24; $I^2 = 0.0\%$), respectively, indicating no heterogeneity (Figure 2). However, when using the fibrosis-4 (FIB-4) score, no association was observed between sarcopenia and advanced fibrosis with an OR of 1.38 (95%CI: 0.96-1.99; $I^2 = 63.3\%$), indicating moderate heterogeneity.

Risk of other events in NAFLD with sarcopenia

Eight studies explored the outcome of NAFLD with sarcopenia in addition to the increased risk of advanced hepatic fibrosis. Koo *et al*^[11] reported a higher risk of nonalcoholic steatohepatitis (NASH) in NAFLD with sarcopenia (aOR: 2.59; 95%CI: 1.22-5.48). Petta *et al*^[12] reported that the prevalence of NASH was higher in the presence of sarcopenia (88.7% vs 76.3% in nonsarcopenic cases, $P = 0.01$). Two studies identified sarcopenia as a predictor of mortality in patients with NAFLD, with adjusted hazard ratios (aHRs) of 1.78 (95%CI: 1.16-2.73)^[14] and 1.44 (95%CI: 1.16-1.80)^[23]. Kang *et al*^[22] reported sarcopenia as an independent predictor of increased intima-media thickness with an OR of 2.26 (95%CI: 1.26-4.04)². Furthermore, Kang *et al*^[22] and Zhu *et al*^[29] reported that sarcopenia in NAFLD was associated with a higher risk of carotid plaque, with aORs of 2.74 (95%CI: 1.30-5.78) and 2.22 (95%CI: 1.23-4.02), respectively. However, Cho *et al*^[30] reported higher odds

of carotid plaque development with sarcopenia in those without carotid plaque at baseline (OR: 2.02, 95%CI: 1.32-3.08). Finally, Zhou *et al*^[34] reported that NAFLD patients with sarcopenia had a higher risk of insulin resistance in both men and women (OR: 2.14, 95%CI: 1.16-3.97; OR: 4.26, 95%CI: 1.29-14.02).

DISCUSSION

An increasing number of studies have indicated the association between sarcopenia and NAFLD. However, the exact prevalence of sarcopenia in the NAFLD population remains unclear. This systematic review is the first to summarize the current evidence on the prevalence of sarcopenia in NAFLD patients. Out of the 24 studies reporting on prevalence, 14, 4, and 2 reported prevalence rates of 10%-30%, 30%-50%, and more than 50%, respectively, whereas only four studies demonstrated a prevalence rate of less than 10%. This finding indicates that a considerable number of patients with NAFLD develop sarcopenia. In addition, sarcopenia was associated with an increased risk of advanced fibrosis in NAFLD, with an OR of 1.97 (95%CI: 1.44-2.70). Furthermore, sarcopenia in patients with NAFLD was associated with increased risks of NASH, insulin resistance, carotid plaque, and mortality.

Our systematic review highlighted a considerable variation in the reported prevalence of sarcopenia among patients with NAFLD. This variation is attributed to the diagnostic modality used-from 1.6% using MRI to 63.0% with DEXA. This discrepancy is amplified by the use of different cutoff values and indices within the same diagnostic modality, such as the normalization of ASM to BMI, weight, or height squared. This substantial variability in sarcopenia prevalence emphasizes the need for standardized diagnostic criteria and measurement techniques for sarcopenia in NAFLD patients. The European Working Group on Sarcopenia in Older People has proposed criteria and cutoffs for the three essential components of sarcopenia: muscle mass, muscle strength, and physical performance^[2]. The choice of diagnostic modality and cutoff criteria markedly affects the reported prevalence rates, highlighting the necessity for consensus guidelines to ensure consistency across studies and populations. The variation in prevalence across different studies is primarily influenced by the distribution of muscle mass index in the population

and the absolute values of the cutoff points. By contrast, variations in cutoff points for gait speed and grip strength appear to have a weak impact on the prevalence rates of sarcopenia^[39].

The results of this systematic review revealed a significant relationship between sarcopenia and an increased risk of advanced fibrosis in NAFLD patients, despite noticeable heterogeneity across the included studies. Upon examining the various modalities used for assessing fibrosis (such as biopsy, NFS/BARD scores, TE, and FIB-4 scores), a consistent association with sarcopenia was observed for all modalities except for the FIB-4 score. The absence of an association with the FIB-4 score indicates the necessity of selecting the appropriate fibrosis assessment method when exploring the relationship between sarcopenia and advanced fibrosis in NAFLD. Additionally, a recent study examined the effectiveness of noninvasive tests for estimating fibrosis, particularly in Asian populations. A recent multicentric study highlighted that only TE and TE-based combination tests accurately predicted liver fibrosis, whereas the internationally accepted thresholds for other NITs exhibited high false-negative rates^[40].

Our systematic review sheds light on the extensive range of outcomes associated with sarcopenia in NAFLD patients. Key findings included an increased risk of NASH and a higher incidence of NASH in those with sarcopenia. Additionally, our meta-analysis revealed the predictive value of sarcopenia for mortality in NAFLD, as demonstrated by two studies with aHR of 1.78 (95%CI: 1.16-2.73) and 1.44 (95%CI: 1.16-1.80^[14,23]). Sarcopenia in NAFLD was also associated with cardiovascular risk factors, such as increased intima-media thickness and a higher likelihood of carotid plaque formation. Moreover, sarcopenia was associated with a higher prevalence of insulin resistance, a key player in NAFLD pathogenesis. The relationship between sarcopenia and cardiovascular risks is particularly significant, considering the established relationship between NAFLD and adverse cardiovascular events^[41]. This association not only highlights the multifaceted impact of sarcopenia in NAFLD but also opens avenues for future research aimed at reducing cardiovascular morbidity and mortality in this patient population.

The strengths of our systematic review include acknowledging the significant heterogeneity in sarcopenia prevalence reports among NAFLD patients and emphasizing the necessity for standardized guidelines in this area. In addition, we examined non-liver-related events in NAFLD patients and their correlation with sarcopenia. Variability in sarcopenia and liver fibrosis assessment methods contributes to the diverse results observed in our review. The inclusion of studies with varied designs and the demographic differences among patient populations could also affect the observed prevalence of sarcopenia and its association with NAFLD outcomes. Although our review explores the association of sarcopenia with mortality and cardiovascular risks, some specific outcomes, such as carotid plaque risk and progression, were only addressed in a limited number of studies, affecting the conclusiveness of these findings. For instance, Zhang et al. noted a higher sarcopenia prevalence in lean versus non-lean NAFLD patients^[28], a detail we could not further analyze due to data limitations. Moreover, the review's focus on studies primarily from Asian populations, especially South Korea, may limit the generalizability of the findings to Western populations.

CONCLUSION

This systematic review highlights the multifaceted impact of sarcopenia on patients with NAFLD, extending beyond liver-related issues to include cardiovascular risks, mortality, and metabolic complications. The observed variations in prevalence and associations indicate the urgent need for standardized diagnostic criteria and measurement techniques. Our review offers crucial insights into the clinical implications of sarcopenia within the NAFLD context, potentially guiding future research and clinical practice.

REFERENCES

- 1 **Lomba R, Friedman SL, Shulman GI.** Mechanisms and disease consequences of nonalcoholic fatty liver disease. *Cell* 2021; **184**: 2537-2564 [PMID: 33989548 DOI: 10.1016/j.cell.2021.04.015]

- 2 **Cruz-Jentoft AJ**, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, Cooper C, Landi F, Rolland Y, Sayer AA, Schneider SM, Sieber CC, Topinkova E, Vandewoude M, Visser M, Zamboni M; Writing Group for the European Working Group on Sarcopenia in Older People 2 (EWGSOP2), and the Extended Group for EWGSOP2. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing* 2019; **48**: 16-31 [PMID: 30312372 DOI: 10.1093/ageing/afy169]
- 3 **Donini LM**, Busetto L, Bischoff SC, Cederholm T, Ballesteros-Pomar MD, Batsis JA, Bauer JM, Boirie Y, Cruz-Jentoft AJ, Dicker D, Frara S, Frühbeck G, Genton L, Gepner Y, Giustina A, Gonzalez MC, Han HS, Heymsfield SB, Higashiguchi T, Laviano A, Lenzi A, Nyulasi I, Parrinello E, Poggiogalle E, Prado CM, Salvador J, Rolland Y, Santini F, Serlie MJ, Shi H, Sieber CC, Siervo M, Vettor R, Villareal DT, Volkert D, Yu J, Zamboni M, Barazzoni R. Definition and Diagnostic Criteria for Sarcopenic Obesity: ESPEN and EASO Consensus Statement. *Obes Facts* 2022; **15**: 321-335 [PMID: 35196654 DOI: 10.1159/000521241]
- 4 **Dhaliwal A**, Armstrong MJ. Sarcopenia in cirrhosis: A practical overview. *Clin Med (Lond)* 2020; **20**: 489-492 [PMID: 32934043 DOI: 10.7861/clinmed.2020-0089]
- 5 **Li AA**, Kim D, Ahmed A. Association of Sarcopenia and NAFLD: An Overview. *Clin Liver Dis (Hoboken)* 2020; **16**: 73-76 [PMID: 32922754 DOI: 10.1002/cld.900]
- 6 **Bhanji RA**, Narayanan P, Allen AM, Malhi H, Watt KD. Sarcopenia in hiding: The risk and consequence of underestimating muscle dysfunction in nonalcoholic steatohepatitis. *Hepatology* 2017; **66**: 2055-2065 [PMID: 28777879 DOI: 10.1002/hep.29420]
- 7 **Wijarnpreecha K**, Panjawan P, Thongprayoon C, Jaruvongvanich V, Ungprasert P. Sarcopenia and risk of nonalcoholic fatty liver disease: A meta-analysis. *Saudi J Gastroenterol* 2018; **24**: 12-17 [PMID: 29451179 DOI: 10.4103/sjg.SJG_237_17]
- 8 **Page MJ**, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D.

The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; **372**: n71 [PMID: 33782057 DOI: 10.1136/bmj.n71]

9 **Stang A.** Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 2010; **25**: 603-605 [PMID: 20652370 DOI: 10.1007/s10654-010-9491-z]

10 **Lee YH, Kim SU, Song K, Park JY, Kim DY, Ahn SH, Lee BW, Kang ES, Cha BS, Han KH.** Sarcopenia is associated with significant liver fibrosis independently of obesity and insulin resistance in nonalcoholic fatty liver disease: Nationwide surveys (KNHANES 2008-2011). *Hepatology* 2016; **63**: 776-786 [PMID: 26638128 DOI: 10.1002/hep.28376]

11 **Koo BK, Kim D, Joo SK, Kim JH, Chang MS, Kim BG, Lee KL, Kim W.** Sarcopenia is an independent risk factor for non-alcoholic steatohepatitis and significant fibrosis. *J Hepatol* 2017; **66**: 123-131 [PMID: 27599824 DOI: 10.1016/j.jhep.2016.08.019]

12 **Petta S, Ciminnisi S, Di Marco V, Cabibi D, Cammà C, Licata A, Marchesini G, Craxi A.** Sarcopenia is associated with severe liver fibrosis in patients with non-alcoholic fatty liver disease. *Aliment Pharmacol Ther* 2017; **45**: 510-518 [PMID: 28028821 DOI: 10.1111/apt.13889]

13 **Hsieh YC, Joo SK, Koo BK, Lin HC, Kim W.** Muscle alterations are independently associated with significant fibrosis in patients with nonalcoholic fatty liver disease. *Liver Int* 2021; **41**: 494-504 [PMID: 33164304 DOI: 10.1111/liv.14719]

14 **Golabi P, Gerber L, Paik JM, Deshpande R, de Avila L, Younossi ZM.** Contribution of sarcopenia and physical inactivity to mortality in people with non-alcoholic fatty liver disease. *JHEP Rep* 2020; **2**: 100171 [PMID: 32964202 DOI: 10.1016/j.jhepr.2020.100171]

15 **Kang S, Moon MK, Kim W, Koo BK.** Association between muscle strength and advanced fibrosis in non-alcoholic fatty liver disease: a Korean nationwide survey. *J Cachexia Sarcopenia Muscle* 2020; **11**: 1232-1241 [PMID: 32638541 DOI: 10.1002/jcsm.12598]

16 **Choe HJ, Lee H, Lee D, Kwak SH, Koo BK.** Different effects of low muscle mass on the risk of non-alcoholic fatty liver disease and hepatic fibrosis in a prospective

- cohort. *J Cachexia Sarcopenia Muscle* 2023; **14**: 260-269 [PMID: 36403577 DOI: 10.1002/jcsm.13125]
- 17 **Kang MK**, Park JG, Lee HJ, Kim MC. Association of low skeletal muscle mass with advanced liver fibrosis in patients with non-alcoholic fatty liver disease. *J Gastroenterol Hepatol* 2019; **34**: 1633-1640 [PMID: 30667551 DOI: 10.1111/jgh.14607]
- 18 **Wijarnpreecha K**, Kim D, Raymond P, Scribani M, Ahmed A. Associations between sarcopenia and nonalcoholic fatty liver disease and advanced fibrosis in the USA. *Eur J Gastroenterol Hepatol* 2019; **31**: 1121-1128 [PMID: 30888971 DOI: 10.1097/MEG.0000000000001397]
- 19 **Gan D**, Wang L, Jia M, Ru Y, Ma Y, Zheng W, Zhao X, Yang F, Wang T, Mu Y, Zhu S. Low muscle mass and low muscle strength associate with nonalcoholic fatty liver disease. *Clin Nutr* 2020; **39**: 1124-1130 [PMID: 31053512 DOI: 10.1016/j.clnu.2019.04.023]
- 20 **Park H**, Jun DW, Park HK, Park KY, Kim M, Hwang HS. A Critical Appraisal of the Definition of Sarcopenia in Patients with Non-Alcoholic Fatty Liver Disease: Pitfall of Adjusted Muscle Mass by Body Weight. *Life (Basel)* 2020; **10** [PMID: 32977654 DOI: 10.3390/life10100218]
- 21 **Seo DH**, Lee YH, Park SW, Choi YJ, Huh BW, Lee E, Huh KB, Kim SH, Cha BS. Sarcopenia is associated with non-alcoholic fatty liver disease in men with type 2 diabetes. *Diabetes Metab* 2020; **46**: 362-369 [PMID: 31689496 DOI: 10.1016/j.diabet.2019.10.004]
- 22 **Kang MK**, Park JG. Low Skeletal Muscle Mass Is a Risk Factor for Subclinical Atherosclerosis in Patients with Nonalcoholic Fatty Liver Disease. *Diagnostics (Basel)* 2021; **11** [PMID: 34068776 DOI: 10.3390/diagnostics11050854]
- 23 **Kim D**, Wijarnpreecha K, Sandhu KK, Cholankeril G, Ahmed A. Sarcopenia in nonalcoholic fatty liver disease and all-cause and cause-specific mortality in the United States. *Liver Int* 2021; **41**: 1832-1840 [PMID: 33641244 DOI: 10.1111/liv.14852]
- 24 **Lee HJ**, Chang JS, Ahn JH, Kim MY, Park KS, Ahn YS, Koh SB. Association Between Low Muscle Mass and Non-alcoholic Fatty Liver Disease Diagnosed Using Ultrasonography, Magnetic Resonance Imaging Derived Proton Density Fat

Fraction, and Comprehensive NAFLD Score in Korea. *J Prev Med Public Health* 2021; **54**: 412-421 [PMID: 34875824 DOI: 10.3961/jpmph.21.387]

25 **Lee JH**, Lee HS, Lee BK, Kwon YJ, Lee JW. Relationship between Muscle Mass and Non-Alcoholic Fatty Liver Disease. *Biology (Basel)* 2021; **10** [PMID: 33562473 DOI: 10.3390/biology10020122]

26 **Wang YM**, Zhu KF, Zhou WJ, Zhang Q, Deng DF, Yang YC, Lu WW, Xu J, Yang YM. Sarcopenia is associated with the presence of nonalcoholic fatty liver disease in Zhejiang Province, China: a cross-sectional observational study. *BMC Geriatr* 2021; **21**: 55 [PMID: 33446095 DOI: 10.1186/s12877-020-01910-3]

27 **Almeida NS**, Rocha R, de Souza CA, da Cruz ACS, Ribeiro BDR, Vieira LV, Daltro C, Silva R, Sarno M, Cotrim HP. Prevalence of sarcopenia using different methods in patients with non-alcoholic fatty liver disease. *World J Hepatol* 2022; **14**: 1643-1651 [PMID: 36157861 DOI: 10.4254/wjh.v14.i8.1643]

28 **Zhang X**, He Z, Si Q, Hu X, Yang L, Gu X, Du L, Wang L, Pan L, Li Y, Li J, Yang B, Gu X. The Association of Sarcopenia and Visceral Obesity with Lean Nonalcoholic Fatty Liver Disease in Chinese Patients with Type 2 Diabetes Mellitus. *J Diabetes Res* 2022; **2022**: 2229139 [PMID: 36387941 DOI: 10.1155/2022/2229139]

29 **Zhu X**, Huang Q, Ma S, Chen L, Wu Q, Wu L, Ma H, Li X, Li Q, Aleteng Q, Hu Y, He W, Gao J, Lin H, Tang H, Gao X, Xia M. Presence of sarcopenia identifies a special group of lean NAFLD in middle-aged and older people. *Hepatol Int* 2023; **17**: 313-325 [PMID: 36327053 DOI: 10.1007/s12072-022-10439-z]

30 **Cho Y**, Park HS, Huh BW, Lee YH, Seo SH, Seo DH, Ahn SH, Hong S, Kim SH. Non-Alcoholic Fatty Liver Disease with Sarcopenia and Carotid Plaque Progression Risk in Patients with Type 2 Diabetes Mellitus. *Diabetes Metab J* 2023; **47**: 232-241 [PMID: 36653888 DOI: 10.4093/dmj.2021.0355]

31 **Chun HS**, Lee M, Lee HA, Lee S, Kim S, Jung YJ, Lee C, Kim H, Lee HA, Kim HY, Yoo K, Kim TH, Ahn SH, Kim SU. Risk Stratification for Sarcopenic Obesity in Subjects With Nonalcoholic Fatty Liver Disease. *Clin Gastroenterol Hepatol* 2023; **21**: 2298-2307.e18 [PMID: 36462755 DOI: 10.1016/j.cgh.2022.11.031]

32 **Lu Y**, Xia Q, Wu L, Xie Z. Gender difference in association between low muscle mass and risk of non-alcoholic fatty liver disease among Chinese adults with visceral

- obesity. *Front Nutr* 2023; **10**: 1026054 [PMID: 36713086 DOI: 10.3389/fnut.2023.1026054]
- 33 **Linge J**, Ekstedt M, Dahlqvist Leinhard O. Adverse muscle composition is linked to poor functional performance and metabolic comorbidities in NAFLD. *JHEP Rep* 2021; **3**: 100197 [PMID: 33598647 DOI: 10.1016/j.jhepr.2020.100197]
- 34 **Zhou T**, Ye J, Lin Y, Wang W, Feng S, Zhuo S, Zhong B. Impact of skeletal muscle mass evaluating methods on severity of metabolic associated fatty liver disease in non-elderly adults. *Br J Nutr* 2023; **130**: 1373-1384 [PMID: 36896599 DOI: 10.1017/S0007114523000399]
- 35 **Guo W**, Zhao X, Miao M, Liang X, Li X, Qin P, Lu J, Zhu W, Wu J, Zhu C, Xu N, Zhang Q. Association Between Skeletal Muscle Mass and Severity of Steatosis and Fibrosis in Non-alcoholic Fatty Liver Disease. *Front Nutr* 2022; **9**: 883015 [PMID: 35558748 DOI: 10.3389/fnut.2022.883015]
- 36 **Seo JY**, Cho EJ, Kim MJ, Kwak MS, Yang JI, Chung SJ, Yim JY, Yoon JW, Chung GE. The relationship between metabolic dysfunction-associated fatty liver disease and low muscle mass in an asymptomatic Korean population. *J Cachexia Sarcopenia Muscle* 2022; **13**: 2953-2960 [PMID: 36222309 DOI: 10.1002/jcsm.13099]
- 37 **Song W**, Yoo SH, Jang J, Baik SJ, Lee BK, Lee HW, Park JS. Association between Sarcopenic Obesity Status and Nonalcoholic Fatty Liver Disease and Fibrosis. *Gut Liver* 2023; **17**: 130-138 [PMID: 36472070 DOI: 10.5009/gnl220041]
- 38 **Harring M**, Golabi P, Paik JM, Shah D, Racila A, Cable R, Srishord M, Younossi ZM. Sarcopenia Among Patients With Nonalcoholic Fatty Liver Disease (NAFLD) Is Associated With Advanced Fibrosis. *Clin Gastroenterol Hepatol* 2023; **21**: 2876-2888.e5 [PMID: 36848980 DOI: 10.1016/j.cgh.2023.02.013]
- 39 **Masanés F**, Rojano I, Luque X, Salvà A, Serra-Rexach JA, Artaza I, Formiga F, Cuesta F, López Soto A, Ruiz D, Cruz-Jentoft AJ. Cut-off Points for Muscle Mass - Not Grip Strength or Gait Speed - Determine Variations in Sarcopenia Prevalence. *J Nutr Health Aging* 2017; **21**: 825-829 [PMID: 28717813 DOI: 10.1007/s12603-016-0844-5]
- 40 **Arora U**, Goyal RM, Teh KKJ, Pei Y, Goh GBB, Lin S, Kumar R, Biswas S, Vaishnav M, Swaroop S, Pathak P, Sheikh S, Bharadiya V, Elhence A, Gamanagatti S,

Yadav R, Das P, Aggarwal S, Choudhary N, Anirvan P, Singh SP, De A, Duseja A, Shalimar. Poor Performance of Non-invasive Tests for Advanced Fibrosis in Nonalcoholic Fatty Liver Disease: A Multicentric Asian Study. *Dig Dis Sci* 2023; **68**: 4485-4498 [PMID: 37733130 DOI: 10.1007/s10620-023-08085-y]

41 **Wu S**, Wu F, Ding Y, Hou J, Bi J, Zhang Z. Association of non-alcoholic fatty liver disease with major adverse cardiovascular events: A systematic review and meta-

Figure Legends

Figure 1 PRISMA flowchart for study identification, selection, and inclusion process. NAFLD: Nonalcoholic fatty liver disease.

Figure 2 Forest plot showing the odds of advanced fibrosis with sarcopenia in patients with non-alcoholic fatty liver disease with subgroup analysis based on the method of fibrosis assessment. OR: Odds ratio; 95%CI: 95% confidence interval; FIB-4: Fibrosis-4; NFS/BARD: Non-alcoholic fatty liver disease fibrosis scores/body mass index, the ratio of aspartate aminotransferase to alanine aminotransferase, diabetes; DL:

Table 1 Baseline characteristics of the included studies showing the prevalence and outcome of sarcopenia in patients with non-alcoholic fatty liver disease

Ref.	Country, study design	Population and size	Age, in years, male sex, in %	Comorbidities	Definition of NAFLD	Definition and prevalence of sarcopenia	Outcome	Study quality
Lee <i>et al</i> ^[10] , 2016	South Korea, retrospective	Korean National Health and Nutrition Examination Surveys 2008-2011, <i>n</i> = 2761	55.8 ± 14.3, 45%	BMI: 25.8 ± 3.1; MS: 81%; DM: 30%	NAFLD liver fat score	DEXA was used for the calculation of SI = ASM/BMI. Sarcopenia was defined using a cut-off point of SI < 0.789 in men and < 0.521 in women; <i>n</i> = 337 (12.2%)	Significant fibrosis was defined as FIB-4 ≥ 2.67. After adjusting for all covariates, a higher value of SI was associated with a lower risk of significant fibrosis with aOR: 0.67	Fair

										(95%CI: 0.49-0.91)	
Koo et al ^[11] , 2017	South Korea, prospective	Boramae NAFLD registry, n = 240	53.3 ± 14.3, 48.7%	BMI: 27.4 ± 3.5; 39.6%; 40.4%; smoking: 22.5%	DM: macrovesicular steatosis on liver biopsy	HTN: 5%	BIA was used to calculate ASM, which was divided by weight = ASM%. ASM% < 29.0 in men or < 22.9 in women was considered as sarcopenia. n = 64 (26.7%)	Among patients with NAFLD, sarcopenia was associated with a higher risk of NASH (aOR: 2.59; 95%CI: 1.22-5.48).	Good		
							(21/117 in NAFLD and 43/123 in NASH)	Sarcopenia was also associated with the presence of significant fibrosis (F2-F4) on liver			

										biopsy (aOR: 2.21; 95%CI: 1.10-4.44)	
Petta <i>et al</i> ^[12] , Italy, 2017	prospective	Consecutive patients with NAFLD at a single center, <i>n</i> = 225	48.3 ± 13.4, 62.7%	BMI: 30.3 ± 5.2; 45.3%; HTN: 32.9%; obesity: 71.1%	≥ 5%	DM: macrovesicular steatosis on liver biopsy	BIA was used to calculate ASM, which was divided by weight × 100 = SMI. Sarcopenia was defined as fibrosis (F3- an SMI ≤ 37 in males and ≤ 28 in females. <i>n</i> = 98 (43.6%)	Sarcopenia was also associated with the presence of advanced fibrosis (F3- an SMI ≤ 37 in males and ≤ 28 in biopsy (aOR: 2.36; 95%CI: 1.16-4.77). The prevalence of NASH was higher in the presence of sarcopenia (88.7% <i>vs</i> 76.3% in nonsarcopenic	Good		

Kang <i>et al</i> ^[17] , 2019	South Korea, retrospective	Adults undergoing comprehensive health screening at a single center from 2010-2017, <i>n</i> = 10711	47.9 ± 11.6, 52.8%	BMI: 23.9 ± 2.9; MS: 12.5%; DM: 5.9%; HTN: 11.6%; obesity: 34.1%	Abdominal ultrasound ¹	BIA was used to calculate ASM, which was divided by weight = ASM/BW%; ASM/BW% < 29.0 in men or < 22.9 in women was considered as sarcopenia; <i>n</i> = 615 (5.7%)	cases, <i>P</i> = 0.01)	Advanced fibrosis was defined as NFS ≥ 0.676 and FIB-4 ≥ 2.670. Sarcopenia was also associated with the presence of advanced fibrosis (F3-F4) as defined by NFS with aOR: 2.68 (95%CI: 1.28-5.59), but not using FIB-4 (aOR: 1.58,
							Fair	

								95%CI: 0.87-2.85)	
Wijarnpreecha <i>et al</i> ^[18] , 2019	United States, retrospective	Analysis of the third National Health and Nutrition Examination Survey (NHANES), conducted from 1988 to 1994, <i>n</i> = 4188	45.4 ± 0.42, 50.4%	BMI: 28.9 ± 0.22; HTN: 31.6%; DM: 7.5%	Abdominal ultrasound ¹	BIA was used to calculate ASM, which was divided by weight × 100 = SMI. Sarcopenia was defined as an SMI ≤ 37 in males and ≤ 28 in females; <i>n</i> = 2023 (48.3%)	Advanced fibrosis was defined as NFS ≥ 0.676; sarcopenia was significantly associated with advanced fibrosis (aOR: 2.39, 95%CI: 1.50-3.84)	Fair	
Gan <i>et al</i> ^[19] , 2020	China, prospective	Lanxi cohort, a community-based prospective cohort with a focus on obesity-related	55.2 ± 11.5, 32.9%	BMI: 25.9 ± 2.9; MS: 59.5%; DM: 12.9%; HTN: 48.1%	Abdominal ultrasound ¹	DEXA was used for the calculation of SMI = total appendicular lean mass (ALM)/weight.	-	Fair	

diseases, $n =$
1088

The cut-off
points for
sarcopenia were
28.64% for men
and 24.12% for
women; $n = 246$
(22.6%)

Golabi <i>et al</i> ^[14] , 2020	United State, retrospective	Analysis of the National Health and Nutrition Examination Survey (NHANES), from 1999 to 2004, $n = 1351$	50.7 ± 0.7^2 , 60.0%	BMI: $32.5 \pm$ 0.32; 60.6%; 68.4%; 63.9%; 20.7%	obesity: index (FLI) $\geq$ 30 based on age, race/ethnicity, waist circumference, GGT, activity, fasting insulin, and fasting glucose	Fatty liver DEXA was used to calculate SI = ASM/BMI. Sarcopenia was defined using a cut-off point of SI < 0.789 in men and < 0.521 in women. $n = 239$ (17.7%)	Sarcopenia was an independent predictor of mortality in NAFLD with aHR 1.78 (95%CI: 1.16- 2.73)	Fair
---	--------------------------------	---	-----------------------------	---	--	--	--	------

Hsieh <i>et al</i> ^[13] , 2021	Taiwan, prospective	Boramae NAFLD cohort, <i>n</i> = 521	52.0 ± 15.0, 50.9%	BMI: 27.8 ± 3.8; 39.3%; 42.4%	≥ 5% DM: macrovesicular HTN: steatosis on liver biopsy	Cross-sectional CT images at L3 was used to calculate SMI; Sarcopenia defined by L3- SMI < 50 cm ² /m ² for men and < 39 cm ² /m ² for women. <i>n</i> = 122 (23.4%)	Sarcopenia was also associated with the presence of significant fibrosis (F2- F4) on liver biopsy (aOR: 1.72; 95% CI: 1.05-2.84)	Good
Kang <i>et al</i> ^[15] , 2020	South Korea, retrospective	Korean National Health and Nutrition Examination Surveys 2014- 2016 with age 35-65 yr, <i>n</i> = 2092	45.6 ± 0.22, 42.4%	BMI: 23.8 ± 0.02; 10.7%; 24.1%; obesity: 33.6%	HIS was calculated based on ALT, AST, BMI, DM, sex, NAFLD defined by HIS > 36	Hand grip strength was calculated using a dynamometer, and sarcopenia was defined for individuals in the 1 st quartile (Q1) of muscle strength	Advanced fibrosis was defined as either a FIB-4 score ≥ 1.30 or a BARD score ≥ 2.00. Sarcopenia was also associated with the	Fair

						presence of advanced fibrosis as defined by BARD with aOR: 1.68 (95%CI: 1.07-2.62), but not using FIB-4 (aOR: 1.35, 95%CI: 0.75-2.45)	
Park <i>et al</i> ^[20] , 2020	South Korea, retrospective	Patients attending annual health examination at a single center, <i>n</i> = 747	48.9 ± 10.8, BMI: 24.9 ± 3.1	Abdominal ultrasound ¹	BIA was used to calculate ASM, which was divided by weight × 100 = SMI. ASM/BW% < 29.1 in men or < 23.0 in women was considered as sarcopenia. <i>n</i>	-	Fair

= 66 (8.8%)

Seo <i>et al</i> ^[21] , 2020	South Korea, retrospective	Seoul Metabolic Syndrome Cohort, <i>n</i> = 1278	55.8 10.8, 53.6%	± BMI: 26.5 ± 3.3; DM: 100%	Abdominal ultrasound ¹	BIA was used to - calculate ASM, which was divided by weight = ASM/BW%. ASM/BW% < 29.0 in men or < 22.9 in women was considered as sarcopenia. <i>n</i> = 528 (41.3%)	Fair
Kang <i>et al</i> ^[22] , 2021	South Korea, retrospective	Patients undergoing carotid ultrasound at a single center, <i>n</i>	49.1 10.0, 86.1%	± BMI: 26.4 ± 2.6; DM: 15.2%; obesity: 67%; HTN: 29.1%; MS:	Abdominal ultrasound ¹	BIA was used to Sarcopenia calculate SI = was an ASM/BMI. independent Sarcopenia was predictor of defined using a increased	Fair

			= 683		43.6%		cut-off point of SI	intima-media	
							< 0.789 in men	thickness (OR:	
							and < 0.521 in	2.26, (95%CI:	
							women. <i>n</i> = 75	1.26-4.04) and	
							(11.0%)	carotid plaque	
								(OR: 2.74,	
								95%CI: 1.30-	
								5.78)	
Kim <i>et al</i> ^[23] , 2021	United States, retrospective	Analysis of the third National Health and Nutrition Examination Survey (NHANES), conducted from 1988 to 1994, <i>n</i> = 3773	45.5 ± 0.45 ² , 50.5%	± BMI: 29.0 ± 0.23 ² ; 12.1%; HTN: 30.9%	Abdominal ultrasound ¹	BIA was used to calculate ASM, which was divided by weight × 100 = SMI. Sarcopenia was defined as an SMI ≤ 37 in males and ≤ 28 in females. <i>n</i> = 1822 (48.3%)	Sarcopenia was an independent predictor of mortality in NAFLD with aHR 1.44 (95%CI: 1.16-1.80)	Fair	
Lee <i>et al</i> ^[24] , 2021	South Korea, retrospective	Gangnam Severance Hospital	51.2 ± 11.5, 65.5%	± BMI: 26.1 ± 3.5	Abdominal ultrasound ¹	<i>n</i> = 1288 (30.9%)	-	Poor	

		Check-up (GSHC) dataset from 2016 to 2019, $n = 4168$							
Lee <i>et al</i> ^[25] , 2021	South Korea, retrospective	Korean Genome and Epidemiology Study on Atherosclerosis Risk of Rural Areas in the Korean General Population data, $n = 320$	65.7 ± 7.6 , 63.6%	BMI: $26.9 \pm$ 2.9; 67.9%; 60.5%	Abdominal ultrasound ¹ DM: HTN:	57 (39.6%), 107 (59.8%), and 148 (63.0%) participants had low muscle mass adjusted for height, BMI, and body weight in the NAFLD not group, adjusted respectively height and BMI	Appendicular muscle mass adjusted for body weight only was associated with hepatic fibrosis but not when adjusted for height and BMI	Fair	
Linge <i>et al</i> ^[33] , 2021	United Kingdom, retrospective	Participants of United Kingdom Biobank study, aged 40-69 yr at recruitment	62.9 ± 7.4 , 53.5%	BMI: 30.1 ± 4.8	MRI liver PDFF > 5%	Sarcopenia, defined as low hand grip strength [$< 16/27$ kg (females/males)]	-	Fair	

			in 2006-2010, n = 1204			and low muscle quantity [MRI threshold of 3.0 and 3.6 L/m ² for thigh FFMV/height ² (females/males)]. n = 19 (1.6%)		
Wang <i>et al</i> ^[26] , 2021	China, prospective	Patients attending annual health examination at a single center in 2019, n = 154	67.8 ± 9.3, 19.5%	BMI: 24.9 ± 2.9	Abdominal ultrasound ¹	Sarcopenia, defined as low hand grip strength (< 18 kg in women and < 26 kg in men), a gait speed < 0.8 m/s, and DEXA- based ASM/height ² < 5.4 in women and < 7.0 kg/m ² in men. n = 25 (16.2%)	-	Fair

Almeida <i>et al</i> ^[27] , 2022	Brazil, prospective	Consecutive patients with NAFLD at a single center, <i>n</i> = 57	52.7 ± 11.3, 24.6%	-	Abdominal ultrasound ¹	Probable sarcopenia, defined as low hand grip strength [$< 16/27$ kg (females/males)]. <i>n</i> = 15 (26.3%)	-	Fair
Guo <i>et al</i> ^[35] , 2022	China, prospective	Patients undergoing health checkup at a single center from 2020-2021, <i>n</i> = 1830	47.4 ± 10.5, 80.2%	BMI: 27.1 ± 3.0	Transient elastography with fat attenuation parameter > 240 dB/m	BIA was used to calculate ASM, which was divided by height $\times 100$ = SMI. SMI gradually decreased in a stepwise manner as the severity of hepatic steatosis increased	LSM values > 7.3 kPa were classified as having liver fibrosis. Participants in the tertile 1 of SMI had significantly higher odds of liver fibrosis (aOR: 3.7, 95%CI: 2.6-5.3) compared	Good

								to tertile 3	
Seo <i>et al</i> ^[36] , 2022	South Korea, retrospective	Patients undergoing health checkup at a single center from 2017-2019, <i>n</i> = 3198	54.2 ± 9.6, 89.8%	BMI: 26.2 ± 2.9; 40.2%; 20.1%	Transient elastography with controlled attenuation parameter > 248 dB/m	BIA was used to calculate ASM, which was divided by weight × 100 = SMI. ASM/BW% < 29.1 in men or < 23.0 in women was considered as sarcopenia. <i>n</i> = 517 (16.2%)	-	Poor	
Song <i>et al</i> ^[37] , 2023	South Korea, retrospective	Patients undergoing health checkup at a single center from 2007-2018, <i>n</i> = 1180	53.3 ± 10.3, 71.5%	BMI: 26.7 ± 3.67; 20.7%	Transient elastography with fat attenuation parameter > 260 dB/m	BIA was used to calculate ASM, which was divided by weight × 100 = SMI. ASM/BW% < 30.0 in men or < 26.8 in women was considered as sarcopenia	LSM values ≥ 7.5 kPa (≥ F2) were classified as having liver fibrosis. Sarcopenia was not a predictor of fibrosis in NAFLD with	Good	

							aOR: 3.80 (95% CI: 0.86-16.75)	
Zhang <i>et al</i> ^[28] , 2022	China, retrospective	T2DM patients with BMI < 25 kg/m ² were enrolled from a single center from 2017 to 2021, <i>n</i> = 1112	53.4 ± 10.7, 57.6%	BMI: 22.6; DM: 100%	Abdominal ultrasound ¹	BIA was used to calculate ASM, which was divided by weight × 100 = SMI. ASM/BW% < 32.2 in men or < 25.5 in women was considered as sarcopenia. <i>n</i> = 290 (26.1%)	-	Fair
Zhu <i>et al</i> ^[29] , 2023	China, prospective	Participants of Shanghai Changfeng Study, a community-based prospective cohort study of	62.6 ± 8.9, 33.1%	BMI: 25.7 ± 3.2	Fatty liver was diagnosed when liver fat content by ultrasound exceeded the cut-off value of 9.15%	DEXA was used to calculate SI = ASM/height ² . The cut-off SI for sarcopenia were 6.88 kg/m ² in male and 5.67 kg/m ² in female.	Significant fibrosis was defined as FIB-4 ≥ 2.67. The presence of sarcopenia was associated	Fair

multiple
chronic
diseases Jun
2009 to Dec
2012, with age
> 45 yr, $n =$
1305

$n = 260$ (19.9%) with increased
risk of carotid
plaque (aOR:
2.22; 95%CI:
1.23-4.02) and
liver fibrosis
(aOR: 2.07;
95%CI: 1.24-
3.44)

Cho *et al*^[30], South Korea, Patients with 55.0 $\pm$ 9.4, BMI: 25.7 $\pm$ Abdominal
2023 retrospective T2DM from the 46.3% 2.8; DM: 100%; ultrasound¹
Seoul HTN: 36.0%
Metabolic
Syndrome
Cohort, $n = 456$

BIA was used to Sarcopenia Good
calculate ASM, was an
which was independent
divided by predictor
weight = carotid plaque
ASM/BW%; progression
ASM/BW% < (OR: 2.02,
29.0 in men or < 95%CI: 1.32-
22.9 in women 3.08)
was considered
as sarcopenia. n
= 123 (27.0%)

Choe *et al*^[16], South Korea, Korean 51.7 $\pm$ 8.5, BMI: 27.9 $\pm$ Hepatic

- Fibrosis was Poor

2023	retrospective	Genome and Epidemiology Study (KoGES) Ansung-Ansan cohort, $n = 1442$	40.0%	2.5; 28.4%; 34.7%; 69.7%	DM: steatosis index HTN: (HSI) based on MS: ALT, AST, BMI, DM, sex. NAFLD defined by HSI > 36	defined as FIB-4 ≥ 1.3 and APRI ≥ 0.5 . In the adjusted model, low muscle mass (lowest quartile) did not contribute to progression to hepatic fibrosis (HR: 1.02, 95%CI: 0.85-1.22)
------	---------------	---	-------	-----------------------------------	---	--

Chun <i>et al</i> ^[31] , 2023	South Korea, retrospective	Patients undergoing health checkup at a three center from 2014-2020, $n = 23889$	50.0 $\pm$ 11.0, 69.5%	BMI: 25.9 $\pm$ 3.3; 14.4%; 37%; 56.9%; 47.1%	Abdominal ultrasound ¹ DM: HTN: obesity: MS:	BIA was used to - calculate ASM, which was divided by weight = ASM/BW%; ASM/BW% <	Fair
---	-------------------------------	--	---------------------------	---	--	---	------

29.0 in men or < 22.9 in women was considered as sarcopenia. $n = 3092$ (12.9%). Sarcopenia was defined using a cut-off point of $ASM/BMI = SI < 0.789$ in men and < 0.521 in women. $n = 1577$ (6.6%)

Harring <i>al</i> ^[38] , 2023	<i>et</i> United States, retrospective	Analysis of the third National Health and Nutrition Examination Survey (NHANES), from 2017 to 2018, $n = 1056$	41.9 ± 0.42 ² , 54.8%	BMI: 33.5 ± 0.37 ² ; obesity: 78.7%; DM: 18.1%; HTN: 53.9%; MS: 64.8%	Transient elastography with fat attenuation parameter > 263 dB/m	DEXA was used - to calculate $SI = ASM/BMI$. Sarcopenia was defined using a cut-off point of $SI < 0.789$ in men and < 0.512 in women. $n = 303$	Good
---	--	--	----------------------------------	--	--	---	------

Lu <i>et al</i> ^[32] , 2023	China, retrospective	Patients diagnosed with obesity during health checkup at a single center from 2020-2021, <i>n</i> = 476	51.0 13.7, 52.7%	± BMI: 27.9 ± 3.3; 100%	Abdominal ultrasound ¹	(28.7%)	BIA was used to - calculate SMI = appendicular skeletal mass/height ² . Sarcopenia defined as SMI ≤ 7.0 kg/m ² for males and ≤ 5.7 kg/m ² for females; <i>n</i> = 261	Fair
						(54.8%)		
Zhou <i>et al</i> ^[34] , 2023	China, prospective	Consecutively enrolled subjects who underwent BIA at a single center, between May 2017 and July 2022, <i>n</i> =	37.8 10.6, 58.7%	± BMI: 28.9 ± 5.1; DM: 17.6%	MRI liver PDFF > 5%	BIA was used to calculate the appendicular skeletal mass (ASM). Sarcopenia was defined as ASM/height ² or	The MAFLD patients with lower quartiles of ASM/W had a higher risk OR for insulin resistance,	Fair

ASM/weight or both in male
 ASM/BMI less and female
 than 2 SD. $n = 50$ (OR: 2.14,
 (4.4%) 95%CI: 1.16-
 3.97), and OR:
 4.26, 95%CI:
 1.29, 14.02) for
 Q4 vs Q1

¹Abdominal ultrasound showing at least two of the following three abnormal findings: (1) Diffusely increased echogenicity in liver near field ('bright liver') with greater liver echogenicity than kidney or spleen; (2) vascular blurring; and (3) poor visualization of the posterior portion of the right lobe because of deep attenuation.

²Standard error.

ALT/AST/GGT: Alanine transaminase/aspartate transaminase/gamma-glutamyltransferase; aOR: Adjusted odds ratio; 95%CI: 95% confidence interval; ASM: Appendicular skeletal muscle mass; BIA: Bioelectrical impedance analysis; BMI: Body mass index; BW: Body weight; DM/HTN/MS: Diabetes mellitus/hypertension/metabolic syndrome; DEXA: Dual X-ray absorptiometry; LSM: Liver stiffness measurement; MRI-PDFF: Magnetic resonance imaging proton density fat fraction; SMI: Skeletal muscle mass index; HIS: Hepatic steatosis index; FIB-4: Fibrosis-4; NASH: Nonalcoholic steatohepatitis; NAFLD: Nonalcoholic fatty liver disease.

ORIGINALITY REPORT

10%

SIMILARITY INDEX

PRIMARY SOURCES

1 Suprabhat Giri, Prajna Anirvan, Mansi Chaudhary, Taraprasad Tripathy et al. "Impact of nutritional status on the outcome of transjugular intrahepatic portosystemic shunt in patients with cirrhosis: A systematic review", British Journal of Radiology, 2023 88 words — 3%

Crossref

2 bmcgeriatr.biomedcentral.com 29 words — 1%

Internet

3 Yoo Jin Choo, Min Cheol Chang. "Prevalence of Sarcopenia Among the Elderly in Korea: A Meta-Analysis", Journal of Preventive Medicine and Public Health, 2021 25 words — 1%

Crossref

4 e-ce.org 23 words — 1%

Internet

5 www.mdpi.com 23 words — 1%

Internet

6

www.science.gov

Internet

23 words — 1%

7

S. Petta, S. Ciminnisi, V. Di Marco, D. Cabibi, C. Cammà, A. Licata, G. Marchesini, A. Craxì. "Sarcopenia is associated with severe liver fibrosis in patients with non-alcoholic fatty liver disease", *Alimentary Pharmacology & Therapeutics*, 2017

Crossref

21 words — 1%

8

rcastoragev2.blob.core.windows.net

Internet

18 words — 1%

9

Ji Min Choi, Hyo Eun Park, Yoo Min Han, Jooyoung Lee, Heesun Lee, Su Jin Chung, Seon Hee Lim, Jeong Yoon Yim, Goh Eun Chung. "Non-alcoholic/Metabolic-Associated Fatty Liver Disease and Helicobacter pylori Additively Increase the Risk of Arterial Stiffness", *Frontiers in Medicine*, 2022

Crossref

16 words — 1%

10

worldhealth.net

Internet

15 words — 1%

11

www.ncbi.nlm.nih.gov

Internet

15 words — 1%

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES

< 15 WORDS

EXCLUDE MATCHES

< 10 WORDS