

91430\_Auto\_Edited-check.docx

**Progress of mitochondrial and endoplasmic reticulum-associated signaling and its regulation of chronic liver disease by Chinese medicine**

Zheng Y *et al.* Endoplasmic reticulum MAM and CLD

Yang Zheng, Yi-Hui Zheng, Jia-Hui Wang, Tie-Jian Zhao, Lei Wang, Tian-Jian Liang

**Abstract**

**Abstract:** The endoplasmic reticulum (ER) is connected to mitochondria through mitochondria-associated ER membranes (MAMs). MAMs provide a framework for crosstalk between the ER and mitochondria, playing a crucial role in regulating cellular calcium balance, lipid metabolism, and cell death. Dysregulation of MAMs is involved in the development of chronic liver disease (CLD). In CLD, changes in MAMs structure and function occur due to factors such as cellular stress, inflammation, and oxidative stress, leading to abnormal interactions between mitochondria and the ER, resulting in liver cell injury, fibrosis, and impaired liver function. Traditional Chinese medicine has shown some research progress in regulating MAMs signaling and treating CLD. This paper reviews the literature on the association between mitochondria and the ER, as well as the intervention of traditional Chinese medicine in regulating CLD.

**Key Words:** Mitochondria; Endoplasmic reticulum; Mitochondria-associated ER membranes; Traditional Chinese medicine; Chronic liver disease

Zheng Y, Zheng YH, Wang JH, Zhao TJ, Wang L, Liang TJ. Progress of mitochondrial and endoplasmic reticulum-associated signaling and its regulation of chronic liver disease by Chinese medicine. *World J Hepatol* 2024; In press

**Core Tip:** Endoplasmic reticulum mitochondria-associated membranes play a very important role in the pathogenesis of chronic liver disease.

## **INTRODUCTION**

Chronic liver disease (CLD) is a global health threat and a leading cause of human mortality, encompassing non-alcoholic fatty liver disease (NAFLD), alcoholic fatty liver disease, viral hepatitis, liver fibrosis, and hepatocellular carcinoma (HCC)<sup>[1]</sup>. While early-stage CLD can be reversed, advanced liver cirrhosis and HCC severely impact patients' quality of life and even pose life-threatening risks. Mitochondria, known as the powerhouse of cells, are responsible for generating cellular energy through oxidative phosphorylation. Additionally, mitochondria play a role in calcium homeostasis, reactive oxygen species production, and cell apoptosis. Disruption of mitochondrial function is associated with the progression of CLD, including liver fibrosis, lipid degeneration, and hepatocyte injury. On the other hand, the endoplasmic reticulum (ER) is an interconnected membrane network involved in protein synthesis, folding, and transport. It also plays a crucial role in lipid metabolism, calcium signaling, and cellular stress response. Disturbance of ER homeostasis leads to the accumulation of misfolded proteins, triggering a cellular stress response known as ER stress. ER stress has emerged as a key factor in liver injury, inflammation, and the development of CLD. There is crosstalk between the ER and mitochondrial signaling pathways, with mitochondria-associated ER membranes (MAMs) connecting the ER and mitochondria in terms of structure and function. MAMs play important roles in cellular signaling and function, including calcium signaling, mitochondrial dynamics, energy metabolism, inflammation response, lipid transport, and cell apoptosis. Thus, they have a significant impact on the interplay and regulation of CLD<sup>[2]</sup>. Given the interdependence and

coordinated functions of mitochondria and the ER, understanding the potential mechanisms of mitochondria-ER signaling in CLD has become an area of in-depth research. Targeted therapeutic approaches may offer potential strategies to improve liver damage and prevent disease progression.

## **MITOCHONDRIA AND CLDS**

Mitochondria, also known as the "powerhouses" of cells, play a crucial role in various metabolic processes<sup>[3]</sup>. They serve as the site for cellular oxidative phosphorylation and adenosine triphosphate (ATP) synthesis, and are involved in a wide range of important biological functions, including energy conversion, the citric acid cycle, regulation of cellular calcium concentration, as well as being the primary source of reactive oxygen species (ROS) and a regulatory center for cell apoptosis<sup>[4]</sup>.

### ***Mitochondrial dysfunction and NAFLD***

Research suggests that NAFLD is characterized by excessive accumulation of fat in liver cells, leading to abnormal fatty acid oxidation, significant increase in mitochondrial reactive ROS, and alterations in mitochondrial membrane lipids and proteins<sup>[5,6]</sup>. Zeng *et al*<sup>[7]</sup> found that fatty acid translocase (FAT/CD36) on the mitochondrial membrane is heavily glycosylated in NAFLD, reducing its ability to transport long-chain fatty acids into the mitochondria and inhibiting fatty acid oxidation. Additionally, the lipid composition of the mitochondrial membrane changes with the continuous accumulation of lipids<sup>[7]</sup>. The increased electron transport chain (ETC) activity leads to excessive ROS production, which directly attacks biomolecules, reduces intracellular antioxidant enzyme levels, and causes oxidative damage to cells. This results in metabolic changes and ultimately leads to ETC dysfunction. The combination of increased rates of fatty acid beta-oxidation and elevated ROS levels leads to damage to the mitochondrial respiratory chain, insulin resistance, and inflammation<sup>[8]</sup>.

### ***Mitochondrial dysfunction and alcoholic liver disease***

In AFLD, various changes occur in liver mitochondria, such as increased levels of reactive ROS, decreased mitochondrial membrane potential, abnormal fatty acid beta-oxidation, and mitotic abnormalities<sup>[9]</sup>. Alcohol metabolism in the liver interferes with the oxidation of fatty acids, leading to the accumulation of triglycerides (TG). Simultaneously, alcohol metabolism generates a large amount of ROS, which causes oxidative damage to cells, resulting in metabolic changes, liver cell injury, and inflammation leading to the development of alcoholic liver disease (ALD)<sup>[10]</sup>. Furthermore, alcohol-induced mitochondrial damage exacerbates liver injury and steatosis. Mitochondria undergo structural remodeling to enhance alcohol metabolism. Studies have shown that alcohol can damage mtDNA, impair cellular energy metabolism, and increase ROS production. Additionally, alcohol induces excessive accumulation of iron in the liver. Alcohol metabolism also reduces the levels of the antioxidant GSH in mitochondria, leading to increased ROS production, lipid peroxidation in liver cells, and iron-induced cell death<sup>[11]</sup>.

#### ***Mitochondrial dysfunction and viral hepatitis***

The incidence and contagiousness of viral hepatitis are high, with hepatitis B (HBV) and hepatitis C (HCV) being common forms of chronic hepatitis. Gao *et al*<sup>[12]</sup> demonstrated through their experiments that morphological and structural changes in mitochondria can typically be detected in patients with hepatitis B<sup>[13]</sup>. HBV can induce perinuclear aggregation of mitochondria and trigger their translocation through phosphorylation of the core protein at Ser616, leading to mitochondrial division<sup>[14]</sup>. Glick *et al*<sup>[15]</sup> found that the absence of BNIP3 promotes excessive accumulation of reactive ROS, inflammatory reactions, and the formation of fatty liver in liver tissue. The characteristic of HCV-related CLD is an increase in ROS, which causes metabolic disorders, resulting in insulin resistance, hepatic steatosis, and iron accumulation in the liver. Amako *et al*<sup>[16]</sup> discovered that the HCV core protein NS5A can induce mitochondrial division, inhibit the translocation of Parkin to mitochondria, reduce autophagic activity, and maintain HCV-induced mitochondrial damage.

### ***Mitochondrial dysfunction and liver fibrosis***

Yong *et al*<sup>[17]</sup> discovered that pathological factors can cause mitochondrial damage, leading to the release of mtDNA and triggering an inflammatory response through the cyclic GMP-AMP synthase (cGAS)-stimulator of interferon genes (STING) pathway. In their experiments, abnormal mitochondrial morphology was observed, and RT-PCR revealed an increase in mtDNA content in the cytoplasm, making mtDNA highly susceptible to ROS attack<sup>[18]</sup>. Mutations in mtDNA can easily affect genes encoding important functional regions, leading to the occurrence of mitochondrial dysfunction-related diseases and inducing activation of the hepatic inflammatory signaling pathway<sup>[19]</sup>. In the liver, the expression levels of NLRP3, caspase, and IL-1 $\beta$  are significantly increased, along with the expression levels of key molecules in the cGAS-STING signaling pathway, including cGAS, STING, IRF3, and downstream NF- $\kappa$ B nuclear translocation. This also leads to the accumulation of TDP-43 in liver cells, inducing mitochondrial damage. The release of mtDNA from mitochondrial damage can activate the cGAS-STING signaling pathway, collectively triggering an inflammatory response and promoting the formation of liver fibrosis. The progression of liver fibrosis is believed to be associated with iron overload, which promotes hepatocyte iron death through excessive induction of lipid peroxidation<sup>[20]</sup>. Inhibition of hepatic stellate cell (HSC) activation and alleviation of liver fibrosis can be achieved by regulating iron death<sup>[21]</sup>.

### ***Mitochondrial dysfunction and HCC***

Mitochondrial dysfunction-mediated accumulation of reactive ROS and damage to mitochondrial DNA may contribute to the development of liver cancer. Research has shown that mitochondrial fission promotes the survival of liver cancer cells by enhancing autophagy and suppressing mitochondria-dependent cell apoptosis, and inhibitors of mitochondrial fission significantly inhibit tumor growth<sup>[18]</sup>. The characteristic feature of mitochondria in cancer cells is the excessive production of ROS,



5

which promotes cancer development by inducing genomic instability, modifying gene expression, and participating in signaling pathways. The relative increase in ROS is due to a slowdown in electron transfer in the respiratory chain, leading to an absolute increase in ROS production, as well as a decrease in antioxidant enzyme activity. The decrease in cellular antioxidant enzyme levels leads to oxidative damage in cells and signaling transduction, resulting in corresponding metabolic changes<sup>[8]</sup>. Mechanistic analysis has shown that mitochondrial fission leads to increased ROS production, mediates the activation of AKT, and synergistically interacts with the TP53 and NF- $\kappa$ B pathways to promote tumor progression<sup>[22]</sup>.

### **ENDOPLASMIC RETICULUM AND CLD**

The ER is a vital organelle involved in calcium storage, lipid metabolism, and synthesis of steroid hormones<sup>[23]</sup>. External stimuli that disrupt protein folding or lipid homeostasis can trigger ER stress (ERS), activating the unfolded protein response (UPR) signaling pathway in cells. Research has shown that ERS is closely associated with the development of various diseases, including inflammation, dysregulation of glucose and lipid metabolism, liver fibrosis, and cancer.

#### ***Endoplasmic reticulum stress and NAFLD***

ERS can contribute to the progression of NAFLD from simple steatosis to NASH, which is characterized by the development of inflammation and varying degrees of fibrosis<sup>[24]</sup>. The "second-strike theory" suggests that the liver undergoes oxidative stress, resulting in increased cellular activity and a massive release of inflammatory cytokines, causing hepatocellular inflammation and the formation of NAFLD<sup>[25]</sup>. In NAFLD, cytotoxic lipids modify the ER structure and directly activate the UPR, leading to dysregulation of protein homeostasis. Although hepatocytes can tolerate TG storage, saturated fatty acids and lysophosphatidylcholine cause lipotoxicity<sup>[26,27]</sup>. Once taken up by hepatocytes, these lipids are stored in cytoplasmic droplets<sup>[28]</sup>, and excess saturated fatty acids trigger hepatocyte ERS through various mechanisms<sup>[41]</sup>. Vander Graaff *et*

*al*<sup>[44]</sup> found that hypoxia-inducible factors stimulate the overexpression of vascular endothelial growth factor, ultimately leading to enhanced neovascularization. ERS and IR trigger NAFLD by acting on hepatocytes and causing massive intracellular lipid deposition.

#### ***Endoplasmic reticulum stress and ALD***

Homocysteine is known to enhance the production of NF- $\kappa$ B, IL-1b, IL-6, and IL-8, induce endoplasmic ERS, and promote various cellular damage processes<sup>[29]</sup>. Homocysteine can activate pathways such as GRP78, CHOP, IRE1 $\alpha$ , sterol regulatory element-binding protein, and JNK<sup>[30]</sup>. Long-term alcohol administration in animals leads to increased expression of cytochrome P450, which in turn enhances the expression of superoxide dismutase and activates nuclear factor erythroid 2-related factor 2, a key factor in the ESR response. Sun *et al*<sup>[31]</sup> found that changes in gene expression in ALD are closely related to its pathogenesis. Gene ontology enrichment analysis revealed two main functional groups: angiogenesis and stress response. Abnormal angiogenesis is also observed in the progression of ALD and exacerbates alcohol-induced liver injury. Studies have shown that excessive ethanol intake can cause chronic inflammation in liver parenchyma, leading to liver fibrosis and pathological angiogenesis.

#### ***Endoplasmic reticulum stress and viral hepatitis***

ERS plays a crucial role in the progression of viral hepatitis, particularly in cases of HBV infection. The HBV inflammatory surface antigen can activate the UPR, leading to hepatocyte apoptosis and precancerous phenotypic changes. Notably, elevated levels of GRP78, an ER chaperone protein, have been observed in HBV patients, suggesting its association with HBV infection<sup>[32]</sup>. Additionally, research indicates that HCV-related liver cancer patients exhibit upregulation of vascular endothelial growth factor (VEGF) in tissue serum, leading to abnormal angiogenesis acceleration<sup>[33]</sup>. Accumulation of misfolded proteins triggers ER stress and the expression of GRP78<sup>[34]</sup>. GRP94, which



binds to  $\text{Ca}^{2+}$ , facilitates translocation and folding of newly synthesized polypeptides, participates in oligomer assembly and degradation, and inhibits the formation of misfolded proteins. Furthermore, GRP94 acts as an antigen-presenting cell, activating the UPR and initiating cellular self-regulation<sup>[35]</sup>. XBP1 promotes the expression of molecular chaperones such as GRP78 and GRP94, enhancing the cell's ability to handle unfolded proteins and activating ER-associated degradation (ERAD) and other ER degradation pathway-related genes.

#### *Endoplasmic reticulum stress and liver fibrosis*

ERS is a cellular response to various stressors that disrupt protein folding in the endoplasmic ER. Three transmembrane proteins on the ER membrane, namely ATF6, PERK, and IRE1 $\alpha$ , play crucial roles in releasing unfolded/misfolded proteins and initiating the UPR. During ERS, BiP dissociates from these transmembrane proteins and binds to unfolded/misfolded proteins, activating downstream signaling pathways<sup>[36]</sup>. In liver cells, PERK-mediated ERS induces the upregulation of dual-specificity phosphatase 5, which reduces the phosphorylation levels of extracellular signal-regulated kinases and inhibits their activity. Additionally, PERK-mediated ERS increases the activation of caspase-3, leading to hepatocyte death and liver fibrosis<sup>[37]</sup>. Overexpression of IRE1 $\alpha$  in HSCs downregulates IRE1 $\alpha$ , XBP1, PERK, and CHOP, preventing HSC activation and reducing liver injury and fibrosis. In PDGF-induced HSCs, the activation of the PI3K/AKT pathway disrupts calcium homeostasis, leading to the accumulation of misfolded proteins and the induction of ERS, ultimately promoting HSC activation, proliferation, and liver fibrosis<sup>[38]</sup>.

#### *Endoplasmic reticulum stress and HCC*

<sup>11</sup> ATF6 is a type II ER transmembrane protein that dissociates from BiP and undergoes cleavage when ER stress occurs<sup>[39]</sup>. The cleaved form of ATF6 $\alpha$  not only activates the transcription of XBP1u, but also induces genes related to protein folding and ERAD<sup>[40]</sup>.  
<sup>1</sup> The rapid proliferation of tumor cells is accompanied by a sharp increase in protein

synthesis, inevitably leading to UPR activation, while miRNA imbalance exists in both hepatitis and HCC. Some studies have also shown that the endoribonuclease activity of IRE1 is not only responsible for cleaving XBP1, but also for cleaving and regulating miRNAs<sup>[41]</sup>. Therefore, ER stress leads to miRNA imbalance in inflammation and cancer, promoting tumor development and progression. Tumor cells can transmit ER stress and UPR signals to neighboring macrophages, upregulating UPR target genes and pro-inflammatory factors, thereby promoting the pro-inflammatory effect of tumors and regulating the tumor microenvironment<sup>[42,43]</sup>. VEGF, as an important regulator of angiogenesis, inhibits tumor growth by restricting blood vessel formation and reducing tumor blood supply when its levels decrease<sup>[44]</sup>.

#### **MITOCHONDRIAL AND ENDOPLASMIC RETICULUM ASSOCIATION SIGNALING AND CLD**

There is crosstalk between ER and mitochondrial signaling pathways, and the opening of endoplasmic reticulum InsP3/Ca ion channels affects mitochondrial Ca<sup>2+</sup> homeostasis, and disruption of mitochondrial Ca<sup>2+</sup> homeostasis leads to alterations in mitochondrial membrane potential, permeability transition pore, and other alterations in mitochondria, which are one of the most important organelles in hepatocytes, providing the majority of the cell's energy<sup>[45]</sup>. The ER is connected to the mitochondria by MAMs connected to the mitochondria, a dynamic structure that is highly sensitive to the cellular physiological environment and is mainly involved in the regulation of ERS, oxidative stress, apoptosis, cellular autophagy, changes in mitochondrial dynamics and inflammation<sup>[46]</sup>.

#### ***MAMs and NAFLD***

The molecular mediators of mitochondrial fusion are primarily Mfn1/2 (mitofusin1/2). Both Mfn1 and Mfn2 are located on the outer mitochondrial membrane, with Mfn2 also being localized on MAMs. Together, they regulate the structure and function of MAMs, inhibit the proximity of ER and mitochondria, and promote mitochondrial fusion<sup>[47]</sup>.

Reduced expression of Mfn2 has been observed in liver biopsies of NASH patients, and specific ablation of liver Mfn2 *in vivo* leads to inflammation, TG accumulation, fibrosis, and liver cancer. This suggests that Mfn2 binds to phosphatidylserine (PS) and can selectively extract PS into membrane domains, facilitating its transfer to mitochondria and the synthesis of mitochondrial phosphatidylethanolamine (PE). Therefore, the deficiency of Mfn2 in the liver reduces PS transfer and phospholipid synthesis, leading to ER stress and the development of NASH. Specifically, PS is mainly synthesized in the ER and enters mitochondria through transient membrane contacts between MAMs and the outer mitochondrial membrane. In mitochondria, PS is converted to PE, which then enters the endoplasmic reticulum to be converted to phosphatidylcholine (PC). Thus, MAMs play a crucial role in phospholipid synthesis and transport<sup>[48]</sup>.

#### ***MAMs and ALD***

Under the influence of alcohol, PERK, as a connecting protein of MAMs, mediates liver cell death through the regulation of MAMs. When PERK in MAMs is knocked out, cells exhibit abnormal ER morphology, as well as MAMs disruption and calcium imbalance<sup>[49]</sup>. After passing through the gastrointestinal tract, alcohol is absorbed into the bloodstream and enters the liver for metabolism. Within liver cells, enzymes such as alcohol dehydrogenase and aldehyde dehydrogenase convert alcohol into acetic acid, which enters the tricarboxylic acid cycle (TCA), leading to a decrease in the NAD<sup>+</sup>/NADH ratio and further mitochondrial dysfunction. Damage to the mitochondria impairs the TCA, leading to disturbances in fatty acid metabolism and excessive fat deposition in liver cells, resulting in fatty liver. Additionally, alcohol can also cause damage to ER morphology and function in liver cells through various pathways. In alcohol-induced liver tissue with severe impairment of both mitochondrial and ER function and morphology, MAMs, as a functional platform where these two organelles interact, may be more sensitive to alcohol exposure<sup>[50]</sup>.

#### ***MAMs and viral hepatitis***

Our findings demonstrate that STING can bind to mitochondrial antiviral signaling protein at MAMs, thereby enhancing the interferon response to viral infections. Additionally, HCV proteins were found to localize to MAMs, potentially leading to increased mitochondrial ROS levels through  $\text{Ca}^{2+}$  manipulation. Furthermore, certain molecules present on the structure of MAMs, such as Ero1 and p66Shc, promote ROS production. Ero1, an important molecule involved in redox homeostasis, consists of two isoforms, Ero1- $\alpha$  and Ero1- $\beta$ . The majority of Ero1- $\alpha$  is localized on MAMs, and its upregulation can result in increased ROS production. Notably, the dephosphorylation of the Ser36 site enables its transfer to MAMs, mediating ROS generation. This process facilitates the flow of  $\text{Ca}^{2+}$  from the endoplasmic ER to mitochondria through MAMs. Consequently, the accumulation of mitochondrial  $\text{Ca}^{2+}$  leads to mitochondrial depolarization and abnormalities in oxidative phosphorylation. As a result, the mitochondrial electron transport chain uncouples with respiratory complexes I and III, further augmenting ROS production. This process can be described by  $\text{ERCa}^{2+}$ , which serves as the basis for mitochondrial ROS production. Importantly, the inhibition of  $\text{ERCa}^{2+}$  channels can effectively block this process<sup>[51,52]</sup>.

### *MAMs and liver fibrosis*

MAMs can have an effect on mitochondrial function, including fusion/fission, mitophagy, and energy metabolism. MAMs play a dual role in maintaining cellular homeostasis by promoting mitochondrial division and mitophagy while also potentially causing pathological damage through excessive mitochondrial fission, calcium overload, or oxidative stress. Evidence suggests that MAMs are involved in PINK1/Parkin-mediated mitophagy, with core proteins associated with MAMs regulating their integrity and functionality<sup>[49]</sup>. *In vivo* studies have demonstrated that liver-specific ablation of Mfn2, a protein involved in MAMs, leads to inflammation, triglyceride accumulation, fibrosis, and HCC. This suggests that Mfn2 binds to PS and selectively extracts it to membrane domains, facilitating PS transfer to mitochondria and mitochondrial phosphatidylethanolamine synthesis<sup>[53]</sup>.



### *MAMs and HCC*

Research has shown that enhancing the physical contact between the endoplasmic ER and mitochondria leads to mitochondrial calcium overload and cell apoptosis. Conversely, disrupting the ER-mitochondria contact points stimulates mitochondrial oxidative respiration and ATP production<sup>[54]</sup>. Increased mitochondrial fission has been observed in liver cancer tissues, and it has been found that enhanced mitochondrial fission promotes the growth of liver cancer cells. Overexpression of Drp1 in liver cancer cells, followed by overexpression of Rab32, leads to a decrease in MAMs structure formation, a significant increase in mitochondrial calcium concentration, and subsequently promotes liver cancer cell apoptosis while inhibiting cell proliferation. Numerous studies have found that MAMs play important roles in both calcium signal transduction and the transfer of calcium from the ER to mitochondria<sup>[55]</sup>. Excessive uptake of mitochondrial calcium can result in mitochondrial calcium overload, opening of the permeability transition pore, mitochondrial swelling, rupture of the outer mitochondrial membrane, and subsequently, release of cytochrome c and cell apoptosis<sup>[54]</sup>.

### TRADITIONAL CHINESE MEDICINE TARGETS MITOCHONDRIAL ENDOPLASMIC RETICULUM MODULATION TO INTERVENE IN CLD

The research on traditional Chinese medicine (TCM) in regulating MAMs signaling and treating CLDs has shown significant progress. Certain TCM components have been proven to have a reparative effect on the structure and function of MAMs, thereby improving the progression of CLDs. Additionally, some active ingredients in TCM exhibit antioxidant, anti-inflammatory, and insulin-sensitivity regulating properties, some of which may be related to their modulation of MAMs signaling. The modulation of MAMs signaling by TCM components is an important research direction for the treatment of CLDs. Current studies have demonstrated that certain TCM components

can regulate MAMs signaling and improve liver pathological responses, providing new insights and approaches for the treatment of CLDs.

### *Silymarin*

Silymarin is a novel flavonoid compound extracted from the seeds of *Silybum marianum*<sup>[56]</sup>. This drug has been shown to protect the liver, promote liver cell regeneration, enhance liver metabolism, and reduce serum transaminase levels. It is suitable for adjunctive treatment of chronic hepatitis, cirrhosis, and other liver diseases. Animal experiments have shown that silymarin can increase the expression of superoxide dismutase (SOD) in lymphocytes of mice with CLD, thereby reducing the production of reactive ROS and protecting liver cell membranes<sup>[57]</sup>. Silymarin can also stabilize endoplasmic ER function and exert anti-inflammatory and hepatoprotective effects by reducing the levels of TNF- $\alpha$ , IL-6, IL-8, and other inflammatory cytokines<sup>[58]</sup>.

### *Tartaric acid*

Tartaric acid can inhibit lipid peroxidation and clear ROS, effectively preventing the occurrence and development of liver fibrosis induced by carbon tetrachloride (CCl<sub>4</sub>). Multiple studies have found that rhein can inhibit the chemotaxis and phagocytosis of neutrophils, as well as the activation of macrophage lipid inflammatory mediators, and suppress the activity of TNF- $\alpha$  and IL-1 $\beta$ <sup>[59,60]</sup>. Rhein has the ability to clear ROS and inhibit lipid peroxidation<sup>[61]</sup>. Research has shown that in animal models of CCl<sub>4</sub> and ethanol-induced liver fibrosis, rhein can significantly reduce malondialdehyde levels and increase SOD levels, thereby reducing the excessive generation of ROS in mitochondria.

### *Quercetin*

Quercetin is a flavonoid compound. Recent studies have found that subcellular mechanisms related to calcium imbalance, endoplasmic ERS, and mitochondrial damage are closely related to the progression of CLD and the activation of NLRP3 inflammasomes. During metabolic-related fatty liver disease, the expression of NLRP3



inflammasomes increases, and the homeostasis of MAMs is disrupted. Quercetin intervention has been shown to effectively protect against liver damage and reverse changes in MAMs. The results suggest that quercetin may exert its protective effects through the MAMs-NLRP3 pathway, providing new insights for nutritional interventions in the treatment of metabolic-related fatty liver disease. Quercetin inhibits alcohol-induced hepatocyte ferroptosis by regulating the PERK-MAMs pathway, thereby alleviating alcohol-induced liver injury<sup>[62]</sup>.

#### ***Vanillic acid***

One such TCM component is vanillic acid (VA), an edible plant compound, which has been studied for its beneficial effects on calcium ( $\text{Ca}^{2+}$ ) complications induced by hyperinsulinemia (HI), intracellular homeostasis, MAM integrity, and liver metabolism under *in vivo* and *in vitro* conditions. VA has been found to possess various pharmacological effects, such as antioxidant, antidiabetic, and anti-inflammatory properties<sup>[63]</sup>. The liver plays a crucial role in regulating lipid metabolism balance in the body, and excessive lipid accumulation in liver cells contributes to the occurrence and development of NAFLD. During the HI phase in HepG2 cells, distorted MAMs lead to insulin resistance (IR) and excessive lipid synthesis, accompanied by inflammation and oxidative stress. VA effectively protects ER and  $\text{Ca}^{2+}$  homeostasis during the HI phase<sup>[64]</sup>.

#### ***Compound glycyrrhizin tablets***

Compound glycyrrhizin tablets have a positive effect on the prevention and treatment of liver diseases, with glycyrrhizin, glycine, and cysteine hydrochloride as its main components<sup>[65]</sup>. Animal experiments have shown that compound glycyrrhizin tablets can protect the liver by inhibiting the transduction of the NF- $\kappa$ B pro-inflammatory signaling pathway and the expression of downstream inflammatory molecules<sup>[66]</sup>. Studies have found that glycyrrhizin not only inhibits the expression of IL-6 and IL-8, significantly increasing the activity of liver cells, but also promotes the expression of

SOD and glutathione, and inhibits malondialdehyde synthesis, thereby alleviating oxidative stress-induced liver cell damage. Glycine can inhibit the generation of reactive ROS, reduce ROS and lipid peroxidation, and alleviate liver damage caused by oxidative stress reactions<sup>[67]</sup>.

## **CONCLUSION**

### ***Treatment and prospect***

The signaling crosstalk between mitochondria and ER, mainly mediated by ROS and calcium loading, influences each other's endoplasmic reticulum and mitochondrial function mechanisms to regulate CLD. The mechanisms of mitochondrial and ER dysfunction are complex and diverse, and they are interrelated and interact with each other. <sup>2</sup> With the deepening of the basic theoretical research on the structure and function of mitochondria and ER, it is possible to combine the application of genomics, proteomics, metabolomics, and other modern technologies to explore in depth the signaling links between mitochondrial and ER function in various CLDs. In this paper, we elucidate the treatment of CLDs by TCM from traditional Chinese medicine to provide new ideas and means for the diagnosis and treatment of CLDs in the future. MAMs are the key factors linking the material and communication signals of mitochondria and ER. However, there are more methods for evaluating MAMs, and the accuracy of their detection varies in different studies, such as subjective judgment errors in techniques such as fluorescence localization. Under different pathologic conditions, the body's compensatory effects on MAMs proteins can vary. Therefore, future studies need to further define the criteria for judging the number of MAMs, such as the promotion of dynamic observation techniques at the cellular level, which will increase the credibility and rigor of the study.

## **REFERENCES**

- 1 **Zhang C**, Zhao Y, Yu M, Qin J, Ye B, Wang Q. Mitochondrial Dysfunction and Chronic Liver Disease. *Curr Issues Mol Biol* 2022; **44**: 3156-3165 [PMID: 35877442 DOI: 10.3390/cimb44070218]
- 2 **Lebeaupin C**, Vallée D, Hazari Y, Hetz C, Chevet E, Bailly-Maitre B. Endoplasmic reticulum stress signalling and the pathogenesis of non-alcoholic fatty liver disease. *J Hepatol* 2018; **69**: 927-947 [PMID: 29940269 DOI: 10.1016/j.jhep.2018.06.008]
- 3 **Rodrigues T**, Ferraz LS. Therapeutic potential of targeting mitochondrial dynamics in cancer. *Biochem Pharmacol* 2020; **182**: 114282 [PMID: 33058754 DOI: 10.1016/j.bcp.2020.114282]
- 4 **Yamada Y**, Takano Y, Satrialdi, Abe J, Hibino M, Harashima H. Therapeutic Strategies for Regulating Mitochondrial Oxidative Stress. *Biomolecules* 2020; **10** [PMID: 31948035 DOI: 10.3390/biom10010083]
- 5 **Prasun P**, Ginevic I, Oishi K. Mitochondrial dysfunction in nonalcoholic fatty liver disease and alcohol related liver disease. *Transl Gastroenterol Hepatol* 2021; **6**: 4 [PMID: 33437892 DOI: 10.21037/tgh-20-125]
- 6 **Durand M**, Coué M, Croyal M, Moyon T, Tesse A, Atger F, Ouguerram K, Jacobi D. Changes in Key Mitochondrial Lipids Accompany Mitochondrial Dysfunction and Oxidative Stress in NAFLD. *Oxid Med Cell Longev* 2021; **2021**: 9986299 [PMID: 34257827 DOI: 10.1155/2021/9986299]
- 7 **Zeng S**, Wu F, Chen M, Li Y, You M, Zhang Y, Yang P, Wei L, Ruan XZ, Zhao L, Chen Y. Inhibition of Fatty Acid Translocase (FAT/CD36) Palmitoylation Enhances Hepatic Fatty Acid  $\beta$ -Oxidation by Increasing Its Localization to Mitochondria and Interaction with Long-Chain Acyl-CoA Synthetase 1. *Antioxid Redox Signal* 2022; **36**: 1081-1100 [PMID: 35044230 DOI: 10.1089/ars.2021.0157]
- 8 **Kharroubi W**, Haj Ahmed S, Nury T, Andreoletti P, Sakly R, Hammami M, Lizard G. Mitochondrial dysfunction, oxidative stress and apoptotic induction in microglial BV-2 cells treated with sodium arsenate. *J Environ Sci (China)* 2017; **51**: 44-51 [PMID: 28115150 DOI: 10.1016/j.jes.2016.08.028]

- 9 **Buyco DG**, Martin J, Jeon S, Hooks R, Lin C, Carr R. Experimental models of metabolic and alcoholic fatty liver disease. *World J Gastroenterol* 2021; **27**: 1-18 [PMID: 33505147 DOI: 10.3748/wjg.v27.i1.1]
- 10 **Xu W**, Lu C, Yao L, Zhang F, Shao J, Zheng S. Dihydroartemisinin protects against alcoholic liver injury through alleviating hepatocyte steatosis in a farnesoid X receptor-dependent manner. *Toxicol Appl Pharmacol* 2017; **315**: 23-34 [PMID: 27939985 DOI: 10.1016/j.taap.2016.12.001]
- 11 **Ali N**, Ferrao K, Mehta KJ. Liver Iron Loading in Alcohol-Associated Liver Disease. *Am J Pathol* 2023; **193**: 1427-1439 [PMID: 36306827 DOI: 10.1016/j.ajpath.2022.08.010]
- 12 **Han D**, Johnson HS, Rao MP, Martin G, Sancheti H, Silkwood KH, Decker CW, Nguyen KT, Casian JG, Cadenas E, Kaplowitz N. Mitochondrial remodeling in the liver following chronic alcohol feeding to rats. *Free Radic Biol Med* 2017; **102**: 100-110 [PMID: 27867097 DOI: 10.1016/j.freeradbiomed.2016.11.020]
- 13 **Zai W**, Hu K, Ye J, Ding J, Huang C, Li Y, Fang Z, Wu M, Wang C, Chen J, Yuan Z. Long-Term Hepatitis B Virus Infection Induces Cytopathic Effects in Primary Human Hepatocytes, and Can Be Partially Reversed by Antiviral Therapy. *Microbiol Spectr* 2022; **10**: e0132821 [PMID: 35171034 DOI: 10.1128/spectrum.01328-21]
- 14 **Kim SJ**, Khan M, Quan J, Till A, Subramani S, Siddiqui A. Hepatitis B virus disrupts mitochondrial dynamics: induces fission and mitophagy to attenuate apoptosis. *PLoS Pathog* 2013; **9**: e1003722 [PMID: 24339771 DOI: 10.1371/journal.ppat.1003722]
- 15 **Hsu CC**, Wu LC, Hsia CY, Yin PH, Chi CW, Yeh TS, Lee HC. Energy metabolism determines the sensitivity of human hepatocellular carcinoma cells to mitochondrial inhibitors and biguanide drugs. *Oncol Rep* 2015; **34**: 1620-1628 [PMID: 26133123 DOI: 10.3892/or.2015.4092]
- 16 **Ezaki J**, Matsumoto N, Takeda-Ezaki M, Komatsu M, Takahashi K, Hiraoka Y, Taka H, Fujimura T, Takehana K, Yoshida M, Iwata J, Tanida I, Furuya N, Zheng DM, Tada N, Tanaka K, Kominami E, Ueno T. Liver autophagy contributes to the maintenance of blood glucose and amino acid levels. *Autophagy* 2011; **7**: 727-736 [PMID: 21471734 DOI: 10.4161/auto.7.7.15371]

- 17 **Yong H**. The role of mitochondrial damage mediated by abnormal TDP-43 aggregation in CCl<sub>4</sub>-induced liver fibrosis and its mechanism. Shandong University 2022 [DOI: 10.27272/d.cnki.gshdu.2022.004748]
- 18 **Ali SS**, Medda N, Dutta SM, Patra R, Maiti S. Protection against Mitochondrial Oxidative-Stress by Flesh-Extract of Edible Freshwater Snail *Bellamya bengalensis* Prevents Arsenic Induced DNA and Tissue Damage. *Anticancer Agents Med Chem* 2020; **20**: 1266-1273 [PMID: 32275495 DOI: 10.2174/1871520620666200410081535]
- 19 **Niemann J**, John C, Schröder S, Koch F, Ibrahim SM, Schultz J, Tiedge M, Baltrusch S. An mtDNA mutation accelerates liver aging by interfering with the ROS response and mitochondrial life cycle. *Free Radic Biol Med* 2017; **102**: 174-187 [PMID: 27890640 DOI: 10.1016/j.freeradbiomed.2016.11.035]
- 20 **Wu A**, Feng B, Yu J, Yan L, Che L, Zhuo Y, Luo Y, Yu B, Wu D, Chen D. Fibroblast growth factor 21 attenuates iron overload-induced liver injury and fibrosis by inhibiting ferroptosis. *Redox Biol* 2021; **46**: 102131 [PMID: 34530349 DOI: 10.1016/j.redox.2021.102131]
- 21 **Zhang Z**, Guo M, Shen M, Kong D, Zhang F, Shao J, Tan S, Wang S, Chen A, Cao P, Zheng S. The BRD7-P53-SLC25A28 axis regulates ferroptosis in hepatic stellate cells. *Redox Biol* 2020; **36**: 101619 [PMID: 32863216 DOI: 10.1016/j.redox.2020.101619]
- 22 **Ren T**, Zhang H, Wang J, Zhu J, Jin M, Wu Y, Guo X, Ji L, Huang Q, Zhang H, Yang H, Xing J. MCU-dependent mitochondrial Ca<sup>2+</sup> inhibits NAD(+)/SIRT3/SOD2 pathway to promote ROS production and metastasis of HCC cells. *Oncogene* 2017; **36**: 5897-5909 [PMID: 28650465 DOI: 10.1038/onc.2017.167]
- 23 **Yu HY**, Song YH, Zhao L, Liu TQ, Bao XQ. Research progress on endoplasmic reticulum stress and alcoholic liver disease. *Huagong Shikan* 2021; **35**: 32-36 [DOI: 10.16597/j.cnki.issn.1002-154x.2021.10.009]
- 24 **Lebeaupin C**, Vallée D, Hazari Y, Hetz C, Chevet E, Bailly-Maitre B. Endoplasmic reticulum stress signalling and the pathogenesis of non-alcoholic fatty liver disease. *J Hepatol* 2018; **69**: 927-947 [PMID: 29940269 DOI: 10.1016/j.jhep.2018.06.008]

- 25 **Day CP**, James OF. Steatohepatitis: a tale of two "hits"? *Gastroenterology* 1998; **114**: 842-845 [PMID: 9547102 DOI: 10.1016/s0016-5085(98)70599-2]
- 26 **Song MJ**, Malhi H. The unfolded protein response and hepatic lipid metabolism in non alcoholic fatty liver disease. *Pharmacol Ther* 2019; **203**: 107401 [PMID: 31419516 DOI: 10.1016/j.pharmthera.2019.107401]
- 27 **Mota M**, Banini BA, Cazanave SC, Sanyal AJ. Molecular mechanisms of lipotoxicity and glucotoxicity in nonalcoholic fatty liver disease. *Metabolism* 2016; **65**: 1049-1061 [PMID: 26997538 DOI: 10.1016/j.metabol.2016.02.014]
- 28 **Flessa CM**, Kyrou I, Nasiri-Ansari N, Kaltsas G, Papavassiliou AG, Kassi E, Randeva HS. Endoplasmic Reticulum Stress and Autophagy in the Pathogenesis of Non-alcoholic Fatty Liver Disease (NAFLD): Current Evidence and Perspectives. *Curr Obes Rep* 2021; **10**: 134-161 [PMID: 33751456 DOI: 10.1007/s13679-021-00431-3]
- 29 **Ji C**, Kaplowitz N. Betaine decreases hyperhomocysteinemia, endoplasmic reticulum stress, and liver injury in alcohol-fed mice. *Gastroenterology* 2003; **124**: 1488-1499 [PMID: 12730887 DOI: 10.1016/s0016-5085(03)00276-2]
- 30 **Esfandiari F**, Villanueva JA, Wong DH, French SW, Halsted CH. Chronic ethanol feeding and folate deficiency activate hepatic endoplasmic reticulum stress pathway in micropigs. *Am J Physiol Gastrointest Liver Physiol* 2005; **289**: G54-G63 [PMID: 15705656 DOI: 10.1152/ajpgi.00542.2004]
- 31 **Sun J**, Li B, Sun A, Zhao K, Ma Y, Zhao J, Pan H, Song Q, Wang Y, Yu C, Wang C, Zhang H, Zhang W, Kong C. Comprehensive analysis of aberrantly expressed profiles of messenger RNA in alcoholic liver disease. *J Cell Biochem* 2019; **120**: 4248-4254 [PMID: 30294942 DOI: 10.1002/jcb.27710]
- 32 **Jäger R**, Bertrand MJ, Gorman AM, Vandenabeele P, Samali A. The unfolded protein response at the crossroads of cellular life and death during endoplasmic reticulum stress. *Biol Cell* 2012; **104**: 259-270 [PMID: 22268789 DOI: 10.1111/boc.201100055]
- 33 **Yvamoto EY**, Ferreira RF, Nogueira V, Pinhe MA, Tenani GD, Andrade JG, Baitello ME, Gregório ML, Fucuta PS, Silva RF, Souza DR, Silva RC. Influence of vascular



endothelial growth factor and alpha-fetoprotein on hepatocellular carcinoma. *Genet Mol Res* 2015; **14**: 17453-17462 [PMID: 26782388 DOI: 10.4238/2015.December.21.16]

34 **Xia C**, Liu Y, Chen Z, Zheng M. Involvement of Interleukin 6 in Hepatitis B Viral Infection. *Cell Physiol Biochem* 2015; **37**: 677-686 [PMID: 26343270 DOI: 10.1159/000430386]

35 **Chen WT**, Tseng CC, Pfaffenbach K, Kanel G, Luo B, Stiles BL, Lee AS. Liver-specific knockout of GRP94 in mice disrupts cell adhesion, activates liver progenitor cells, and accelerates liver tumorigenesis. *Hepatology* 2014; **59**: 947-957 [PMID: 24027047 DOI: 10.1002/hep.26711]

36 **Kropski JA**, Blackwell TS. Endoplasmic reticulum stress in the pathogenesis of fibrotic disease. *J Clin Invest* 2018; **128**: 64-73 [PMID: 29293089 DOI: 10.1172/JCI93560]

37 **Jo HJ**, Yang JW, Park JH, Choi ES, Lim CS, Lee S, Han CY. Endoplasmic Reticulum Stress Increases DUSP5 Expression *via* PERK-CHOP Pathway, Leading to Hepatocyte Death. *Int J Mol Sci* 2019; **20** [PMID: 31491992 DOI: 10.3390/ijms20184369]

38 **Zuo L**, Zhu Y, Hu L, Liu Y, Wang Y, Hu Y, Wang H, Pan X, Li K, Du N, Huang Y. PI3-kinase/Akt pathway-regulated membrane transportation of acid-sensing ion channel 1a/Calcium ion influx/endoplasmic reticulum stress activation on PDGF-induced HSC Activation. *J Cell Mol Med* 2019; **23**: 3940-3950 [PMID: 30938088 DOI: 10.1111/jcmm.14275]

39 **Haze K**, Yoshida H, Yanagi H, Yura T, Mori K. Mammalian transcription factor ATF6 is synthesized as a transmembrane protein and activated by proteolysis in response to endoplasmic reticulum stress. *Mol Biol Cell* 1999; **10**: 3787-3799 [PMID: 10564271 DOI: 10.1091/mbc.10.11.3787]

40 **Yamamoto K**, Sato T, Matsui T, Sato M, Okada T, Yoshida H, Harada A, Mori K. Transcriptional induction of mammalian ER quality control proteins is mediated by single or combined action of ATF6alpha and XBP1. *Dev Cell* 2007; **13**: 365-376 [PMID: 17765680 DOI: 10.1016/j.devcel.2007.07.018]

- 41 **Khaled J**, Kopsida M, Lennernäs H, Heindryckx F. Drug Resistance and Endoplasmic Reticulum Stress in Hepatocellular Carcinoma. *Cells* 2022; **11** [PMID: 35203283 DOI: 10.3390/cells11040632]
- 42 **Friedl P**, Locker J, Sahai E, Segall JE. Classifying collective cancer cell invasion. *Nat Cell Biol* 2012; **14**: 777-783 [PMID: 22854810 DOI: 10.1038/ncb2548]
- 43 **Talmadge JE**, Fidler IJ. AACR centennial series: the biology of cancer metastasis: historical perspective. *Cancer Res* 2010; **70**: 5649-5669 [PMID: 20610625 DOI: 10.1158/0008-5472.CAN-10-1040]
- 44 **Gao LY**, Liang BY, Jin C, Li YJ, Wang XY, Zhang F, Zheng SZ. Advances in the role of neovascularization in the development of liver diseases. *Shengli Kexue Jinzhan* 2020; **51**: 193-197 [DOI: 10.3969/j.issn.0559-7765.2020.03.008]
- 45 **Wang WD**, Chen DF. Endoplasmic reticulum stress mitochondrial dysfunction and nonalcoholic fatty liver disease. *Chongqing Yixue* 2009; **38**: 2 [DOI: 10.3969/j.issn.1671-8348.2009.04.020]
- 46 **Agrawal RR**, Larrea D, Xu Y, Shi L, Zirpoli H, Cummins LG, Emmanuele V, Song D, Yun TD, Macaluso FP, Min W, Kernie SG, Deckelbaum RJ, Area-Gomez E. Alzheimer's-Associated Upregulation of Mitochondria-Associated ER Membranes After Traumatic Brain Injury. *Cell Mol Neurobiol* 2023; **43**: 2219-2241 [PMID: 36571634 DOI: 10.1007/s10571-022-01299-0]
- 47 **Hernández-Alvarez MI**, Sebastián D, Vives S, Ivanova S, Bartoccioni P, Kakimoto P, Plana N, Veiga SR, Hernández V, Vasconcelos N, Peddinti G, Adrover A, Jové M, Pamplona R, Gordaliza-Alaguero I, Calvo E, Cabré N, Castro R, Kuzmanic A, Boutant M, Sala D, Hyotylainen T, Orešič M, Fort J, Errasti-Murugarren E, Rodrigues CMP, Orozco M, Joven J, Cantó C, Palacin M, Fernández-Veledo S, Vendrell J, Zorzano A. Deficient Endoplasmic Reticulum-Mitochondrial Phosphatidylserine Transfer Causes Liver Disease. *Cell* 2019; **177**: 881-895.e17 [PMID: 31051106 DOI: 10.1016/j.cell.2019.04.010]

- 48 **Filadi R**, Greotti E, Turacchio G, Luini A, Pozzan T, Pizzo P. Mitofusin 2 ablation increases endoplasmic reticulum-mitochondria coupling. *Proc Natl Acad Sci U S A* 2015; **112**: E2174-E2181 [PMID: 25870285 DOI: 10.1073/pnas.1504880112]
- 49 **Doghman-Bouguerra M**, Granatiero V, Sbiera S, Sbiera I, Lacas-Gervais S, Brau F, Fassnacht M, Rizzuto R, Lalli E. FATE1 antagonizes calcium- and drug-induced apoptosis by uncoupling ER and mitochondria. *EMBO Rep* 2016; **17**: 1264-1280 [PMID: 27402544 DOI: 10.15252/embr.201541504]
- 50 **Cao Y**, Chen Z, Hu J, Feng J, Zhu Z, Fan Y, Lin Q, Ding G. Mfn2 Regulates High Glucose-Induced MAMs Dysfunction and Apoptosis in Podocytes *via* PERK Pathway. *Front Cell Dev Biol* 2021; **9**: 769213 [PMID: 34988075 DOI: 10.3389/fcell.2021.769213]
- 51 **Broggini-Tenzer A**, Sharma A, Nytko KJ, Bender S, Vuong V, Orlowski K, Hug D, O'Reilly T, Pruschy M. Combined treatment strategies for microtubule stabilizing agent-resistant tumors. *J Natl Cancer Inst* 2015; **107** [PMID: 25694444 DOI: 10.1093/jnci/dju504]
- 52 **Gilady SY**, Bui M, Lynes EM, Benson MD, Watts R, Vance JE, Simmen T. Ero1alpha requires oxidizing and normoxic conditions to localize to the mitochondria-associated membrane (MAM). *Cell Stress Chaperones* 2010; **15**: 619-629 [PMID: 20186508 DOI: 10.1007/s12192-010-0174-1]
- 53 **Voelker DR**. Reconstitution of phosphatidylserine import into rat liver mitochondria. *J Biol Chem* 1989; **264**: 8019-8025 [PMID: 2542259]
- 54 **Doghman-Bouguerra M**, Lalli E. ER-mitochondria interactions: Both strength and weakness within cancer cells. *Biochim Biophys Acta Mol Cell Res* 2019; **1866**: 650-662 [PMID: 30668969 DOI: 10.1016/j.bbamcr.2019.01.009]
- 55 **Huang Q**, Zhan L, Cao H, Li J, Lyu Y, Guo X, Zhang J, Ji L, Ren T, An J, Liu B, Nie Y, Xing J. Increased mitochondrial fission promotes autophagy and hepatocellular carcinoma cell survival through the ROS-modulated coordinated regulation of the NFkB and TP53 pathways. *Autophagy* 2016; **12**: 999-1014 [PMID: 27124102 DOI: 10.1080/15548627.2016.1166318]

- 56 **Zhao H**, Lv FF, San RX, Wang XZ, Li ZH, Xie W. Clinical efficacy of silymarin capsule in the treatment of drug-induced liver injury. *Gansu Yiyao* 2021; **40**: 490-493 [DOI: 10.15975/j.cnki.gsyy.2021.06.003]
- 57 **Gillessen A**, Schmidt HH. Silymarin as Supportive Treatment in Liver Diseases: A Narrative Review. *Adv Ther* 2020; **37**: 1279-1301 [PMID: 32065376 DOI: 10.1007/s12325-020-01251-y]
- 58 **Federico A**, Dallio M, Loguercio C. Silymarin/Silybin and Chronic Liver Disease: A Marriage of Many Years. *Molecules* 2017; **22** [PMID: 28125040 DOI: 10.3390/molecules22020191]
- 59 **Liu S**, Yin R, Yang Z, Wei F, Hu J. The effects of rhein on D-GalN/LPS-induced acute liver injury in mice: Results from gut microbiome-metabolomics and host transcriptome analysis. *Front Immunol* 2022; **13**: 971409 [PMID: 36389730 DOI: 10.3389/fimmu.2022.971409]
- 60 **Wang A**, Jiang H, Liu Y, Chen J, Zhou X, Zhao C, Chen X, Lin M. Rhein induces liver cancer cells apoptosis *via* activating ROS-dependent JNK/Jun/caspase-3 signaling pathway. *J Cancer* 2020; **11**: 500-507 [PMID: 31897245 DOI: 10.7150/jca.30381]
- 61 **Malterud KE**, Farbrot TL, Huse AE, Sund RB. Antioxidant and radical scavenging effects of anthraquinones and anthrones. *Pharmacology* 1993; **47 Suppl 1**: 77-85 [PMID: 8234446 DOI: 10.1159/000139846]
- 62 **Mei GB**, Chen L, Jiang CJ, Yao P. Protection of quercetin against metabolism-related fatty liver injury in T2DM mice. *Huazhong Keji Daxue Xuebao* 2021; **50**: 561-565, 573 [DOI: 10.3870/j.issn.1672-0741.2021.05.002]
- 63 **Grattagliano I**, de Bari O, Bernardo TC, Oliveira PJ, Wang DQ, Portincasa P. Role of mitochondria in nonalcoholic fatty liver disease--from origin to propagation. *Clin Biochem* 2012; **45**: 610-618 [PMID: 22484459 DOI: 10.1016/j.clinbiochem.2012.03.024]
- 64 **Kumar P**, Liu C, Suliburk J, Hsu JW, Muthupillai R, Jahoor F, Minard CG, Taffet GE, Sekhar RV. Supplementing Glycine and N-Acetylcysteine (GlyNAC) in Older Adults Improves Glutathione Deficiency, Oxidative Stress, Mitochondrial Dysfunction, Inflammation, Physical Function, and Aging Hallmarks: A Randomized Clinical Trial. *J*

*Gerontol A Biol Sci Med Sci* 2023; **78**: 75-89 [PMID: 35975308 DOI: 10.1093/gerona/glac135]

65 **Lian H**, Zhou T, Hu L, Zhang HQ. Clinical efficacy of triple therapy with bitter yellow injection, compound glycyrrhizin and adefovir in the treatment of HBeAg-positive chronic hepatitis B and its effect on patients' immune function. *Xiandai Zhongxiyi Jiehe Zazhi* 2017; **26**: 1314-1316 [DOI: 10.3969/j.issn.1008-8849.2017.12.020]

66 **Qiu HP**. Study on the mechanism of glycyrrhizin affecting acute liver injury in mice through NF- $\kappa$ B pathway. *Chongqing Yixue* 2016; **45**: 3 [DOI: 10.3969/j.issn.1671-8348.2016.30.011]

67 **Yang XY**, Su XL. Pathway mechanism of action of hepatoprotective efficacy of glycyrrhizic acid. *Zhongguo Zuzhi HuaXue Yu Xibao Huaxue Zazhi* 2019; **28**: 5 [DOI: 10.16705/j.cnki.1004-1850.2019.01.012]

# 9%

SIMILARITY INDEX

### PRIMARY SOURCES

- 1

Kaitlyn G. Jackson, Grayson W. Way, Jing Zeng, Marissa K. Lipp, Huiping Zhou. "The dynamic role of endoplasmic reticulum stress in chronic liver disease", The American Journal of Pathology, 2023  
Crossref

95 words — 2%
- 2

[www.mdpi.com](http://www.mdpi.com)  
Internet

92 words — 2%
- 3

[cris.vtt.fi](http://cris.vtt.fi)  
Internet

45 words — 1%
- 4

Si-Wei Xia, Zhi-Min Wang, Su-Min Sun, Ying Su et al. "Endoplasmic reticulum stress and protein degradation in chronic liver disease", Pharmacological Research, 2020  
Crossref

17 words — < 1%
- 5

[www.science.gov](http://www.science.gov)  
Internet

16 words — < 1%
- 6

Cheng Ji. "Mechanisms of Alcohol-Induced Endoplasmic Reticulum Stress and Organ Injuries", Biochemistry Research International, 2012  
Crossref

14 words — < 1%
- 7

Fabiola Lilí Sarmiento-Salinas, Alam Delgado-Magallón, José Benito Montes-Alvarado, Dalia

14 words — < 1%



Ramírez-Ramírez et al. "Breast Cancer Subtypes Present a Differential Production of Reactive Oxygen Species (ROS) and Susceptibility to Antioxidant Treatment", *Frontiers in Oncology*, 2019

Crossref

8 [pubmed.ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov)

Internet

14 words — < 1%

9 "Surgery: Adult", *Epilepsia*, 10/2005

Crossref

13 words — < 1%

10 [trialsjournal.biomedcentral.com](https://trialsjournal.biomedcentral.com)

Internet

13 words — < 1%

11 Entaz Bahar, Hyongsuk Kim, Hyonok Yoon. "ER Stress-Mediated Signaling: Action Potential and Ca<sup>2+</sup> as Key Players", *International Journal of Molecular Sciences*, 2016

Crossref

12 words — < 1%

12 Hristina Ivanova, Martijn Kerkhofs, Rita M. La Rovere, Geert Bultynck. "Endoplasmic Reticulum–Mitochondrial Ca<sup>2+</sup> Fluxes Underlying Cancer Cell Survival", *Frontiers in Oncology*, 2017

Crossref

12 words — < 1%

13 Lei Zhan, Haiyan Cao, Gang Wang, Yinghua Lyu, Xiacheng Sun, Jiase An, Zhenbiao Wu, Qichao Huang, Bingrong Liu, Jinliang Xing. "Drp1-mediated mitochondrial fission promotes cell proliferation through crosstalk of p53 and NF- $\kappa$ B pathways in hepatocellular carcinoma", *Oncotarget*, 2016

Crossref

12 words — < 1%

14 Wenting Dai, Lei Jiang. "Dysregulated Mitochondrial Dynamics and Metabolism in

12 words — < 1%

---

15

[f6publishing.blob.core.windows.net](#)  
Internet

12 words — < 1%

---

16

[www.frontiersin.org](#)  
Internet

12 words — < 1%