

Name of journal: *World Journal of Gastrointestinal Oncology*

ESPS Manuscript NO: 30806

Manuscript Type: Original Article

Observational Study

Macroscopic appearance of Type IV and giant Type III is a high risk for a poor prognosis in pathological stage II/III advanced gastric cancer with postoperative adjuvant chemotherapy

Keishi Yamashita, Akira Ema, Kei Hosoda, Hiroaki Mieno, Hiromitsu Moriya,
Natsuya Katada, Masahiko Watanabe

Match Overview

- 1** **Internet** 20 words
crawled on 23-Dec-2016
www.wjgnet.com 1%
- 2** **Crossref** 14 words
Waraya, Mina, Keishi Yamashita, Akira Ema, Natsuya Katada, Shiro Kikuchi, and Masahiko Watanabe. "Exclusive Ass" <1%
- 3** **Crossref** 13 words
Yamashita, K., H. Ushiku, N. Katada, K. Hosoda, H. Moriya, H. Mieno, S. Kikuchi, K. Hoshi, and M. Watanabe. "Reduced" <1%

Macroscopic appearance of Type IV and giant Type III is a high risk for a poor prc

全部 图片 视频 新闻 地图 图书

找到约 2,110,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Gastric Adenocarcinoma - NCBI - National Institutes of Health](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1356843/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1356843/>

Despite the decreasing worldwide incidence, **gastric cancer** accounts for 3% to ... present with **advanced pathologic stage** and can expect a median **survival** of 24 ... as a **risk factor** for **gastric cancer**, identified 3 groups: those with **high** (Koreans, ... The Lauren classification divides **gastric cancer** into 2 major histologic **types**: ...

[Prognostic factors in patients with diffuse type gastric cancer \(linitis ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3351946/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3351946/>

27 Jan 2011 ... Patients with a diffuse **type gastric cancer** with a signet ring ... all included patients presented with a locally **advanced tumor stage** ... The **macroscopic appearance** was classified during preoperative endoscopy as Borrmann **type III** or **IV**. ... optimal cut-off points for discrimination of low- and **high-risk** patients.

[Histological Complete Response in a Case of Advanced Gastric ...](https://jjco.oxfordjournals.org/content/33/11/584.full)

<https://jjco.oxfordjournals.org/content/33/11/584.full>

The patient was treated with **chemotherapy** consisting of S-1 and low-dose ... lymph nodes, confirming a **pathological** complete response (CR grade 3). ... However, for **stage III** and **IV advanced gastric cancer**, surgical **treatment** ... Endoscopy of the upper gastrointestinal tract (GIF) showed **type 3 advanced gastric carcinoma** ...

[Stomach cancer - Wikipedia](https://en.wikipedia.org/wiki/Stomach_cancer)

https://en.wikipedia.org/wiki/Stomach_cancer

Stomach cancer, also known as **gastric cancer**, is cancer developing from the lining of the ... Certain **types** of *H. pylori* have greater **risks** than others. ... Most of the time, **stomach cancer** develops in **stages** over years. ... endoscopy in Scotland, patients were diagnosed 7% in **Stage I** 17% in **Stage II**, and 28% in **Stage III**.

[Undifferentiated Pleomorphic Sarcoma: Diagnosis, Treatment and ...](#)

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 2,450,000 条结果 (用时 0.97 秒)

Gastric Adenocarcinoma - NCBI - National Institutes of Health

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: BJ Dicken - 2005 - 被引用次数: 585 - 相关文章

Despite the decreasing worldwide incidence, **gastric cancer** accounts for 3% to ... present with **advanced pathologic stage** and can expect a median **survival** of 24 ... as a **risk factor** for **gastric cancer**, identified 3 groups: those with **high** (Koreans, ... The Lauren classification divides **gastric cancer** into 2 major histologic **types**: ...

Full Text (HTML) - Japanese Journal of Clinical Oncology - Oxford ...

<https://jjco.oxfordjournals.org/content/33/11/584.full> - 翻译此页

作者: T Takahashi - 2003 - 被引用次数: 21 - 相关文章

The patient was **treated with chemotherapy** consisting of S-1 and low-dose ... lymph nodes, confirming a **pathological** complete response (CR grade 3). ... However, for **stage III** and **IV advanced gastric cancer**, surgical **treatment** ... Endoscopy of the upper gastrointestinal tract (GIF) showed **type 3 advanced gastric carcinoma** ...

[PDF] Histological Complete Response in a Case of Advanced ... - CiteSeerX

citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.485.2756...type... ▼ 翻译此页

作者: T Takahashi - 2003 - 被引用次数: 21 - 相关文章

Cancer Treated by **Chemotherapy** with S-1 Plus Low-dose Cisplatin ... with a **type 3 tumor** in the cardia with lymph node metastases, determined by gastrofiberscope ... primary lesion and lymph nodes, confirming a **pathological** complete response (CR grade 3). ... **survival rates** of **stage III** and **IV** patients were reported to be.

[PDF] Gastric Cancer: Introduction - Johns Hopkins Medicine

www.hopkinsmedicine.org/gastroenterology...stomach/gastric_cancer.pdf ▼ 翻译此页

Often, a delay in diagnosis may account for the poor prognosis. ... Gastric cancer consists of two