WJG

World Journal of *Gastroenterology*

Submit a Manuscript: https://www.f6publishing.com

World J Gastroenterol 2023 December 28; 29(48): 6168-6178

DOI: 10.3748/wjg.v29.i48.6168

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

OPINION REVIEW

Challenges involved in the application of artificial intelligence in gastroenterology: The race is on!

Chrysanthos D Christou, Georgios Tsoulfas

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): B Grade C (Good): C, C, C Grade D (Fair): D Grade E (Poor): 0

P-Reviewer: Lei JJ, China; Morya AK, India; Poddighe A, Kazakhstan; Wan XH, China; Wang ZF, China

Received: July 25, 2023 Peer-review started: July 25, 2023 First decision: October 9, 2023 Revised: November 6, 2023 Accepted: December 18, 2023 Article in press: December 18, 2023 Published online: December 28, 2023



Chrysanthos D Christou, Georgios Tsoulfas, Department of Transplantation Surgery, Hippokration General Hospital, School of Medicine, Aristotle University of Thessaloniki, Thessaloniki 54622, Greece

Chrysanthos D Christou, Georgios Tsoulfas, Center for Research and Innovation in Solid Organ Transplantation, School of Medicine, Aristotle University of Thessaloniki, Thessaloniki 54622, Greece

Corresponding author: Georgios Tsoulfas, FACS, FICS, MD, PhD, Chief Doctor, Professor, Surgeon, Department of Transplantation Surgery, Hippokration General Hospital, School of Medicine, Aristotle University of Thessaloniki, 49 Konstantinoupoleos Street, Thessaloniki 54622, Greece. tsoulfasg@gmail.com

Abstract

Gastroenterology is a particularly data-rich field, generating vast repositories of data that are a fruitful ground for artificial intelligence (AI) and machine learning (ML) applications. In this opinion review, we initially elaborate on the current status of the application of AI/ML-based software in gastroenterology. Currently, AI/ML-based models have been developed in the following applications: Models integrated into the clinical setting following real-time patient data flagging patients at high risk for developing a gastrointestinal disease, models employing non-invasive parameters that provide accurate diagnoses aiming to either replace, minimize, or refine the indications of endoscopy, models utilizing genomic data to diagnose various gastrointestinal diseases, computer-aided diagnosis systems facilitating the interpretation of endoscopy images, models to facilitate treatment allocation and predict the response to treatment, and finally, models in prognosis predicting complications, recurrence following treatment, and overall survival. Then, we elaborate on several challenges and how they may negatively impact the widespread application of AI in healthcare and gastroenterology. Specifically, we elaborate on concerns regarding accuracy, cost-effectiveness, cybersecurity, interpretability, oversight, and liability. While AI is unlikely to replace physicians, it will transform the skillset demanded by future physicians to practice. Thus, physicians are expected to engage with AI to avoid becoming obsolete.

Key Words: Artificial intelligence; Machine learning; Gastroenterology; Costeffectiveness; Interpretability; Accuracy

Raisbideng® WJG | https://www.wjgnet.com

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Currently, artificial intelligence (AI) and machine learning (ML) have several applications in the prevention, diagnosis, treatment, and prognosis of various gastrointestinal diseases, including gastroesophageal reflux disease, esophageal cancer, gastric cancer, gastrointestinal bleeding, inflammatory bowel diseases, polyps, colorectal cancer, and others. Despite their promising results, AI/ML applications in gastroenterology are hindered by several challenges, including accuracy, cost-effectiveness, cybersecurity, interpretability, oversight, and liability concerns. In this opinion review, we elaborate on these challenges and present different ways to overcome them.

Citation: Christou CD, Tsoulfas G. Challenges involved in the application of artificial intelligence in gastroenterology: The race is on! World J Gastroenterol 2023; 29(48): 6168-6178 URL: https://www.wjgnet.com/1007-9327/full/v29/i48/6168.htm

DOI: https://dx.doi.org/10.3748/wjg.v29.i48.6168

INTRODUCTION

Gastroenterology is a particularly data-rich field, with data being omnipresent. Several gastroenterology norms, including patient electronic records, imaging, endoscopy images and videos, capsule endoscopy videos, genomic analyses, microbiome data, and histopathology images, generate vast data repositories[1-3]. Conforming with big data features, these repositories provide a fruitful ground for artificial intelligence (AI) applications and, more specifically, AI's subfield, machine learning (ML)[4]. By learning from these data, ML algorithms are able to predict future data points [5]. AI/ML applications have been described in many studies as the potential solution to long-standing healthcare challenges, including the need for cost reduction, enhancing diagnostic accuracy, facilitating individualized and evidence-based care, and more importantly, significantly ameliorating the morbidity and mortality associated with various diseases[6-8].

In this opinion review, we first present the current status of the application of AI in the field of gastroenterology. Then, we pose a series of challenges that these applications face and explore ways to overcome them. Specifically, we elaborate on the following challenges: Accuracy concerns, cost-effectiveness, cybersecurity, interpretability, oversight, and liability.

CURRENT STATUS

Physicians use experience, intuition, and quantifiable and non-quantifiable variables for decision-making. ML algorithms, on the other hand, perform a series of precise calculations of all the quantifiable variables to perform a certain task. Computer systems, in general, and AI/ML tools, particularly, surpass, by far, physicians in their ability to quantify multiple correlations, even in fields where the physicians hold in-depth expertise. Thus, the application of ML tools is particularly useful for performing tasks requiring extensive analytical skills such as unraveling correlations in datasets, following laboratory results trends for long periods of time, and recognizing patterns in various imaging modalities. AI classifiers commonly employed in ML tools in gastroenterology are support vector machines (SVMs), artificial neural networks (ANNs), and convolutional neural networks (CNNs). SVMs are used in supervised learning. Data points of the instances are mapped in a high-dimensional space, where the hyperplane that separates the instances based on their assigned label, retaining the highest performance, is selected[9]. SVMs can be used for linear and non-linear problems using kernel functions[10]. ANNs are inspired by the human brain and consist of an input and output layer with at least one hidden layer in between[11]. As the architectures of ANNs became more sophisticated with the addition of multiple hidden layers and even more layer connections, the concept of deep neural networks (DNNs) emerged[12,13]. A particular type of DNN, the CNN, has found profound application in gastroenterology since the demonstrated high performance in image interpretation. CNNs are based on convolution, where the image is processed using a series of filters or kernels to detect patterns within the image, such as edges and textures [14].

PREVENTION

Table 1 presents a series of studies where AI/ML models have been applied in the field of gastroenterology. These models could be integrated into the clinical setting and follow real-time patient data trends. Then, these models could flag patients at high risk of developing a plethora of gastrointestinal diseases. A notable example is ColonFlag, an ML algorithm used to stratify the risk of developing colorectal cancer. Impressively, the tool has been shown to identify patients with colorectal cancer who would otherwise have avoided screening and to identify patients at the early stages of the disease 18 to 24 mo prior to the usual diagnosis time [15,16]. Other recent studies focus on the stratification of gastric cancer development by employing electronic health care records [17,18]. These models demonstrate how AI/ML models

Table 1 Studies employing artificial intelligence/machine learning tools in the field of gastroenterology

Ref.	Parameters employed/study design	AI classifier	Sample size, control group, validation	Outcomes	Performance ¹
Prevention					
Goshen <i>et al</i> [15]	EHR data/prospective validation	LR, DT, GB	688 flagged patients, NA, NA	High risk of CRC development	19 (7.5%) CRCs were found in 254 colonoscopies
Holt <i>et al</i> [<mark>16</mark>]	EHR data/case-control study	LR, DT, GB	1893641 patients, 15 controls matched to 1 case, NA	Early detection of CRC	0.536-0.624 ^{c-index}
Huang et al[17]	EHR data/case-control study	LR, LASSO, SVM, KNN, RF	1035 patients, 3 controls matched to 1 NCGC case, 10-fold CV	Early detection of NCGC	LR: 72.4 ^{acc} , 79.3 ^{sen} , 70.4 ^{spe} . LASSO: 75.1 ^{acc} , 80 ^{sen} , 73.6 ^{spe} . SVM: 75.1 ^{acc} , 76.3 ^{sen} , 74.7 ^{spe} . KNN: 78.1 ^{acc} , 68.9 ^{sen} , 80.8 ^{spe} . RF: 72.2 ^{acc} , 77.8 ^{sen} , 70.6 ^{spe}
Briggs <i>et al</i> [18]	EHR data/case-control study	RF, SVM, LR, NB, EGBDT	40348, 7471 cases/32877 controls), hold-out validation (75:25), 5-fold CV	Early detection of oesophago-gastric cancer	87-89 ^{acc} , 0.81-0.87 ^{c-index}
Diagnosis					
Manandhar et al[20]	Gut microbiome data (fecal 16S metagenomic data)/case-control study	RF, SVM, DT, ANN	1429 patients, 729 IBD and 700 non-IBD patients, hold- out validation	Diagnosis of IBD	77-84 ^{acc} , 0.41-0.82 ^{sen} , 77-84 ^{sen} , 46-64 ^{spe}
Papa et al <mark>[21</mark>]	Gut microbiome data/case-control study	SVM, RF	105 children and young adults, 91 with IBD and 24 controls, 10-fold CV	Diagnosis of IBD in child population	0.83-0.91 ^{c-index}
Pace <i>et a</i> l[22]	Laboratory results, clinicopatho- logical parameters/retrospective cohort study	ANN	159 patients, 103 with gastroesophageal reflux and 56 controls, 20-fold CV	Diagnosis of gastroeso- phageal reflux disease	78-100 ^{acc}
Yepes et al[25]	Expression levels of 1046/315 microRNAs in gastric samples/retrospective cohort study	SVM, RF	648 gastric samples, 479 cancer and 169 controls, leave-one-out CV	Classification of non- tumor mucosa <i>vs</i> tumor sample	SVM: 94 ^{acc} , RF: 89.3 ^{acc}
Huang et al[<mark>26</mark>]	Serum expression levels of miR-21-5p, miR-22-3p, and miR-29c- 3p/retrospective cohort study	Several	192, 122 cancer samples and 70 controls, leave-one-out CV	Identification of patients with gastric cancer	56-93 ^{acc}
Alizadeh <i>et al</i> [27]	Serum expression levels of miR-663a, miR-1469, miR-92a-2-5p, miR-125b- 3p, and miR-532-5p/retrospective cohort study	ANN	671, NR, 5-fold CV	Identification of patients with pancreatic ductal adenocarcinoma	93 ^{acc} , 93 ^{sen} , 92 ^{spe}
Afshar <i>et al</i> [<mark>28</mark>]	Serum expression levels of miR-6726- 5p, miR-7111-5p, miR-1247-3p, and miR-614/retrospective cohort study	ANN	200, 50 with colorectal cancer and 150 healthy controls, hold-out validation (70:15:15)	Identification of patients with colorectal cancer	100 ^{acc} , 1 ^{c-index} , 100 ^{sen,spe}
Duttagupta et al[29]	Expression levels of 847 microRNAs in the peripheral blood/case-control study	SVM	40, 20 patients with ulcerative colitis and 20 controls, 10-fold CV	Identification of patients with ulcerative colitis	92.3-92.8 ^{acc} , 87.8- 89.5 ^{sen} , 96.2-96.8 ^{spe}
Treatment					
Morilla <i>et al</i> [30]	Colonic microRNA profiles (9 microRNAs) and five clinical factors/retrospective cohort study	DNN	76 patients, 22 responders and 54 non-responders, hold-out validation (47:29)	Prediction of response to treatment of patients with active severe ulcerative colitis	80-93 ¹ , 0.80 -0.91 ^{c-index}
Takiyama <i>et al</i> [31]	Esophago-gastro-duodenoscopy imaging/retrospective cohort study	CNN	17080 images, 363 larynx/2142 esophagus/13048 stomach/1528 duodenum, hold-out validation	Anatomical classification among the larynx, esophagus, stomach, and duodenum	0.99-1.00 ^{c-index}
Rogers <i>et al</i> [35]	Data from baseline impedance, nocturnal baseline impedance, and acid exposure time/retrospective cohort study	DT	335 patients, 210 with gastroesophageal reflux and 115 controls, NR	Prediction of response to treatment with proton pump inhibitors for patients with gastroeso-	0.31-0.938 ^{c-index}



				phageal reflux disease	
Takayama <i>et al</i> [<mark>36</mark>]	Clinicopathological parameters, treatment data/retrospective cohort study	ANN	90 patients, 32 non- responders and 58 remission-effect, hold-out validation (54:36)	Prediction of the need for operation for UC patients treated with cytoapheresis	96 ^{sen} , 97 ^{spe}
Das et al <mark>[37]</mark>	Laboratory results, clinicopatho- logical parameters/retrospective cohort study	ANN	587 patients, 246 patients with major stigmata- emergent endoscopy to 162 patients, hold-out validation (194:193:200)	Prediction of major stigmata of recent hemorrhage	89 ^{acc} , 89-96 ^{sen} , 63 -89 ^{spe}
				Prediction of the need for emergent endoscopy	61-81 ^{acc} , 61-94 ^{sen} , 48-82 ^{spe}
Chu et al[<mark>38</mark>]	Laboratory results, clinicopatho- logical parameters/retrospective cohort study	Several	189 patients, NR, hold-out validation (122:67)	Prediction of the source of GIB	69.7-94.3 ^{acc} , 0.658- 0.999 ^{c-index} , 90.1 -98.0 ^{sen} , 89-100 ^{spe}
				Prediction of the need for blood resuscitation	64.7-94.1 ^{acc} , 0.381- 0.993 ^{c-index} , 90.3 -93.9 ^{sen} , 18.4-95.5 ^{spe}
				Prediction of the need for emergent endoscopy	62.7-83.3 ^{acc} , 0.404- 0.913 ^{c-index} , 80.1 -89.1 ^{sen} , 13.8-85.7 ^{spe}
				Prediction of disposition	58.4-89.7 ^{acc} , 0.324- 0.972 ^{c-index} , 81.9 -92.9 ^{sen} , 18.4-90.9 ^{spe}
Loftus <i>et al</i> [39]	Laboratory results, clinicopatho- logical parameters/retrospective cohort study	ANN	147 patients, 41% of patients with severe lower GIB and 13 patients needed surgical intervention, hold-out validation (103:44)	Prediction of severe lower GIB	0.979 ^{c-index}
				Prediction of the need for surgical intervention	0.954 ^{c-index}
Ayaru <i>et al</i> [40]	Laboratory results, clinicopatho- logical parameters/retrospective cohort study	GB	300 patients, 88 with severe bleeding, 53 with recurrent bleeding, and 35 requiring intervention, hold-out validation (170:130)	Prediction of severe lower GIB	78-83 ^{acc}
				Prediction of recurrent bleeding	88-88 ^{acc}
				Prediction of the need for intervention	88-91 ^{acc}
Prognosis					
Sato et al[41]	Laboratory results, clinicopatho- logical parameters, tumor character- istics/retrospective cohort study	ANN	395 patients, 281 alive at 1- year-89 alive at 5 years, hold-out validation (53:27:20)	1-yr and 5-yr survival of patients with esophageal cancer following surgery	0.883-0.884 ^{c-index} , 78.1-80.7 ^{sen} , 84.7- 86.5 ^{spe}
Shung et al[42]	Laboratory results, clinicopatho- logical parameters/retrospective cohort study	GB	2357 patients, 1109 requiring either transfusion or hemostatic intervention, hold-out validation (1958:399)	Stratification of risk of patients with gastrointestinal bleeding	0.91 ^{c-index}
Zhou et al[43]	Laboratory results, clinicopatho- logical parameters, tumor character- istics/retrospective cohort study	Several	2012 patients, 405 patients with recurrence, hold-out validation (80:20)	Recurrence of gastric cancer following surgery	0.77-0.80 ^{c-index}

¹Test set results are reported.

Acc: Accuracy (%); c-index: under the receiver operating curve or C-index; Sen: Sensitivity (%); Spe: Specificity (%); ANN: Artificial neural network; CD: Chron's disease; CNN: Convolutional neural network; CRC: Colorectal cancer; CV: Cross-validation; DNN: Deep neural network; DT: Decision tree; EGBDT: Extreme gradient boosting decision tree; EHR: Electronic healthcare record; GB: Gradient boosting; IBD: Inflammatory bowel disease; KNN: Knearest neighbor; LASSO: Least absolute shrinkage and selection operator; LR: Logistic regression; NA: Not applicable; NB: Naïve Bayes; NBI: Narrowband imaging; NCGC: Noncardia gastric cancer; NPV: Negative predictive value; NR: Not recorded; RF: Random forest; SVM: Support vector machine; UC: Ulcerative colitis.

can be used at the primary care level to lessen the burden of a plethora of gastrointestinal diseases significantly.

DIAGNOSIS

Currently, endoscopy constitutes the gold standard in the diagnosis of various gastrointestinal diseases. Nevertheless, endoscopy is an unpleasant intervention for the patient, has a significant cost, and is related to a series of potential complications[19]. AI/ML models utilizing non-invasive parameters that provide comparably accurate diagnoses could



Baishideng® WJG | https://www.wjgnet.com

either replace, minimize, or refine the indications of endoscopy. Two studies have used data from the microbiota and microbiome to identify patients with inflammatory bowel disease (IBD)[20,21]. Although these models could not replace endoscopy in the diagnosis of IBD, they can be used to identify patients requiring endoscopic evaluation. However, in the diagnosis of gastroesophageal reflux disease, the employment of ML tools could substitute the use of endoscopy for patients at low risk of developing cancer[22]. These efforts demonstrate how ML tools could be used to significantly reduce the cost and complications related to endoscopy, avoiding, at the same time, an unpleasant experience for the patient.

The introduction of next-generation sequencing has widened our capabilities regarding detecting host genetic factors related to certain diseases and allowed us to unveil the complex hereditary background of several diseases by clarifying the cascade of gene interactions and the impact of the environment [23,24]. This progress allowed for the emergence of novel biomarkers, including microRNAs. In gastroenterology, various studies have employed microRNA profiling data to develop tools able to diagnose various gastrointestinal diseases. Specifically, tools have been developed that classify non-tumor from tumor mucosa from gastric samples^[25]. In addition, by employing microRNA profiling, data from serum models have been developed that identify patients with gastric cancer[26], pancreatic ductal adenocarcinoma[27], colorectal cancer^[28], and ulcerative colitis^[29]. Finally, a model has been developed using microRNA profiling from colonic samples to predict the response to treatment of patients with active severe ulcerative colitis[30]. These models again demonstrate how AI/ML tools can provide a non-invasive endoscopy alternative.

Except for replacing endoscopy, AI/ML tools could significantly enhance the endoscopy's clinical outcomes. Such models could be used for developing Computer Aided Detection or Diagnosis systems (CAD systems) to facilitate the interpretation of endoscopy images. First, these CAD systems could aid the endoscopist to navigate in the gastrointestinal tract. Existing models have demonstrated outstanding performance[31]. These CAD systems could be used as a "third eye" for the endoscopist, an "eye" that does not get tired, is not susceptible to any distractions, and can identify lesions missed by the endoscopist. Following the identification of lesions, these models could be used to classify the lesions, for example, as benign or malignant. In capsule endoscopy, CAD systems could automatically identify and then classify lesions from capsule endoscopy videos and then provide the images to the gastroenterologist for review. Such models could significantly decrease the cost associated with the labor hours required to review the capsule endoscopy videos. Several models have been developed for a plethora of gastrointestinal diseases, including IBD, gastric cancer, small bowel lesions (ulcerative/hemorrhagic), colon cancer, and others[6,32]. Another application of CAD systems is proposing an ideal site for biopsy when needed. CAD systems also find a particular application in colonoscopy in classifying benign from malignant polyps[33]. By avoiding the removal of benign polyps and correctly removing all malignant polyps with the assistance of CAD systems, the cost-efficiency of coloscopy significantly increases. Notably, regarding polyp classification, the reported negative predictive value in the literature of CAD systems is above the 90% recommended by the American Society of Gastrointestinal Endoscopy for adenoma detection[34].

TREATMENT

In addition to diagnosis, AI/ML-based tools can be employed to facilitate treatment allocation for various gastrointestinal diseases. Data regarding impedance and acid exposure time were employed in a study to develop an ML tool predicting treatment response[35]. Other studies focusing on IBD have developed tools that predict treatment response to specific drugs or the need for operation[35,36]. A different group of studies, which focused on gastrointestinal bleeding management, achieved the development of models identifying patients in need of a transfusion, emergent endoscopy, emergent surgery, and intensive care unit triage[37-40]. These studies demonstrate how AI/ML tools could be used to create frameworks for individualized, evidence-based treatment allocations.

PROGNOSIS

AI/ML algorithms have also been employed in the development of predictive tools able to predict complications, recurrence following treatment, and overall survival. Examples of such studies include a study developing an ML tool to predict overall survival in patients with esophageal cancer[41], a model predicting in-hospital death for patients with gastrointestinal bleeding[42], and an ML model predicting the chance of recurrence in operated gastric cancer patients [43]. Such models could be used for family and patient counseling.

ACCURACY CONCERNS

The accuracy of a developed AI/ML tool largely depends on the quality of the training data. The notion "garbage in, garbage out" is particularly true in ML models. Flaws and weaknesses of the datasets, such as wrongly labeled instances, low image quality, small dataset size and variability of instances, and discrepancies in the data collection processes, will be inadvertently built into the end model. For example, capsule endoscopy has significantly lower image resolution compared with other types of endoscopy images, and thus, a model trained from capsule endoscopy imaging is expected to have lower classification accuracy. Standardization of data collection methods is essential to acquire datasets of clean, high-quality data that are representative of a diverse patient population. A main challenge encountered in ML



development is class imbalance. Most of the time, normal findings represent the majority class, while patients with a disease are usually the minority class. This problem is amplified in multi-class situations where accurate classification of rare classes is particularly challenging. A series of techniques to undersample the majority class and oversample the minority class in medical datasets have been proposed that, to some extent, resolve this issue[44]. In addition, in a study, researchers developed a model able to add polyps in the endoscopic images, which can then be used for training ML models^[45]. Such methodologies of creating synthetic instances could again resolve the imbalance challenge. A different challenge in ML model development is underfitting, which can be defined as the incapability of the model to capture the variability of the data[46]. On the other hand, overfitting can be described as the inability of the model to retain its performance in datasets other than the training dataset[47]. In other words, the model's performance is significantly superior in the training set and drops in new datasets. CNNs, which are widely used in gastroenterology, are vulnerable to overfitting[14]. There is a series of proposed methodologies to resolve underfitting and overfitting challenges, including early stopping/dropout, penalty method/regularization, adjusting the learning rate and the optimizer, weight matrix initialization, data augmentation, and batch normalization [46,48]. Another intriguing point is that ML models are trained from datasets not practically available in the clinical setting. The different training variables are derived at various time points during the treatment of the patient and are not readily available at a given time. These missing parameters could become a shortcoming anchoring the use of ML in the clinical setting. All the above challenges have a common "true" solution. Physicians should be included in the development of ML models from the early stages of the procedure. Effective communication among physicians, the developing team, and the investors could address and resolve any potential shortcomings of these models. The key is to involve physicians in all the stages of the process. Physicians should be involved in problem identification to ensure that the developed tool addresses actual needs and in data collection and annotation to ensure that the data are labelled correctly, which is crucial in supervised learning. Additionally, we believe that the following steps should be followed during model development to avoid potential drawbacks: Physicians should provide a detailed description of the problem at hand to the developing team, explain what is requested by the model (the endpoint), describe the features thoroughly, cooperate to identify and engineer features of high predictive value, discuss the sample size needed based on the nature of the task (binary classification, multi-class classification, regression), ensure that the patients included in the developing procedure are a representative sample of the targeted group, conduct a rigorous clinical validation before the clinical application of the model, provide input on how the tool can be seamlessly be integrated into the existing clinical workflows without causing disruption, discuss the prospects of real-time learning, where the model continues to learn and improves following clinical application, and provide insights into ethical considerations including ensuring patient privacy and avoiding potential biases. Even, following the model's application in the clinical setting is essential to establish a framework of continuous monitoring and feedback from healthcare professionals to address model shortcomings and improve the model's efficiency. Finally, when applying AI/ML-based tools in the clinical setting is crucial to collect data on the impact of the use of such tools on the clinical outcome, the well-being of the patient, and the patient-physician relation. Failing to cooperate efficiently and to communicate what exactly is expected from the model could result in an end product that does not meet the expectations of the physicians and jeopardizes the safety of the patients.

COST-EFFECTIVENESS

Unfortunately, resources are not infinite. Especially in an era of technological advancement, with other cutting-edge technologies, such as three-dimensional printing and bioprinting, virtual and augmented reality, regenerative medicine, and robotics that also promise to facilitate patient care, AI must prove that it is worth investing in. In reality, AI, which has been at the forefront for many decades, has not yet been able to provide the expected results in healthcare. This is evident during AI winters, where investors demonstrate their frustration by cutting funding[4]. The expectations are often placed extremely high, and from their perspective, AI overpromises and constantly underdelivers. As mentioned above, effective communication among the physicians, the developing team, and the investors is crucial in agreeing on the expectations and avoiding disappointment in the end product. Nevertheless, even if physicians, developers, and investors are satisfied, the true success of the model should be assessed in association with the clinical outcome. Thus, conducting a series of cost-effective analyses is crucial to determine whether AI applications' benefits in healthcare are worth the associated cost. The costing of new technologies is generally challenging due to the lack of pre-existing data, and it involves bottom-up costing approaches, which typically require more resources and are time-consuming[49]. The cost associated with new technologies tends to drop over time; thus, cost estimation studies will need to be updated throughout the lifecycle of the AI application. Despite its importance, the topic of the cost-effectiveness of the application of AI in medicine is poorly investigated, mainly due to the lack of real AI applications in the clinical setting. There are several ways to improve the cost-effectiveness of implementing AI/ML-based technologies in healthcare. First, a series of models has been developed and patented with a high performance in performing several tasks, such as CAD systems that classify benign from malignant polyps. Thus, employing existing or pre-trained models rather than developing new tools from scratch could significantly decrease the cost of adopting these technologies in clinical practice. When, however, healthcare institutions decide to invest in developing AI/ML-based tools from scratch, a plethora of approaches could be adopted to improve the cost-effectiveness, including the utilization of collaborative platforms and open-source tools, running models on existing hardware when possible to reduce the need of specialized hardware, use of existing datasets when available to avoid data acquisition costs, and finally focus on developing tools that allow for the automatization of routine, time-consuming tasks while prioritizing high-incidence conditions with the potential for significant cost savings. One of the few studies addressing this issue is a recent study in the field of gastroenterology, where researchers

performed a Markov model microsimulation of AI application in screening colonoscopy and concluded that at the United States population level, AI application could prevent more than seven thousand colorectal cancer cases and more than two thousand related deaths, and provide a yearly saving of 290 million dollars[50]. These results are very promising, and as soon as AI models are widely applied in the clinical setting, studies proving their cost-effectiveness must be conducted.

CYBERSECURITY

Cybersecurity is a challenge not exclusive to AI but to the tendency of digitalization in general. Nevertheless, AI introduces further concerns since it requires a wholly electronic tracking of patient records. This raises concerns that a huge amount of sensitive data is vulnerable to massive disclosures[51]. For example, a transfer of 1.6 million patient records to Google DeepMind was ruled illegal in the United Kingdom^[52]. Additionally, due to the sensitive nature of data (including financial data), healthcare facilities are essentially attractive targets for cyberattacks. The introduction of AI in healthcare allows for the emergence of new vulnerabilities. Imagine an intrusion into an ML, which is now manipulated into making wrong decisions or even introducing malevolent data. For example, in a study, the researchers demonstrated how a DNN was used to add or remove lung tumors from computed tomography scans at a level of accuracy that remained undetected by specialized radiologists^[53]. Imagine a similar intrusion that will introduce colorectal cancer images during real-time colonoscopy. Such incidents could severely undermine any trust in AI in healthcare and have severe consequences on patient outcomes. How could patients and physicians be confident about any AI model suggestions, knowing that they can be manipulated? Evidently, the information technology infrastructure will need to be expanded in order to accommodate AI applications and shield these models from potential threats.

INTERPRETABILITY

Interpretability in ML refers to the degree of explainability and transparency of the model^[54]. In other words, whether the logic behind the model's decisions is understandable or not. Imagine interpretability as a spectrum, where at the "white" end, we find "white-box" models, where their inner workings and decision-making processes are fully explainable and interpretable. At the other end of the spectrum, we find "black-box" models, which are characterized by a highly complex architecture to the extent that we are unable to comprehend their inner mechanisms and decisionmaking processes. In simple tasks, for example, when the features and target variable have a linear relation, white box models demonstrate a sufficiently high performance, rendering the use of black box models redundant. However, in complex tasks with high-dimensional data, black boxes exhibit superior performance to white boxes due to their ability to capture intricate patterns and relationships within the data. However, the application of black box models in healthcare introduces a series of challenges, including undermining the trust towards AI, inability to oversight any biases or prejudiced decisions towards minority groups, failure to justify our AI-assisted decision to patients, and violation of fundamental ethical tenets that physicians should hold a basic comprehension of the devices that they use. Avoiding using "black-box" models due to the above challenges could significantly handicap the potential of AI applications in healthcare. Thus, we should aim for methodologies that allow for a level of reasoning of "black-box" model predictions. Introducing a level of interpretability in black boxes applied in healthcare is crucial to enhance patient and physician trust in these tools and facilitate clinical decision-making. Interpretability can be divided into global and local. Global interpretability offers a level of explainability of the model as a whole [55]. Ways to improve global interpretability include feature importance analysis, surrogate models, and interactive visualization tools developed to allow users to explore how the various features influence the model's predictions [55-58]. For example, a model stratifying the risk of colorectal cancer development could provide its prediction along with a notification that the prediction is based mainly on the patient's sex, age, and blood count. On the other hand, local interpretability provides the reasoning behind individual predictions[55]. Methodologies to enhance local interpretability include surrogate models, Shapley values, saliency regions, and visualization techniques[55-58]. Shapley values could be used to demonstrate how each value of each feature contributes to the model's prediction, while saliency maps provide a post-hoc visualization to comprehend which parts of the input were used by the model to reach its decision and are particularly helpful for CNNs to visualize which parts of the image were used in the model's interpretation. For example, a saliency map would highlight exactly which parts of a biopsy that the model focused on to reach its diagnosis. Providing a level of explainability for "blackbox" models is key to gaining the trust of both physicians and patients and will play a pivotal role in determining which models will dominate the industry.

OVERSIGHT AND LIABILITY CONCERNS

As acknowledged by the Food and Drug Administration (FDA), applications regarding AI/ML-based software marketing in healthcare have increased exponentially [59,60]. In early 2021, the FDA published the Artificial Intelligence/Machine Learning (AI/ML)-Based Software as a Medical Device (SaMD) Action Plan, while in 2023, it published the Marketing Submission Recommendations for a Predetermined Change Control Plan for Artificial Intelligence/Machine Learning (AI/ML)-Enabled Device Software Functions[59,61]. Similarly, the European Union has announced its first regulation on



AI applications, the EU AI Act^[62]. Evidently, governments feel the need for AI oversight and regulation. Regarding law and regulation, the true challenge for the future is to find the golden mean between too little oversight, which may have devastating and severe consequences for the well-being of our patients, and too strict regulation, which will handicap innovation. AI/ML-based tools are unique in the way that they constantly evolve following the initial audit and marketing. Thus, as the model encompasses more data and is retrained, it diverges from the original product to the extent that it can be considered an entirely different product at different time points. Who ensures the safety and credibility of the product at different time points? Evidently, a lifecycle regulatory framework is needed that provides post-marketing oversight, or alternative time-limited authorizations should be granted with periodic audits when several modifications have been made to the initial product^[63].

Liability is the main physician-related concern that could undermine the widespread application of AI in healthcare. Even if the research succeeds in providing highly accurate, reliable AI models, for AI to impact the clinical outcome, physicians will need to accept and adopt these models in their everyday lives. Liability could be the main concern that discourages physicians from doing so. To evade malpractice liability, a physician is expected to provide care at the same level as a competent, same-specialty physician, considering the available resources[64]. However, the introduction of AI makes liability a far more complicated matter. An intriguing concept to consider is the potential legal binding of AI's decisions in the future. As AI/ML-based software continues to advance and grow, it is reasonable to assume that they may eventually outperform physicians, at least in certain tasks. In such a scenario, how can physicians justify ignoring the decisions presented by AI, especially when AI's decisions are solely data-driven and lack any subjective bias?

Furthermore, who should be held liable if following the AI model's decision leads to patient harm? Currently, no legal precedent clarifies liability in cases where a patient suffers an injury due to an inaccurate output generated by AI/MLbased software [65]. In a published legal analysis, the authors present a series of scenarios based on the various combinations of whether the physicians follow the AI recommendation, whether the AI recommendation is correct, and whether a patient injury occurs. The authors' conclusion highlights that since existing laws protect physicians from liability when adhering to standard care, it incentivizes them to downplay AI's actual usefulness, potentially turning AI into a mere confirmatory tool instead of a tool to enhance the quality of care[65]. Until a comprehensive legal framework regarding liability is established, healthcare facilities have valid concerns and may hesitate to adopt AI technologies due to fears and uncertainties about potential liabilities that they may expose the facility and its staff to.

CONCLUSION

In this opinion review, we initially elaborate on the current status of the application of AI/ML-based software in gastroenterology. Currently, AI/ML-based models have been developed in the following applications: Models integrated into the clinical setting following real-time patient data flagging patients at high risk for developing a gastrointestinal disease, models employing non-invasive parameters that provide accurate diagnoses aiming to either replace, minimize, or refine the indications of endoscopy, models utilizing genomic data to diagnose various gastrointestinal diseases, CAD systems facilitating the interpretation of endoscopy images, models to facilitate treatment allocation and predict the response to treatment, and finally, models in prognosis predicting complications, recurrence following treatment, and overall survival that can be used for family and patient counseling. Then, we elaborate on several challenges and how they may negatively impact the widespread application of AI in healthcare and gastroenterology, offering possible ways to overcome them. Specifically, we have elaborated on concerns regarding accuracy, cost-effectiveness, cybersecurity, interpretability, oversight, and liability.

It is worth noting that the vast majority of AI research originates from high-income countries[66]. Despite being described as the solution that could narrow the gap in quality of care between high and low-income countries, AI may become one more brick in the wall of inequality. To avoid that, future AI initiatives should include middle and lowincome countries. In any healthcare system where AI/ML-based software will be integrated, it is essential to simultaneously integrate equally sophisticated oversight mechanisms to ensure the safety of patients; otherwise, physicians may encounter unintended ramifications.

As mentioned before, AI/ML models' decision-making is based on quantifiable parameters. However, clinicians rely on non-quantifiable parameters for decision-making. It is crucial to state that overreliance on AI could impair our critical thinking and reasoning. Thus, the ultimate goal is not AI-driven but AI-assisted clinical practice. AI could be the means to augment our intelligence by embracing the inherent complexity of medicine. Envision the hospital of the future where hospitals are fully robotic, AI and robots conduct the majority of procedures, 3D printers are used to print almost everything, from medical equipment to artificial parts, telemedicine allows patients to stay at home, virtual and augmented reality will assist physicians at all procedures, and biosensors are placed at birth diagnosing all diseases at an early stage and alerting physicians[67-69]. It is unlikely that AI will replace physicians. Nevertheless, it is also unlikely that the skillset demanded by future physicians will be in any way similar to the present. In this upcoming transformation of healthcare with the integration of cutting-edge technologies, AI is expected to be at the center. Thus, physicians are expected to engage with AI to avoid becoming obsolete.

FOOTNOTES

Author contributions: Christou CD and Tsoulfas G performed the screening of articles for eligibility; Christou CD drafted the manuscript; Tsoulfas G edited the manuscript.



Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: Greece

ORCID number: Chrysanthos D Christou 0000-0002-5417-8686; Georgios Tsoulfas 0000-0001-5043-7962.

S-Editor: Wang JJ L-Editor: Wang TQ P-Editor: Yu HG

REFERENCES

- Alizadeh M, Sampaio Moura N, Schledwitz A, Patil SA, Ravel J, Raufman JP. Big Data in Gastroenterology Research. Int J Mol Sci 2023; 24 1 [PMID: 36768780 DOI: 10.3390/ijms24032458]
- 2 Raghupathi W, Raghupathi V. Big data analytics in healthcare: promise and potential. Health Inf Sci Syst 2014; 2: 3 [PMID: 25825667 DOI: 10.1186/2047-2501-2-3
- Dash S, Shakyawar SK, Sharma M, Kaushik S. Big data in healthcare: management, analysis and future prospects. J Big Data 2019; 6: 54 [DOI: 10.1186/s40537-019-0217-0]
- SDG resource centre. Artificial Intelligence: How knowledge is created, transferred, and used. [cited 25 May 2023]. Available from: https:// 4 sdgresources.relx.com/artificial-intelligence-how-knowledge-created-transferred-and-used
- 5 Cruz JA, Wishart DS. Applications of machine learning in cancer prediction and prognosis. Cancer Inform 2007; 2: 59-77 [PMID: 19458758]
- Christou CD, Tsoulfas G. Challenges and opportunities in the application of artificial intelligence in gastroenterology and hepatology. World J 6 Gastroenterol 2021; 27: 6191-6223 [PMID: 34712027 DOI: 10.3748/wjg.v27.i37.6191]
- 7 Erratum for the Research Article: "Squalene epoxidase drives NAFLD-induced hepatocellular carcinoma and is a pharmaceutical target" by D. Liu, C. C. Wong, L. Fu, H. Chen, L. Zhao, C. Li, Y. Zhou, Y. Zhang, W. Xu, Y. Yang, B. Wu, G. Cheng, P. B.-S. Lai, N. Wong, J. J. Y. Sung, J. Yu. Sci Transl Med 2018; 10 [PMID: 30257951 DOI: 10.1126/scitranslmed.aav4522]
- Ruffle JK, Farmer AD, Aziz Q. Artificial Intelligence-Assisted Gastroenterology- Promises and Pitfalls. Am J Gastroenterol 2019; 114: 422-8 428 [PMID: 30315284 DOI: 10.1038/s41395-018-0268-4]
- Guo CY, Chou YC. A novel machine learning strategy for model selections Stepwise Support Vector Machine (StepSVM). PLoS One 2020; 9 15: e0238384 [PMID: 32853243 DOI: 10.1371/journal.pone.0238384]
- Aizerman M. Theoretical foundations of the potential function method in pattern recognition learning. Autom Remote Control 1964; 25: 821-10 837
- 11 Yang YJ, Bang CS. Application of artificial intelligence in gastroenterology. World J Gastroenterol 2019; 25: 1666-1683 [PMID: 31011253 DOI: 10.3748/wjg.v25.i14.1666]
- 12 Kriegeskorte N, Golan T. Neural network models and deep learning. Curr Biol 2019; 29: R231-R236 [PMID: 30939301 DOI: 10.1016/j.cub.2019.02.034]
- 13 LeCun Y, Bengio Y, Hinton G. Deep learning. Nature 2015; 521: 436-444 [PMID: 26017442 DOI: 10.1038/nature14539]
- Anwar SM, Majid M, Qayyum A, Awais M, Alnowami M, Khan MK. Medical Image Analysis using Convolutional Neural Networks: A 14 Review. J Med Syst 2018; 42: 226 [PMID: 30298337 DOI: 10.1007/s10916-018-1088-1]
- Goshen R, Choman E, Ran A, Muller E, Kariv R, Chodick G, Ash N, Narod S, Shalev V. Computer-Assisted Flagging of Individuals at High 15 Risk of Colorectal Cancer in a Large Health Maintenance Organization Using the ColonFlag Test. JCO Clin Cancer Inform 2018; 2: 1-8 [PMID: 30652563 DOI: 10.1200/CCI.17.00130]
- Holt TA, Virdee PS, Bankhead C, Patnick J, Nicholson BD, Fuller A, Birks J. Early detection of colorectal cancer using symptoms and the 16 ColonFlag: case-control and cohort studies [version 1; peer review: 1 approved with reservations]. NIHR Open Res 2023; 3: 6 [DOI: 10.3310/nihropenres.13360.1]
- Huang RJ, Kwon NS, Tomizawa Y, Choi AY, Hernandez-Boussard T, Hwang JH. A Comparison of Logistic Regression Against Machine 17 Learning Algorithms for Gastric Cancer Risk Prediction Within Real-World Clinical Data Streams. JCO Clin Cancer Inform 2022; 6: e2200039 [PMID: 35763703 DOI: 10.1200/CCI.22.00039]
- Briggs E, de Kamps M, Hamilton W, Johnson O, McInerney CD, Neal RD. Machine Learning for Risk Prediction of Oesophago-Gastric 18 Cancer in Primary Care: Comparison with Existing Risk-Assessment Tools. Cancers (Basel) 2022; 14 [PMID: 36291807 DOI: 10.3390/cancers14205023]
- Kavic SM, Basson MD. Complications of endoscopy. Am J Surg 2001; 181: 319-332 [PMID: 11438266 DOI: 19 10.1016/S0002-9610(01)00589-X
- Manandhar I, Alimadadi A, Aryal S, Munroe PB, Joe B, Cheng X. Gut microbiome-based supervised machine learning for clinical diagnosis 20 of inflammatory bowel diseases. Am J Physiol Gastrointest Liver Physiol 2021; 320: G328-G337 [PMID: 33439104 DOI: 10.1152/ajpgi.00360.2020]
- Papa E, Docktor M, Smillie C, Weber S, Preheim SP, Gevers D, Giannoukos G, Ciulla D, Tabbaa D, Ingram J, Schauer DB, Ward DV, 21 Korzenik JR, Xavier RJ, Bousvaros A, Alm EJ. Non-invasive mapping of the gastrointestinal microbiota identifies children with inflammatory bowel disease. PLoS One 2012; 7: e39242 [PMID: 22768065 DOI: 10.1371/journal.pone.0039242]
- 22 Pace F, Buscema M, Dominici P, Intraligi M, Baldi F, Cestari R, Passaretti S, Bianchi Porro G, Grossi E. Artificial neural networks are able to



recognize gastro-oesophageal reflux disease patients solely on the basis of clinical data. Eur J Gastroenterol Hepatol 2005; 17: 605-610 [PMID: 15879721 DOI: 10.1097/00042737-200506000-00003]

- 23 Mardis ER. Next-generation DNA sequencing methods. Annu Rev Genomics Hum Genet 2008; 9: 387-402 [PMID: 18576944 DOI: 10.1146/annurev.genom.9.081307.164359]
- Mardis ER. DNA sequencing technologies: 2006-2016. Nat Protoc 2017; 12: 213-218 [PMID: 28055035 DOI: 10.1038/nprot.2016.182] 24
- Yepes S, López R, Andrade RE, Rodriguez-Urrego PA, López-Kleine L, Torres MM. Co-expressed miRNAs in gastric adenocarcinoma. 25 Genomics 2016; 108: 93-101 [PMID: 27422560 DOI: 10.1016/j.ygeno.2016.07.002]
- 26 Huang Y, Zhu J, Li W, Zhang Z, Xiong P, Wang H, Zhang J. Serum microRNA panel excavated by machine learning as a potential biomarker for the detection of gastric cancer. Oncol Rep 2018; 39: 1338-1346 [PMID: 29286167 DOI: 10.3892/or.2017.6163]
- Alizadeh Savareh B, Asadzadeh Aghdaie H, Behmanesh A, Bashiri A, Sadeghi A, Zali M, Shams R. A machine learning approach identified a 27 diagnostic model for pancreatic cancer through using circulating microRNA signatures. Pancreatology 2020; 20: 1195-1204 [PMID: 32800647 DOI: 10.1016/j.pan.2020.07.399]
- Afshar S, Afshar S, Warden E, Manochehri H, Saidijam M. Application of Artificial Neural Network in miRNA Biomarker Selection and 28 Precise Diagnosis of Colorectal Cancer. Iran Biomed J 2019; 23: 175-183 [PMID: 30056689 DOI: 10.29252/ibj.23.3.175]
- 29 Duttagupta R, DiRienzo S, Jiang R, Bowers J, Gollub J, Kao J, Kearney K, Rudolph D, Dawany NB, Showe MK, Stamato T, Getts RC, Jones KW. Genome-wide maps of circulating miRNA biomarkers for ulcerative colitis. PLoS One 2012; 7: e31241 [PMID: 22359580 DOI: 10.1371/journal.pone.0031241]
- 30 Morilla I, Uzzan M, Laharie D, Cazals-Hatem D, Denost Q, Daniel F, Belleannee G, Bouhnik Y, Wainrib G, Panis Y, Ogier-Denis E, Treton X. Colonic MicroRNA Profiles, Identified by a Deep Learning Algorithm, That Predict Responses to Therapy of Patients With Acute Severe Ulcerative Colitis. Clin Gastroenterol Hepatol 2019; 17: 905-913 [PMID: 30223112 DOI: 10.1016/j.cgh.2018.08.068]
- Takiyama H, Ozawa T, Ishihara S, Fujishiro M, Shichijo S, Nomura S, Miura M, Tada T. Automatic anatomical classification of 31 esophagogastroduodenoscopy images using deep convolutional neural networks. Sci Rep 2018; 8: 7497 [PMID: 29760397 DOI: 10.1038/s41598-018-25842-6]
- Pang X, Zhao Z, Weng Y. The Role and Impact of Deep Learning Methods in Computer-Aided Diagnosis Using Gastrointestinal Endoscopy. 32 Diagnostics (Basel) 2021; 11 [PMID: 33919669 DOI: 10.3390/diagnostics11040694]
- Mori Y, Kudo SE, Berzin TM, Misawa M, Takeda K. Computer-aided diagnosis for colonoscopy. Endoscopy 2017; 49: 813-819 [PMID: 33 28561195 DOI: 10.1055/s-0043-109430]
- 34 Rex DK, Kahi C, O'Brien M, Levin TR, Pohl H, Rastogi A, Burgart L, Imperiale T, Ladabaum U, Cohen J, Lieberman DA. The American Society for Gastrointestinal Endoscopy PIVI (Preservation and Incorporation of Valuable Endoscopic Innovations) on real-time endoscopic assessment of the histology of diminutive colorectal polyps. Gastrointest Endosc 2011; 73: 419-422 [PMID: 21353837 DOI: 10.1016/j.gie.2011.01.023
- 35 Rogers B, Samanta S, Ghobadi K, Patel A, Savarino E, Roman S, Sifrim D, Gyawali CP. Artificial intelligence automates and augments baseline impedance measurements from pH-impedance studies in gastroesophageal reflux disease. J Gastroenterol 2021; 56: 34-41 [PMID: 33151406 DOI: 10.1007/s00535-020-01743-2]
- Takayama T, Okamoto S, Hisamatsu T, Naganuma M, Matsuoka K, Mizuno S, Bessho R, Hibi T, Kanai T. Computer-Aided Prediction of 36 Long-Term Prognosis of Patients with Ulcerative Colitis after Cytoapheresis Therapy. PLoS One 2015; 10: e0131197 [PMID: 26111148 DOI: 10.1371/journal.pone.0131197
- Das A, Ben-Menachem T, Farooq FT, Cooper GS, Chak A, Sivak MV Jr, Wong RC. Artificial neural network as a predictive instrument in 37 patients with acute nonvariceal upper gastrointestinal hemorrhage. Gastroenterology 2008; 134: 65-74 [PMID: 18061180 DOI: 10.1053/j.gastro.2007.10.037]
- Chu A, Ahn H, Halwan B, Kalmin B, Artifon EL, Barkun A, Lagoudakis MG, Kumar A. A decision support system to facilitate management 38 of patients with acute gastrointestinal bleeding. Artif Intell Med 2008; 42: 247-259 [PMID: 18063351 DOI: 10.1016/j.artmed.2007.10.003]
- 39 Loftus TJ, Brakenridge SC, Croft CA, Smith RS, Efron PA, Moore FA, Mohr AM, Jordan JR. Neural network prediction of severe lower intestinal bleeding and the need for surgical intervention. J Surg Res 2017; 212: 42-47 [PMID: 28550920 DOI: 10.1016/j.jss.2016.12.032]
- Ayaru L, Ypsilantis PP, Nanapragasam A, Choi RC, Thillanathan A, Min-Ho L, Montana G. Prediction of Outcome in Acute Lower 40 Gastrointestinal Bleeding Using Gradient Boosting. PLoS One 2015; 10: e0132485 [PMID: 26172121 DOI: 10.1371/journal.pone.0132485]
- Sato F, Shimada Y, Selaru FM, Shibata D, Maeda M, Watanabe G, Mori Y, Stass SA, Imamura M, Meltzer SJ. Prediction of survival in 41 patients with esophageal carcinoma using artificial neural networks. Cancer 2005; 103: 1596-1605 [PMID: 15751017 DOI: 10.1002/cncr.20938]
- Shung DL, Au B, Taylor RA, Tay JK, Laursen SB, Stanley AJ, Dalton HR, Ngu J, Schultz M, Laine L. Validation of a Machine Learning 42 Model That Outperforms Clinical Risk Scoring Systems for Upper Gastrointestinal Bleeding. Gastroenterology 2020; 158: 160-167 [PMID: 31562847 DOI: 10.1053/j.gastro.2019.09.009]
- Zhou C, Hu J, Wang Y, Ji MH, Tong J, Yang JJ, Xia H. A machine learning-based predictor for the identification of the recurrence of patients 43 with gastric cancer after operation. Sci Rep 2021; 11: 1571 [PMID: 33452440 DOI: 10.1038/s41598-021-81188-6]
- Rahman MM, Davis DN. Addressing the class imbalance problem in medical datasets. Int J Mach Learn Comput 2013; 3: 224 [DOI: 44 10.7763/IJMLC.2013.V3.307]
- 45 de Almeida Thomaz V, Sierra-Franco CA, Raposo AB. Training data enhancements for improving colonic polyp detection using deep convolutional neural networks. Artif Intell Med 2021; 111: 101988 [PMID: 33461694 DOI: 10.1016/j.artmed.2020.101988]
- 46 Jabbar H, Khan RZ. Methods to avoid over-fitting and under-fitting in supervised machine learning (comparative study). Comput Sci Commun Instrum Devices 2015; 70: 978-981 [DOI: 10.3850/978-981-09-5247-1_017]
- 47 Park SH, Han K. Methodologic Guide for Evaluating Clinical Performance and Effect of Artificial Intelligence Technology for Medical Diagnosis and Prediction. Radiology 2018; 286: 800-809 [PMID: 29309734 DOI: 10.1148/radiol.2017171920]
- Li Q, Yan M, Xu J. Optimizing Convolutional Neural Network Performance by Mitigating Underfitting and Overfitting. Proceedings of the 48 2021 IEEE/ACIS 19th International Conference on Computer and Information Science (ICIS); 2021 Jun 23-25; Shanghai, China: IEEE, 2001: 126-131
- Christou CD, Athanasiadou EC, Tooulias AI, Tzamalis A, Tsoulfas G. The process of estimating the cost of surgery: Providing a practical 49 framework for surgeons. Int J Health Plann Manage 2022; 37: 1926-1940 [PMID: 35191067 DOI: 10.1002/hpm.3431]
- Areia M, Mori Y, Correale L, Repici A, Bretthauer M, Sharma P, Taveira F, Spadaccini M, Antonelli G, Ebigbo A, Kudo SE, Arribas J, Barua 50 I, Kaminski MF, Messmann H, Rex DK, Dinis-Ribeiro M, Hassan C. Cost-effectiveness of artificial intelligence for screening colonoscopy: a



modelling study. Lancet Digit Health 2022; 4: e436-e444 [PMID: 35430151 DOI: 10.1016/S2589-7500(22)00042-5]

- For the Record Protecting Electronic Health Information. Washington (DC): National Academies Press (US); 1997 [PMID: 25121276] 51
- The Guardian. Royal Free breached UK data law in 1.6m patient deal with Google's DeepMind. [cited 15 May 2023]. Available from: https:// 52 /cdgma.com/nhs-breaking-data-protection-act
- Mirsky Y, Mahler T, Shelef I, Elovici Y. CT-GAN: Malicious tampering of 3D medical imagery using deep learning. 2019 Preprint. 53 Available from: arXiv:1901.03597 [DOI: 10.48550/arXiv.1901.03597]
- 54 Gilpin LH, Bau D, Yuan BZ, Bajwa A, Specter M, Kagal L. Explaining explanations: An overview of interpretability of machine learning. Proceedings of the 2018 IEEE 5th International Conference on data science and advanced analytics (DSAA); 2018 Oct 23-25; Turin, Italy: IEEE, 2018: 80-89
- Du M, Liu N, Hu X. Techniques for interpretable machine learning. 2018 Preprint. Available from: arXiv:1808.00033 [DOI: 55 10.48550/arXiv.1808.00033]
- Neto MP, Paulovich FV. Explainable Matrix Visualization for Global and Local Interpretability of Random Forest Classification Ensembles. 56 IEEE Trans Vis Comput Graph 2021; 27: 1427-1437 [PMID: 33048689 DOI: 10.1109/TVCG.2020.3030354]
- 57 Kopitar L, Cilar L, Kocbek P, Stiglic G. Local vs. global interpretability of machine learning models in type 2 diabetes mellitus screening. In: Artificial Intelligence in Medicine: Knowledge Representation and Transparent and Explainable System. Cham: Springer, 2019: 108-119
- 58 Park SH. Artificial Intelligence in Medicine: Beginners Guide. J Korean Society Radiol 2018; 78: 301 [DOI: 10.3348/jksr.2018.78.5.301]
- 59 News Medical Life Sciences. FDA's AI/ML-Based Software as a Medical Device Action Plan released. 2021. [cited 25 May 2023]. Available from: https://www.news-medical.net/news/20210112/FDAs-AIML-Based-Software-as-a-Medical-Device-Action-Plan-released.aspx#:~:text= The%20AI%2FML-Based%20Software%20as%20a%20Medical%20Device%20Action,approach%2C%20including%20device% 20transparency%20to%20users%3B%20More%20items
- Topol EJ. High-performance medicine: the convergence of human and artificial intelligence. Nat Med 2019; 25: 44-56 [PMID: 30617339 DOI: 60 10.1038/s41591-018-0300-7]
- Federal Register. Marketing Submission Recommendations for a Predetermined Change Control Plan for Artificial Intelligence/Machine 61 Learning (AI/ML) - Enabled Device Software Functions Draft Guidance for Industry and Food and Drug Administration Staff; Availability.[cited 25 May 2023]. Available from: https://www.federalregister.gov/documents/2023/04/03/2023-06786/marketing-submissionrecommendations-for-a-predetermined-change-control-plan-for-artificial
- Neuwirth RJ. The EU Artificial Intelligence Act. EU Artif Intell Act 2022; 0106 [DOI: 10.4324/9781003319436] 62
- Hwang TJ, Kesselheim AS, Vokinger KN. Lifecycle Regulation of Artificial Intelligence- and Machine Learning-Based Software Devices in 63 Medicine. JAMA 2019; 322: 2285-2286 [PMID: 31755907 DOI: 10.1001/jama.2019.16842]
- Froomkin AM, Kerr I, Pineau J. When ais outperform doctors: CONFRONTING The challenges of a tort-induced over-reliance on machine 64 learning. University of Miami Law School 2019
- Price WN 2nd, Gerke S, Cohen IG. Potential Liability for Physicians Using Artificial Intelligence. JAMA 2019; 322: 1765-1766 [PMID: 65 31584609 DOI: 10.1001/jama.2019.15064]
- Zhang L, Wang H, Li Q, Zhao MH, Zhan QM. Big data and medical research in China. BMJ 2018; 360: j5910 [PMID: 29437562 DOI: 66 10.1136/bmj.j5910]
- Vatandoost M, Litkouhi S. The future of healthcare facilities: how technology and medical advances may shape hospitals of the future. Hosp 67 *Pract Res* 2018; **4** [DOI: 10.15171/HPR.2019.01]
- Christou CD, Tsoulfas G. Role of three-dimensional printing and artificial intelligence in the management of hepatocellular carcinoma: 68 Challenges and opportunities. World J Gastrointest Oncol 2022; 14: 765-793 [PMID: 35582107 DOI: 10.4251/wjgo.v14.i4.765]
- Christou CD, Vasileiadou S, Sotiroudis G, Tsoulfas G. Three-Dimensional Printing and Bioprinting in Renal Transplantation and 69 Regenerative Medicine: Current Perspectives. J Clin Med 2023; 12 [PMID: 37892658 DOI: 10.3390/jcm12206520]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

