



## 学术搜索

找到约 10,800 条结果 (用时0.10秒)

试用新功能

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

 包括专利 包含引用

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[PDF] Routine coagulation of visible vessels may prevent delayed bleeding after endoscopic submucosal dissection-An analysis of risk factors\*

K Takizawa, I Oda, T Gotoda, C Yokoi, T Matsuda... - ..., 2008 - researchgate.net

... intervention. Blood transfusion was required in only one patient. ... gation. In this study, age and gender had no relevance to delayed bleeding after ESD. ... al. Predictors of bleeding after endoscopic mucosal resection of gastric tumors. ...

被引用次数: 211 相关文章 所有 7 个版本 引用

researchgate.net 中的 [PDF]

[PDF] Risk factors for bleeding after endoscopic submucosal dissection for gastric lesions

Y Tsuji, K Ohata, T Ito, H Chiba, T Ohya, T Gunji... - World J..., 2010 - wjgnet.com

... did not require surgical intervention; one case required blood transfusion. ... sites of immediate bleeding, suggesting that well-coagulated arteries seldom bleed. ... Asaka M. Effects of preoperative administration of omeprazole on bleeding after endoscopic submucosal dissection ...

被引用次数: 83 相关文章 所有 9 个版本 引用

wjgnet.com 中的 [PDF]

Effect of a proton pump inhibitor or an H2-receptor antagonist on prevention of bleeding from ulcer after endoscopic submucosal dissection of early gastric cancer: a ...

N Uedo, Y Takeuchi, T Yamada, R Ishihara... - The American journal of..., 2007 - nature.com

... Pepsin digests blood clots overlying the ulcer crater and its activity is pH ... trial of either famotidine or omeprazole for the prevention of bleeding after endoscopic mucosal resection ... Omeprazole versus placebo for acute upper gastrointestinal bleeding: Randomized double blind ...

被引用次数: 129 相关文章 所有 10 个版本 引用

nature.com 中的 [HTML]

Risk factors for delayed bleeding after endoscopic submucosal dissection for gastric neoplasm

K Okada, Y Yamamoto, A Kasuga, M Omae... - ... endoscopy, 2011 - Springer

... started drinking water on the day after ESD and began eating a liquid diet after second-look EGD ... of bleeding with occurrence of melena and/or hematemesis confirmed by hospital staff, or confirmed blood or coagulants in the stomach and bleeding spots confirmed ...

被引用次数: 97 相关文章 所有 11 个版本 引用



Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 26485

Manuscript Type: TOPIC HIGHLIGHT

2016 Gastrointestinal Endoscopy: Global view

Bleeding after endoscopic submucosal dissection: Risk factors and preventive methods

Kataoka Y *et al.* An overview of post-ESD bleeding

Yosuke Kataoka, Yosuke Tsuji, Yoshiki Sakaguchi, Chihiro Minatsuki,  
 Itsuko Asada-Hirayama, Keiko Niimi, Satoshi Ono, Shinya Kodashima,  
 Nobutake Yamamichi, Mitsuhiro Fujishiro, Kazuhiko Koike

### Abstract

Endoscopic submucosal dissection (ESD) has become widely accepted as a

### Match Overview

1	Internet 48 words crawled on 09-Mar-2016 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	2%
2	CrossCheck 19 words Tsuji, Yosuke, Ken Ohata, Toshiaki Gunji, Mieko Shozu, hima, Jun Hamanaka, Akiko Ohno, Takafumi Ito, Nobuta	1%
3	CrossCheck 19 words Lim, J.H. "Do antiplatelets increase the risk of bleeding... after endoscopic submucosal dissection of gastric neop...	1%
4	CrossCheck 18 words Niishizawa, Toshihiro, Hidekazu Suzuki, Satoshi Kinoshita, Osamu Goto, Takanori Kanai, and Naohisa Yahagi. "S...	1%
5	CrossCheck 18 words Matsumura, Tomoaki, Makoto Arai, Daisuke Maruoka, Kenichiro Okimoto, Shoko Minemura, Hideaki Ishigami, Kei...	1%
6	CrossCheck 17 words Fujishiro, Mitsuhiro, Philip WY Chiu, and Hsui-Po Wang. "Role of antisecretory agents for gastric endoscopic s...	1%
7	CrossCheck 17 words Mabe, Katsuhiro, Mototsugu Kato, Koji Oba, Soichi Naka...	1%

[全部](#) [图片](#) [视频](#) [新闻](#) [购物](#) [更多 ▾](#) [搜索工具](#)

找到约 167,000 条结果 (用时 0.60 秒)

## Google 学术: Bleeding after endoscopic submucosal dissection: Risk factors and preventive methods

[... bleeding after endoscopic submucosal dissection-An ... - Takizawa - 被引用次数: 216](#)

[Risk factors for bleeding after endoscopic submucosal ... - Tsuji - 被引用次数: 86](#)

[... after endoscopic submucosal dissection of early gastric ... - Uedo - 被引用次数: 133](#)

## Risk factors for bleeding after endoscopic submucosal dissection for ...

[www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central \(PMC\) ▾ 翻译此页](#)

作者: Y Tsuji - 2010 - 被引用次数: 84 - 相关文章

2010年6月21日 - AIM: To assess **risk factors for bleeding after gastric endoscopic submucosal dissection (ESD)** and to develop **preventive measures**.

## Routine coagulation of visible vessels may prevent delayed bleeding ...

[www.ncbi.nlm.nih.gov/pubmed/18322872 - 翻译此页](#)

作者: K Takizawa - 2008 - 被引用次数: 214 - 相关文章

Routine coagulation of visible vessels may **prevent** delayed **bleeding after endoscopic submucosal dissection**--an analysis of risk factors. Takizawa K(1), Oda I, ...

## Analysis of Delayed Bleeding after Endoscopic Submucosal ...

[www.hindawi.com/journals/grp/2012/875323/ ▾ 翻译此页](#)

作者: S Mukai - 2012 - 被引用次数: 25 - 相关文章

2011年11月3日 - Delayed bleeding after endoscopic submucosal dissection (ESD) ... We investigated

... have reported investigations of



全部 图片 视频 新闻 购物 更多 ▾ 搜索工具

找到约 171,000 条结果 (用时 0.64 秒)

### Google 学术 : Bleeding after endoscopic submucosal dissection: Risk factors and preventive methods

... bleeding after endoscopic submucosal dissection-An ... - Takizawa - 被引用次数 : 217

Risk factors for bleeding after endoscopic submucosal ... - Tsuji - 被引用次数 : 86

... after endoscopic submucosal dissection of early gastric ... - Uedo - 被引用次数 : 134

### Risk factors for bleeding after endoscopic submucosal dissection for ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者 : Y Tsuji - 2010 - 被引用次数 : 86 - 相关文章

2010年6月21日 - AIM: To assess risk factors for bleeding after gastric endoscopic submucosal dissection (ESD) and to develop preventive measures.

### Routine coagulation of visible vessels may prevent delayed bleeding ...

[www.ncbi.nlm.nih.gov/pubmed/18322872](http://www.ncbi.nlm.nih.gov/pubmed/18322872) - 翻译此页

作者 : K Takizawa - 2008 - 被引用次数 : 217 - 相关文章

Routine coagulation of visible vessels may prevent delayed bleeding after endoscopic submucosal dissection--an analysis of risk factors. Takizawa K(1), Oda I, ...

### Analysis of Delayed Bleeding after Endoscopic Submucosal ...

[www.hindawi.com/journals/grp/2012/875323/](http://www.hindawi.com/journals/grp/2012/875323/) ▾ 翻译此页

作者 : S Mukai - 2012 - 被引用次数 : 25 - 相关文章

2011年11月3日 - Delayed bleeding after endoscopic submucosal dissection (ESD) ... We investigated factors related to post-ESD bleeding to identify preventive measures. ... have reported investigations of risk factors for post-ESD bleeding [6, ...]

### Delayed bleeding after endoscopic submucosal dissection for non ...

[onlinelibrary.wiley.com](http://onlinelibrary.wiley.com) > > Digestive Endoscopy > Vol 27 Issue 3 - 翻译此页