



All

Images

Videos

翻译成中文

关闭取词

27,900 Results

Any time ▾

## Neoadjuvant Chemoradiation with IMRT in Resectable and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4530020>

Induction chemotherapy was used in 32 (82%) of the 39 patients with BLR disease prior to chemoXRT. All resectable patients were treated with chemoXRT alone. Following neoadjuvant treatment, 48(70%) of the 69 patients underwent successful pancreatic resection with 47 (98%) being margin negative ...

Cited by: 18

Author: Jordan Kharofa, Susan Tsai, Tracy Kelly, ...

Publish Year: 2014

## The role of radiotherapy in management of pancreatic cancer

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3397613>

Jul 20, 2011 · However, some patients with borderline resectable pancreatic cancer may benefit from neoadjuvant CRT if the resection can be performed. The assessment of resectability after neoadjuvant CRT is critical to determining the need for surgery, which can have a ...

Cited by: 20

Author: Fen Wang, Parvesh Kumar

Publish Year: 2011

## Neoadjuvant chemoradiation with IMRT in resectable and ...

<https://www.sciencedirect.com/science/article/pii/S0167814014003971>

Induction chemotherapy was used in 32 (82%) of the 39 patients with BLR disease prior to chemoXRT. All resectable patients were treated with chemoXRT alone. Following neoadjuvant treatment, 48 (70%) of the 69 patients underwent successful pancreatic resection with 47 (98%) being margin negative ...

Cited by: 18

Author: Jordan Kharofa, Susan Tsai, Tracy Kelly, ...

Publish Year: 2014

## Meta-analysis comparing upfront surgery with neoadjuvant ...

<https://onlinelibrary.wiley.com/doi/abs/10.1002/bjs.10870>

Apr 30, 2018 · Michiaki Unno, Tatsuo Hata and Fuyuhiko Motoi, Long-term outcome following neoadjuvant therapy for resectable and borderline resectable pancreatic cancer compared to upfront surgery: a meta-analysis of comparative studies by intention-to-treat analysis, Surgery Today, 10.1007/s00595-019-01786-w, 49, 4, (295-299), (2019).

Cited by: 20

Author: E. Versteijne, J. A. Vogel, M. G. Besselin...

Publish Year: 2018

## [PDF] JMAOncology | OriginalInvestigation ...

[https://www.surgonc.org/wp-content/uploads/2019/04/jamaoncology\\_murphy\\_2018\\_oi\\_180015.pdf](https://www.surgonc.org/wp-content/uploads/2019/04/jamaoncology_murphy_2018_oi_180015.pdf)

## Match Overview

1	<b>Crossref</b> 47 words Kathryn E Hitchcock, R Charles Nichols, Christopher G Morris, Debashish Bose, Steven J Hughes, John A Stauffer, Scott A	3%
2	<b>Crossref</b> 10 words Varela, J.E.. "Comparison of laparoscopic and open gastrectomy for gastric cancer", The American Journal of Surgery, 200	1%
3	<b>Internet</b> 10 words crawled on 29-Apr-2019 f6publishing.blob.core.windows.net	1%

**Name of Journal:** *World Journal of Gastrointestinal Surgery*

**Manuscript NO:** 49163

**Manuscript Type:** EDITORIAL

## Optimizing neoadjuvant radiotherapy for resectable and borderline resectable pancreatic cancer using protons

Romaine Charles Nichols, Michael Rutenberg

### Abstract

Approximately 25% of patients diagnosed with pancreatic cancer present with non-metastatic resectable or borderline resectable disease. Unfortunately, the cure rate for these "curable" patients is only in the range of 20%. Local-regional failure rates may exceed 50% after margin-negative, node-negative pancreatectomy, but up to 80% of





All

Images

Videos

翻译成中文

开启取词

29,200 Results

Any time ▾

## Neoadjuvant Chemoradiation with IMRT in Resectable and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4530020>

Induction chemotherapy was used in 32 (82%) of the 39 patients with BLR disease prior to chemoXRT. All resectable patients were treated with chemoXRT alone. Following neoadjuvant treatment, 48 (70%) of the 69 patients underwent successful pancreatic resection with 47 (98%) being margin negative ...

Cited by: 20

Author: Jordan Kharofa, Susan Tsai, Tracy Kelly, ...

Publish Year: 2014

## Neoadjuvant chemoradiation with IMRT in resectable and ...

<https://www.sciencedirect.com/science/article/pii/S0167814014003971>

Induction chemotherapy was used in 32 (82%) of the 39 patients with BLR disease prior to chemoXRT. All resectable patients were treated with chemoXRT alone. Following neoadjuvant treatment, 48 (70%) of the 69 patients underwent successful pancreatic resection with 47 (98%) being margin negative ...

Cited by: 20

Author: Jordan Kharofa, Susan Tsai, Tracy Kelly, ...

Publish Year: 2014

## The role of radiotherapy in management of pancreatic cancer

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3397613>

Jul 20, 2011 · Neoadjuvant radiotherapy. A neoadjuvant treatment strategy in pancreatic cancer may offer several theoretical advantages: 1. Pancreatic cancer is more likely a systemic disease with high incidence of distal and local regional failure. By starting systemic treatment early we may be able to reduce the incidence of distal metastasis and improve survival.

Cited by: 20

Author: Fen Wang, Parvesh Kumar

Publish Year: 2011

## Neoadjuvant vs Adjuvant Therapy for Resectable Pancreatic ...

<https://www.sciencedirect.com/science/article/pii/S1053429613000969>

Furthermore, in a malignancy with such high rates of distant metastasis, investigators are now exploring the feasibility and outcomes of delivering therapy in the neoadjuvant setting, both for clearly resectable as well as borderline resectable tumors.

Cited by: 9

Author: Sarah Hoffs, Nikhil Rao, Ravi Shridhar

Publish Year: 2014

## Meta-analysis comparing upfront surgery with neoadjuvant ...

<https://www.onlinelibrary.wiley.com/doi/full/10.1002/bjs.10870>

Apr 30, 2018 · After neoadjuvant treatment, the R0 resection rate was 85.0 per cent among patients with resectable pancreatic cancer and 88.6 per cent for those with borderline resectable cancer.

Cited by: 24

Author: E. Versteijne, J. A. Vogel, M. G. Besselink...

Publish Year: 2018