

Name of Journal: *World Journal of Biological Chemistry*

ESPS Manuscript NO: 29780

Manuscript Type: Original Article

Basic Study

B-1 cells modulate the murine macrophage response to *Leishmania major* infection

Angelica F Arcanjo, Marise P Nunes, Elias B Silva-Junior, Monique Leandro, Juliana Dutra Barbosa da Rocha, Alexandre Morrot, Debora Decote-Ricardo, Celio Geraldo Freire-de-Lima

Abstract

Match Overview

1	Crossref 39 words Arcanjo, Angélica F., Isabel F. LaRocque-de-Freitas, Juliana Dutra B. Rocha, Daniel Zamith, Ana Caroline Costa-	1%
2	Internet 20 words intimm.oupjournals.org	<1%
3	Internet 13 words crawled on 18-Apr-2016 arca.icict.fiocruz.br	<1%
4	Internet 13 words crawled on 17-Jan-2017 www.jimmunol.org	<1%
5	Internet 13 words crawled on 08-Jul-2010 www.jleukbio.org	<1%
6	Crossref 13 words W.-W. Zhang. "Immunization with a Toll-Like Receptor 7 and/or 8 Agonist Vaccine Adjuvant Increases Protective ...	<1%
7	Crossref 12 words Érico Vinícius de Souza Carmo. "Neutrophils Reduce t ... Parasite Burden in <i>Leishmania</i> (<i>Leishmania</i>) amazonen	<1%



All

Images

News

Shopping

Videos

More ▾

Search tools

About 115,000 results (0.85 seconds)

Scholarly articles for **B-1 cells modulate the innate response of murine macrophages to Leishmania major infection**

... T-bet+ Foxp3- Th1 **cells** are the **major** source of host- ... - Jankovic - Cited by 468

Linkage analysis of IL4 and other chromosome 5q31. 1 ... - Marsh - Cited by 1091

... growth factor- β regulation of immune **responses** - Li - Cited by 1608

Simultaneous transcriptional profiling of *Leishmania major* and its ...

bmcbgenomics.biomedcentral.com/articles/10.1186/s12864-015-2237-2 ▾

by LAL Dillon - 2015 - Cited by 2 - Related articles

Additionally, a CD4+ T helper (Th) type **1 response** by the host leads to ... to identify host and parasite genes that were **modulated** over time, and pathway and RNA-seq was carried out on **mouse macrophages** infected with *L. major* at 4, 24, of the **innate** immune **response** in Th1 cell development following *Leishmania* ...

Avoidance of Innate Immune Mechanisms by the Protozoan Parasite ...

www.ncbi.nlm.nih.gov > ... > Bookshelf ▾ National Center for Biotechnology Information ▾

by DM Mosser - 2000 - Cited by 4 - Related articles

This IL-10 can prevent macrophage responses to IFN- γ allowing the parasites ... points about the activation and **modulation** of **innate immunity** by *Leishmania*: First, **Cellular** SHP-1 activity is increased following **infection**, and subsequent and avirulent *Leishmania donovani* promastigotes in **mouse macrophage** cells.

Mechanisms of Immune Evasion in Leishmaniasis

www.ncbi.nlm.nih.gov/PMC36971... ▾ National Center for Biotechnology Information ▾

by G Gupta - 2013 - Cited by 16 - Related articles

Jan 1, 2014 - **1. IMMUNITY TO LEISHMANIA**. Successful elimination of *Leishmania* ... **Cells of Innate Immunity** ... In *L. major* infection, protection by NKT cells seems to be organ ... state play diverse roles in **modulating** the adaptive immune and affecting In **murine macrophages**, Nrp1 functions as an efflux pump that ...

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 245,000 条结果 (用时 0.74 秒)

Google 学术: B-1 cells modulate the murine macrophage response to Leishmania major infection

... role of CD4+ T cells in the antitumor immune response - Hung - 被引用次数: 1216

... monocyte cytokine responses to Leishmania donovani ... - Reiner - 被引用次数: 102

... of immunity to parasites by T cells and T cell-derived ... - Sher - 被引用次数: 936

Murine Visceral Leishmaniasis: IgM and Polyclonal B-Cell Activation ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: E Deak - 2010 - 被引用次数: 50 - 相关文章

Murine Visceral Leishmaniasis: IgM and Polyclonal B-Cell Activation Lead to Disease ... for infection with L. major [17], while IgG production but not B cell antigen ... Lymph node responses to L. infantum infection. Table 1. B cell activation and ... of this B cell response and mechanisms by which these cells might modulate L.

Leishmania-Infected Macrophages Are Targets of NK Cell-Derived ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: CK Prajeeth - 2011 - 被引用次数: 25 - 相关文章

Previous in vivo studies in murine cutaneous or visceral leishmaniasis showed ... for NK cell activation (58) and released by myeloid DCs in response to Leishmania ... During L. major infection, NK cell-derived IFN- γ was shown to drive the Th1 Qa-1(b)-dependent modulation of dendritic cell and NK cell cross-talk in vivo.

B Cell-Deficient Mice Are Highly Resistant to Leishmania donovani ...

www.jimmunol.org/content/164/7/3681 - 翻译此页

作者: SC Smelt - 2000 - 被引用次数: 158 - 相关文章

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 237,000 条结果 (用时 0.75 秒)

Google 学术: B-1 cells modulate the murine macrophage response to Leishmania major infection

... role of CD4+ T cells in the antitumor immune response - Hung - 被引用次数: 1216

... monocyte cytokine responses to Leishmania donovani ... - Reiner - 被引用次数: 102

... of immunity to parasites by T cells and T cell-derived ... - Sher - 被引用次数: 936

Murine Visceral Leishmaniasis: IgM and Polyclonal B-Cell Activation ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: E Deak - 2010 - 被引用次数: 50 - 相关文章

Murine Visceral Leishmaniasis: IgM and Polyclonal B-Cell Activation Lead to Disease ... for infection with L. major [17], while IgG production but not B cell antigen ... Lymph node responses to L. infantum infection. Table 1. B cell activation and ... of this B cell response and mechanisms by which these cells might modulate L.

Leishmania-Infected Macrophages Are Targets of NK Cell-Derived ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: CK Prajeeth - 2011 - 被引用次数: 25 - 相关文章

Previous in vivo studies in murine cutaneous or visceral leishmaniasis showed ... for NK cell activation (58) and released by myeloid DCs in response to Leishmania ... During L. major infection, NK cell-derived IFN- γ was shown to drive the Th1 Qa-1(b)-dependent modulation of dendritic cell and NK cell cross-talk in vivo.

B Cell-Deficient Mice Are Highly Resistant to Leishmania donovani ...

www.jimmunol.org/content/164/7/3681 - 翻译此页

作者: SC Smelt - 2000 - 被引用次数: 158 - 相关文章

Resolution of Leishmania infection is T cell-dependent, and B lymphocytes have ... This hepatic response