



Unusual cervical foreign body - a neglected thermometer for 5 year





ALL

IMAGES

VIDEOS

12,200 Results

Any time ▼

Neglected foreign body in oesophagus with an unusual ...

https://www.semanticscholar.org/paper/Neglected...

. DOI: 10.1007/BF02907753 Corpus ID: 206824497. Neglected foreign body in oesophagus with anunusual presentation: A case report @article{Verma2006NeglectedFB, title={Neglected foreign body in oesophagus with an unusual presentation: A case report), author={P. Verma and A. Gaur and A. Singhal}, journal={Indian Journal of Otolaryngology and Head and Neck Surgery}, year={2006}, volume={58}, ...

Question Sets and Answers


https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7120678

The larynx is the most common site for ${f foreign\ body}$ aspiration in children younger than 1, while the trachea (C, D) and right mainstem bronchus (A) are the most common sites in older children. The left mainstem bronchus (B) is a less frequent site for foreign body aspiration owing to its acute angle as it enters the lung versus an obtuse angle ...

Cited by: 384

Author: Johanna A.M.G. Tol, Dirk J. Gouma, Claudi...

Publish Year: 2014





Unusual cervical foreign body - a neglected thermometer for 5 years







ALL

IMAGES

VIDEOS

12.200 Results

Any time *

Question Sets and Answers

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7120678

The larynx is the most common site for **foreign body** aspiration in children younger than 1, while the trachea (C, D) and right mainstem bronchus (A) are the most common sites in older children. The left mainstem bronchus (B) is a less frequent site for **foreign body** aspiration owing to its acute angle as it enters the lung versus an obtuse angle ...

Cited by: 384 Author: Johanna A.M.G. Tol, Dirk J. Gouma, Clau...

Publish Year: 2014

Foreign Bodies | RadioGraphics

https://pubs.rsna.org/doi/10.1148/rg.233025137

May 01, 2003 · Foreign Body Ingestions. Foreign body ingestions or insertions are seen in four broad categories of patients: (a) children, (b) mentally handicapped or mentally retarded persons, (c) adults with unusual sexual behavior, and (d) "normal" adults or children with predisposing factors or injurious situational problems. This latter group includes individuals who may abuse drugs or alcohol ...

Bacterial, Fungal, Parasitic, and Viral Myositis

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2493084

a *, the most common cause of myositis in the causative organism category. Of note, S. aureus accounts for 90% of the pyomyositis cases in the tropics and 70% of cases in the developed world. The most common fungal agent is Candida spp., and influenza virus is the most frequent viral cause of myositis identified. The occurrence of parasitic myositis depends on geographic location; cases are ...

Cited by: 319 Author: Nancy F. Crum-Cianflone

Publish Year: 2008

Is a Mask That Covers the Mouth and Nose Free from ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8072811

Apr 20, 2021 - Unlike garments worn over closed skin, masks cover body areas close to the mouth and nose, i.e., body parts that are involved with respiration. Inevitably, this leads not only to a measurable temperature rise [15 , 44 , 85], but also to a severe increase in humidity due to condensation of the exhaled air, which in turn changes the natural ...

Cited by: 4 Author: Kai Kisielinski, Paul Giboni, Andreas Pre...

Publish Year: 2021