

88927_Auto_Edited-check.docx

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 88927

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Combining systemic inflammatory response index and albumin fibrinogen ratio to predict early serious complications and prognosis after respectable gastric cancer

Ren JY *et al.* SIRI and AFR to gastric cancer

Abstract

BACKGROUND

Gastric cancer has a high incidence and fatality rate, and surgery is the preferred course of treatment. Nonetheless, patient survival rates are still low and the incidence of major postoperative complications cannot be disregarded. The systemic inflammatory response, nutritional level, and coagulation status are key factors affecting the postoperative recovery and prognosis of patients with gastric cancer. The systemic inflammatory response index (SIRI) and the albumin fibrinogen ratio (AFR) are 2 valuable comprehensive indicators of the severity and prognosis of systemic inflammation in various medical conditions.

AIM

To assess the clinical importance and prognostic significance of the SIRI scores and the AFR on early postoperative outcomes in patients undergoing radical gastric cancer surgery.

METHODS

We conducted a retrospective analysis of the clinicopathological characteristics and relevant laboratory indices of 568 patients with gastric cancer from January 2018 to December 2019. We calculated and compared two indicators of inflammation and then examined the diagnostic ability combined SIRI and AFR values for early postoperative serious complications. We scored the patients and categorized them into 3 groups based on their SIRI and AFR levels. COX analysis was used to compare three groups of patients the prognostic value of various preoperative SIRI-AFR scores for five-year overall survival (OS) and disease-free survival (DFS).

RESULTS

SIRI-AFR scores were an independent risk factor for prognosis [OS: $P = 0.004$; hazards ratio (HR) = 3.134; DFS: $P < 0.001$; HR = 3.543] and had the highest diagnostic power

(area under the curve: 0.779; 95% confidence interval: 0.737-0.820) for early serious complications in patients with gastric cancer. The tumor-node-metastasis stage ($P = 0.001$), perioperative transfusion ($P = 0.044$), positive carcinoembryonic antigen ($P = 0.014$) findings, and major postoperative complications ($P = 0.011$) were factors associated with prognosis.

CONCLUSION

Preoperative SIRI and AFR values were significantly associated with early postoperative survival and the occurrence of severe complications in patients with gastric cancer.

Key Words: Inflammation; Albumin fibrinogen ratio; Gastric cancer; Complications; Prognosis

Ren JY, Wang D, Zhu LH, Liu S, Yu M, Cai H. Combining systemic inflammatory response index and albumin fibrinogen ratio to predict early serious complications and prognosis after respectable gastric cancer. *World J Gastrointest Oncol* 2024; In press

Core Tip: We conducted a retrospective analysis of the clinicopathological characteristics and relevant laboratory indices of 568 patients with gastric cancer. The aim of this study was to assess the clinical importance and prognostic significance of systemic inflammatory response index (SIRI) combining the albumin fibrinogen ratio (AFR) on early postoperative outcomes in patients with radical gastrectomy. The results demonstrated that the preoperative SIRI and AFR were significantly associated with five-year survival and the occurrence of major complications in gastric cancer patients. We created novel markers in the current study to aid in the early identification and therapy of gastric cancer.

INTRODUCTION

Gastric cancer ranks fifth for morbidity and fourth for fatality for all malignancies and is one of the most prominent diseases worldwide^[1]. Similarly, gastric cancer has made a great contribution to the cancer burden in China. ² Gastric cancer is the second most diagnosed cancer and the third leading cause of cancer-related deaths in China. As a transitioning country, China bears a greater morbidity/mortality and 5-year prevalence rate for gastric cancer compared to most developed countries^[2]. Surgery-based multidisciplinary comprehensive treatment remains the main approach to treating gastric cancer^[3]. An essential course of treatment for non-metastatic gastric cancer is gastrectomy with lymph node dissection^[4].

Despite significant improvements in surgery and anesthetic procedures, postoperative care, and interventional radiology related to stomach cancer gastrectomy has a substantial risk of postoperative complications, such as wound infection, leakage, bleeding, and intestinal obstruction^[5]. And recurrences are common. The rate of postoperative complications following gastric surgery was reported to be 46%^[6]. Thus, these complications may reduce the quality of life, postpone the start of adjuvant treatment, and impede recovery^[7]. Patients with complications are at greater risk of disease recurrence^[8]. Relevant evidence revealed that more than 70% of recurrences and cancer-related mortalities develop within 2 years of surgery, and the recurrence and metastasis of gastric cancer can significantly decrease patient survival rates^[9].

The chronic and sustained inflammation of the gastric cancer not only promotes the occurrence and advancement of gastric cancer^[10], but the inflammatory response stimulates and releases systemic cytokines, which attract the growth of remaining cancer cells and promote postoperative recurrence and metastasis^[8]. Studies revealed that several newly established inflammation-based indicators, including the neutrophil-to-lymphocyte ratio, lymphocyte-to-monocyte ratio, lymphocyte-to-C reactive protein ratio^[11], fibrinogen-to-albumin ratio (FAR)^[12], and systemic inflammatory response index (SIRI)^[13], play an instrumental part in the diagnosis, staging, and prediction of gastric cancer. For example, fibrinogen-neutrophil-to-lymphocyte has served as a prognostic marker for patients with esophageal-gastric junction and superior gastric

cancer after gastrectomy and has shown favorable predictive effects^[14]. Among the above indicators, SIRI, an indicator for assessing a patient's inflammatory status by integrating multiple inflammatory cells into the assessment, has certain advantages and prospects for application. A significant variety of studies have increasingly reported that SIRI values were strongly associated with the prognosis of patients with many different types of tumors^[15-17]. Patients with nasopharyngeal cancer and higher SIRI values had considerably shorter overall survival (OS) compared to those with lower SIRI values^[18]. SIRI values were also found to be a standalone risk prognostic factor in postmenopausal women with breast cancer^[19]. In some solid tumors, such as pancreatic, gastric, and oesophageal malignancies, SIRI values have demonstrated strong predictive performance^[20,21]. The albumin fibrinogen ratio (AFR) is widely used due to its simplicity of measurement, inexpensive nature, and relatively high accuracy^[22]. According to a large retrospective research study of 1196 patients with gastric cancer, serum fibrinogen levels were positively correlated with advanced tumor stage and poor prognosis in patients receiving gastrectomy^[23]. Several studies reported that the FAR or AFR could serve as a point for the clinical prognosis of patients with gastric cancer undergoing first-line chemotherapy^[24], elderly patients with gastric cancer^[25], and in patients with respectable stage II or III gastric cancer^[26-28].

Therefore, to further explore preoperative indicators that can easily and accurately identify the risk of complications in the early post-operative period and prognosis for patients undergoing radical gastrectomy, we propose using both SIRI and AFR values, with the aim of improving the sensitivity of assessing inflammation, nutritional levels and coagulation status and the accuracy and specificity of predicting postoperative outcomes in the short and long-term for patients with gastric cancer.

MATERIALS AND METHODS

Patients and follow-up

This was a retrospective research study on patients at the Gansu Provincial Hospital (Lanzhou, China) with histologically verified gastric cancer from January 2018 to

December 2019. A total of 568 patients met the inclusion criteria. The average age of the study cohort was 60.29 ± 9.79 years and included 442 (77.8%) men and 126 (22.1%) women. The research protocols for the current investigation, which conformed to the principles of the Declaration of Helsinki, received approval from Gansu Provincial Hospital Medical Ethics Committee (ethical consent: 21/10/2022-410). Information was gathered from medical records on sex, age, tumor dimensions, tumor localization, metastatic rate of lymph nodes rate, degree of tumor differentiation, immunohistochemistry results (ki67, P53, and Her2). The process of immunohistochemistry involved staining tissue sections with an antibody specific to the protein of interest, followed by visualization using a chromogenic or fluorescent label^[29]. The p53 expression was defined as positive (Mutant) when staining more than 10% of cancer cell nuclei^[30]. The percentage of cells with Ki67 expression (0%-49%, 50%-74%, 75%-100%) was calculated from the number of malignant cells in the highest labelled field under high magnification (400 \times)^[31]. HER2 expression was evaluated as membrane staining of invasive tumor cells and scored to four classes (0/1+/2+/3+), the expression of grade 3+ or 2+ was defined as positive^[32]. Tumor-node-metastasis (TNM) stage [referring to the American Joint Commission on Cancer (AJCC) gastric cancer TNM staging criteria (eighth edition)], American Society of Anesthesiologists score, surgical approach, extent of resection, duration of surgery, blood loss, periprocedural blood transfusion, length of hospitalization, and duration of postoperative enteral nutrition.

The inclusive criteria were patients: (1) Between 18 and 80 years of age with a clinical diagnosis of preoperative gastric malignancy; (2) With postoperative pathological results confirming primary gastric cancer; and (3) Undergoing D1/D1+/D2 lymph node dissection with radical R0 resection for the first time for radical gastric cancer. The exclusion criteria were: (1) Patients with distant tumor metastasis; (2) Patients with combined hematological diseases, autoimmune diseases, infectious diseases, chronic inflammatory diseases, or liver dysfunction that may affect white blood cells; (3)

Preoperative neoadjuvant therapy (radiotherapy or chemotherapy); (4) The presence of other malignant tumors; and (5) Incomplete data.

The participants in the included studies were followed up by telephone contact, outpatient review, hospitalization, and other methods. The patients were carefully followed up every 3 to 6 months after surgery. Annual follow-up was implemented after 2 years. The follow-up outcomes were OS and disease-free survival (DFS) at 5 years postoperatively. The definition of DFS is the period from diagnosis to any locally recurring disease, distant metastasis, or the last follow-up. OS was defined as the duration between diagnosis and disease-related mortality or the end of the study. The last follow-up was in December 2022.

Laboratory variables and definition of AFR and SIRI index

Relevant indicator levels were assessed in blood samples drawn within a week prior to surgery. Retrospective analysis and data collection from the electronic medical records included additional parameters. SIRI values and AFRs were calculated as the follows: $\text{SIRI} = \text{neutrophil count} \times \text{monocyte count} / \text{lymphocyte count}$; $\text{AFR} = \text{albumin} / \text{fibrinogen}$. Complications occurring in-hospital or within 30 d were categorized as early postoperative complications, and all complications were graded for severity according to the Clavien-Dindo complication grading system^[33], with grade I or II complications were categorized as minor complications, and grade III and higher characterized as major complications. The general post-operative pathology specimen's greatest diameter was used to calculate the tumor size. The primary tumor locations were classified as upper, middle, and lower stomach. accordingly. Differentiation levels were categorized as badly differentiated and moderately/well differentiated.

Statistical analysis

All of the statistical analyses were completed utilizing IBM SPSS for Windows, version 26.0 (IBM Statistics for Windows, version 26, IBM Corporation, Armonk, NY, United States). Categorized data are presented as number (*n*) and percentage (%). For normally

distributed measures, the information is described as the mean \pm SD, and for non-normally distributed continuous variables, it is expressed as the median (interquartile range). Paired groups were compared using either the Mann-Whitney U test or the Student's t -test, depending on the normality of the data distribution. The χ^2 test was used to evaluate categorical group differences. Logistic regression models were employed to identify factors affecting postoperative complications. Receiver operating characteristic (ROC) curves with Youden indices were employed to establish the most favorable cut-off values for each outcome. Youden's index is a global measure of overall diagnostic accuracy and can be used to choose the best cut-point. Its definition is the maximum vertical distance between the ROC curve and the diagonal line^[34]. The values of the area under the curve (AUC) values are supplied with a 95% confidence interval (CI). The hazard ratios (HRs) for disease recurrence or metastasis were calculated applying Cox proportional hazards models. $P < 0.05$ was designated as statistical significance.

RESULTS

Patient characteristics

The flowchart for patient screening is displayed in Figure 1. A total of 568 patients fit the inclusion criteria. No chemotherapy or radiotherapy was administered to any of the patients prior to surgery, and there was no perioperative mortality. This study included 442 men and 126 women with an average age of 60.29 ± 9.79 years (25-87 years). The average body mass index (BMI) ratio prior to surgery for all patients was 22.20 ± 3.37 kg/m². Of the patients, 31.7% ($n = 180$) underwent open surgery, 40.0% ($n = 227$) had a laparoscopic approach, and 28.3% ($n = 161$) underwent robot-assisted surgery. Based on AJCC staging standards, 119 (21.0%) patients were categorized as stage I, 178 (31.3%) had stage II, and 271 (47.7%) had stage III. A mean follow-up time of 45 mo was established for all patients, ranging from 12 to 61 mo. All patients underwent a follow-up assessment.

Postoperative complications

Eighty-nine (15.7%). patients in our study experienced serious complications. The occurrence of early postoperative complications in individuals experiencing radical gastrectomy is shown in Table 1. The complications included a duration of enteral nutrition longer than 2 wk in 26 patients, infection-related complications (incision infection, abdominal infection, pulmonary infection) in 234 patients, an anastomotic fistula in 6 patients, pyloric or intestinal obstruction in 14 patients, thrombosis or embolism in 15 patients, and postoperative shock in 7 patients developed. All resolved after treatment.

The clinical characteristics of the study population are shown in Table 2, along with a comparison of the characteristics and clinical aspects of the 2 groups of patients who had no complications (no) and/or experienced minor complications and those who had major complications. Age ($P = 0.046$), BMI ($P = 0.003$), tumor size ($< 3/\geq 3$ cm) ($P = 0.014$), resection range ($P = 0.019$), perioperative transfusion ($P < 0.001$), and hospital stay ($P < 0.001$) were significantly between the 2 groups (Table 2). For laboratory parameters, lymphocytes ($P < 0.001$), neutrophils ($P < 0.001$), platelets ($P = 0.013$), monocytes ($P = 0.032$), albumin ($P < 0.001$), fibrinogen ($P < 0.001$), carcinoembryonic antigen (CEA) ($P = 0.011$), SIRS ($P < 0.001$), and AFR values ($P < 0.001$) also differed significantly between groups.

Correlations between SIRS, AFR and the clinicopathological characteristics of gastric cancer

Preoperative SIRS scores were related to sex ($P = 0.002$) and resection range ($P = 0.008$) among patients with gastric cancer, as shown in Table 3. AFR was associated with the degree of tumor differentiation ($P = 0.002$) and the duration of enteral nutrition ($P = 0.01$). Both preoperative conditions were related to age, tumor size ($< 3/\geq 3$ cm), TNM stage, perioperative transfusion, carbohydrate antigen 199 (CA199), CEA, amount of bleeding, locoregional recurrence or metastasis ($P < 0.05$). Upon further analysis, among patients under 60 years of age, SIRS levels were lower and AFR levels were higher

compared to patients older than 60 years (SIRI, $P = 0.038$; AFR, $P < 0.001$), and SIRI levels were higher and AFR levels were lower in individuals with a maximum tumor diameter > 3 cm compared to individuals with tumor diameter of 3 cm or less (SIRI, $P < 0.001$; AFR, $P < 0.001$). SIRI values were the highest in patients with stage III disease, and AFRs were the lowest in patients with stage III (SIRI, $P < 0.001$; AFR, $P < 0.001$). SIRI levels were higher and AFRs were lower in the perioperative blood transfusion patients (SIRI, $P < 0.001$; AFR, $P < 0.001$). SIRI scores were higher and AFRs were lower in the CA199 and CEA-positive group of patients (SIRI, $P = 0.023$, $P < 0.001$; AFR, $P = 0.001$, $P < 0.001$). The highest SIRI levels and lowest AFRs levels were observed in the group with intraoperative blood loss > 400 mL (SIRI, $P < 0.001$; AFR, $P < 0.001$). The SIRI levels of patients with gastric cancer with locoregional recurrence or metastasis was noticeably increased ($P < 0.001$), and the AFRs were reduced ($P < 0.001$).

The significance of preoperative SIRI and AFR levels for early serious postoperative complications in respectable gastric cancer

Table 4 lists the outcomes of the univariate and multivariate regression analyses that were executed to establish the odds ratio (OR) values for the complication estimation. The results suggested that high preoperative SIRI values were substantially related to early serious postoperative complications ($P < 0.001$; OR = 1.429; 95%CI: 1.175-1.738), and elevated preoperative AFRs levels were a protective factor against postoperative complications ($P < 0.001$; OR = 0.729; 95%CI: 0.665-0.799). Additionally, the SIRI and AFR components, such as neutrophil count, monocyte count, lymphocyte count, serum albumin, and fibrinogen serum levels, age, BMI, tumor size ($< 3/\geq 3$ cm), resection range, perioperative transfusion, and CEA status ($< 5/\geq 5$ ng/mL) were other variables related to early serious postoperative complications revealed by univariate analysis ($P < 0.05$). Preoperative SIRI values and AFRs remained independent indicators for postoperative complications in multivariable analysis (SIRI: $P = 0.018$; OR = 1.221; 95%CI: 1.031-1.446; AFR: $P < 0.001$; OR = 0.761; 95%CI: 0.693-0.843). Perioperative transfusion ($P = 0.012$; OR = 2.095; 95%CI: 1.179-3.722) was another contributing factor.

Predictive abilities of SIRI and AFR values for postoperative complications

Previous statistical findings concluded that high levels of AFR were a protective parameter for postoperative complications, but a high SIRI value was a risk factor. Thus, to facilitate the calculation of the predictive power of SIRI combined with AFR, we used the FAR in the calculation. ROC curve generation and AUC calculations were used to determine the predictive capability of SIRI and AFR values. The AUC values for SIRI, AFR, and SIRI combined with AFR levels are summarized in Figure 2. The AUC value computed for SIRI was 0.765 (95%CI: 0.714-0.815), 0.743 for AFR (95%CI: 0.689-0.797), and 0.779 for SIRI-AFR (95%CI: 0.737-0.820).

Establishment of SIRI-AFR scores

The patients were grouped based on the appropriate cut-off values for each determinant, established using ROC curves with Youden's index (SIRI: cut-off value: 1.007, sensitivity: 0.966, specificity: 0.532, AFR: cut-off value: 9.849, sensitivity: 0.770, specificity: 0.582). A scoring system was developed according to the SIRI and AFR cut-off values. Patients with a SIRI score of ≥ 1.007 and an AFR of ≤ 9.849 were assigned a SIRI-AFR score of 2, patients with a SIRI score < 1.007 and an AFR > 9.849 were assigned a SIRI-AFR score of 0, and those with a SIRI score of ≥ 1.007 or an AFR of ≤ 9.849 were assigned a SIRI-AFR score of 1. According to the SIRI-AFR system, 219 (38.6%), 224 (39.4%), and 125 (22.0%) patients had scores of 0, 1, and 2, respectively.

Univariate and multivariate Cox regression analyses for OS and DFS

We conducted a COX analysis to investigate the primary variables influencing the prognosis of patients with postoperative gastric cancer. Among patients with gastric cancer, univariate analysis identified that a worse prognosis was profoundly associated with older age (OS: $P = 0.013$; DFS: $P = 0.003$), large tumor size (OS: $P < 0.001$; DFS: $P < 0.001$), later clinical stage (OS: $P < 0.001$; DFS: $P < 0.001$), perioperative transfusion (OS: $P < 0.001$; DFS: $P < 0.001$), positive CA199 (OS: $P = 0.001$; DFS: $P = 0.001$), positive CEA

(OS: $P < 0.001$; DFS: $P = 0.001$), major postoperative complications (OS: $P < 0.001$; DFS: $P < 0.001$), no postoperative adjuvant chemotherapy (OS: $P = 0.003$; DFS: $P = 0.002$), higher SIRI values (OS: $P < 0.001$; DFS: $P < 0.001$), lower AFR values (OS: $P < 0.001$; DFS: $P < 0.001$), and high SIRI-AFR scores (OS: $P < 0.001$; DFS: $P < 0.001$). Multivariate analysis revealed that TNM stage ($P = 0.001$; HR = 5.464, 95%CI: 1.948-15.327), perioperative transfusion ($P = 0.044$; HR = 1.517, 95%CI: 1.011-2.277), positive CEA ($P = 0.014$; HR = 1.605; 95%CI: 1.101-2.338), fibrinogen levels ($P = 0.005$; HR = 1.415, 95%CI: 1.111-1.803), and SIRI-AFR scores ($P = 0.004$; HR = 3.134, 95%CI: 1.445-6.797) were the independently determined prognostic variables for OS (Table 5). Similarly, Cox survival multivariable analysis indicated that TNM stage ($P = 0.001$; HR = 4.071, 95%CI: 1.757-9.435), major postoperative complications ($P = 0.011$; HR = 1.604, 95%CI: 1.115-2.307), albumin levels ($P = 0.044$; HR = 0.959, 95%CI: 0.920-0.999), fibrinogen levels ($P = 0.003$; HR = 1.407, 95%CI: 1.126-1.759), and SIRI-AFR scores ($P < 0.001$; HR = 3.543, 95%CI: 1.844-6.809) were individual prognostic elements for DFS (Table 6). We also observed that SIRI-AFR scores could effectively differentiate patients into 3 distinct risk groups for OS and DFS (Figure 3).

According to the Cox regression model analysis, we performed further subgroup analyses targeting TNM stage, perioperative transfusion, positive CEA, and major postoperative complications, which were several important factors affecting prognosis. The findings demonstrated longer survival in the low SIRI-AFR subgroups with TNM I-II and TNM III (Figures 4A-D). In the subgroups without or with perioperative blood transfusion, patients with low SIRI-AFR levels also had relatively better prognostic ability (Figures 4E-H). Alternatively, patients with lower SIRI-AFR scores exhibited longer survival in the CEA-negative and positive subgroups (Figures 4I-L). Not surprisingly, prognoses were better in the lower SIRI-AFR group than in the high SIRI-AFR group in subgroups with or without postoperative major complications (Figures 4M-P).

DISCUSSION

Gastric cancer is a serious public health issue^[35], and the occurrence of serious complications and recurrence and metastasis after surgery remain difficult problems for clinicians. The development of gastric cancer is a multi-gene, multi-step process and certain key factors may participate in the development of gastric cancer and even infiltration and metastasis at some stages. The systematic inflammatory response and nutritional situation are 2 considerable contributing factors^[36]. SIRI and AFR values are a valuable novel way to evaluate the inflammatory and nutritional conditions of patients. To our knowledge, no studies have examined how SIRI and AFR values in patients who received radical gastric cancer surgery relate to early postoperative serious complications and postoperative survival outcomes. We created novel markers in the current study and evaluated their diagnostic and predictive potential to aid in the early identification and treatment of gastric cancer.

Tumorigenesis involves the establishment of a preneoplastic inflammatory environment^[37]. The Correa sequence, the canonical theory of cancer development in the stomach, indicated that the inflammatory response was an indispensable component of tumor progression^[36]. The epidemiological and clinical investigations provided substantial evidence that inflammation is associated with supporting the growth and dissemination of tumor cells^[38].

The SIRI is unique in reflecting the sophisticated interactions and complementary activity of the major immune cells in the cancer microenvironment. This new metric reflects the state of equilibrium between the immune and inflammatory systems of the host. Neutrophils, as an essential element of the tumor microenvironment, participate in tumor progression *via* multiple mechanisms, and the pathological activation of neutrophils may symbolize the beginning of comprehending the processes behind the reactivation of dormant tumor cells^[39]. Neutrophils produced substances, such as chemokines, cytokines, stromal degrading proteases, and reactive oxygen species, which can alter tumor growth and invasiveness^[40]. Thus, neutrophil physiology at the cellular and molecular levels seems to indicate that their primary function is to facilitate transferential seeding. Neutrophil extracellular traps, shaped by molecularly released

DNA, are intended to capture tumor cells in the circulation^[41]. Such an entanglement of circulating tumor cells may be beneficial to intraluminal survival, adhesion to the endothelium, and extravasation. Monocytes serve as cells bridging innate and adaptive immunity, and can promote cancer immune escape by differentiation into immunomodulatory cells^[42]. They can be involved in the promotion, support, and maintenance of tumor growth by affecting the tumor microenvironment through multiple mechanisms that produce tolerance, angiogenesis, and accelerated tumor cell proliferation^[43]. Lymphocytes played a role in immunologic surveillance and were contributory to identifying and destroying abnormal cells^[44]. Importantly, a biochemical alterations of T cells can modulate cellular activities and promote tumor progression^[45]. Evidence suggests that the magnitude and composition of tumor-infiltrating lymphocytes can affect the survival of oesophageal adenocarcinoma^[46].

With a combination of multiple metrics, the AFR can more accurately assess a patient's inflammatory status, coagulation, and nutritional conditions. Unlike other indicators of inflammation, fibrinogen and albumin levels in the blood are not disturbed by chemotherapy and reflect more accurately reflect the true inflammatory state of the patient after chemotherapy^[47,48]. Abnormal fibrinogen levels can lead to disturbances in the control of normal homeostasis during coagulation. The sedimentation of fibrinogen on cancer cells can form a physical shield to protect cancer cells from recognition and lysis by natural killer cells^[49]. Albumin levels are influenced by nutritional status and metabolism. Hypoalbuminemia can generate immunodeficiency in tumor patients, reducing treatment effectiveness and increasing mortality^[50]. Thus, albumin levels are a recognized prognostic factor for a number of malignancies^[51,52]. Similarly, some research suggested that albumin levels affect the likelihood of postoperative complications^[53] and cancer recurrence^[54].

Mounting data have pointed to the usefulness of SIRS values as a predictor of adverse survival in patients with a range of malignancies, including gastric cancer^[55-57]. In our findings, SIRS values constituted an independently attributable risk for severe postoperative complications in patients with radical gastrectomy. Recently, Schietroma

et al^[58] confirmed that SIRI could predict anastomotic fistulas after total gastrectomy. Similarly, related research has also demonstrated that the AFR could predict the prognoses of patients with pancreatic cancer^[51], gallbladder cancer^[59], and colorectal cancer^[60]. Chen *et al*^[61] reported that the AFR was a distinct risk factor for postoperative delirium in senior gastric cancer patients who underwent laparoscopic gastrectomy, with a cut-off value of 9.95 and an AUC area of 0.614. You *et al*^[25] found that major postoperative complications in senior gastric cancer patients after laparoscopic radical gastrectomy were predicted by the preoperative AFR. The ROC curve's results revealed a cut-off value of 8.49 and an AUC of 0.841. The discrepancy between our cut-off value and the results for the AUC may be due to variations in the data samples and methodological models. Our findings suggested that the AFR was a worthwhile parameter for predicting serious complications and prognosis in patients receiving radical gastrectomy in the early postoperative period. The predictive value of combining SIRI and AFR values for early postoperative serious complications and prognosis of patients undergoing radical gastrectomy was first identified through our study, and suggest that it could be used as a tool to guide cancer treatment strategy decisions.

SIRI and AFR values reflect the complex interactions and synergistic promotion between major immune cells and components of the cancer microenvironment. By integrating risks related to inflammation, coagulation, and nutrition, SIRI and AFR values can deliver a more comprehensive assessment of a patient's overall condition and provide more accurate predictive outcomes. SIRI and AFR values are suitable for frequent testing during follow-up because they have the advantages of easy accessibility, low cost, and good reproducibility. Both the values and the dynamics of SIRI and AFR have the potential to contribute to assessing the efficacy of adjuvant radiotherapy, the selection of suitable patients for specific targeted therapies and immunotherapies, and the monitoring of possible recurrences. In addition, SIRI and AFR values can improve the accuracy and reliability of predictions by continuously learning and updating the models. With the continuous development of medical

technology and the accumulation of clinical data, SIRI values and AFRs can be used to constantly optimize the models to provide more accurate prediction results and better support for patient treatment and rehabilitation.

This investigation had a few limitations. Firstly, the retrospective nature of the study at a single institution restricts its statistical power. Subsequently, we lacked an evaluation of postoperative SIRI and AFR dynamic changes in a relatively large cohort of gastric cancer patients. Therefore, larger multicenter prospective randomized controlled trials are needed to verify our conclusions. Finally, despite the fact that SIRI and AFR values are worthwhile and easily attainable routine blood parameters, the underlying biological and molecular mechanisms that account for their prognostic and predictive nature remain unclear.

CONCLUSION

Overall, the findings of this investigation indicate a significant association between preoperative SIRI and AFR values in individuals with gastric cancer and the occurrence of severe complications, as well as early postoperative survival outcomes. These results may aid surgeons and oncologists in conducting more effective preoperative evaluations and management and developing postoperative monitoring plans for patients with gastric cancer.

ARTICLE HIGHLIGHTS

Research background

Gastric cancer is a serious public health issue, and the occurrence of serious complications and recurrence and metastasis after surgery remain difficult problems for clinicians. Patient survival rates are still low and the incidence of major postoperative complications cannot be disregarded. The systemic inflammatory response, nutritional level, and coagulation status are key factors affecting the postoperative recovery and prognosis of patients with gastric cancer. The systemic inflammatory response index

(SIRI) and the albumin fibrinogen ratio (AFR) are 2 valuable comprehensive indicators of the severity and prognosis of systemic inflammation in various medical conditions.

Research motivation

The aim of this study was to assess the clinical importance and prognostic significance of the SIRI scores and AFR on early postoperative outcomes in patients undergoing radical gastric cancer surgery. These results may aid surgeons and oncologists in conducting more effective preoperative evaluations and management and developing postoperative monitoring plans for patients with gastric cancer.

Research objectives

The object of this study is to assess the clinical importance and prognostic significance of the SIRI scores and the AFR on early postoperative outcomes in patients undergoing radical gastric cancer surgery.

Research methods

We conducted an analysis of the clinicopathological characteristics and relevant laboratory indices of 568 patients with gastric cancer from January 2018 to December 2019. We calculated and compared two indicators of inflammation and then examined the diagnostic ability combined SIRI and AFR values for early postoperative serious complications. We scored the patients and categorized them into 3 groups based on their SIRI and AFR levels.

Research results

SIRI-AFR scores had the highest diagnostic power for early serious complications and were an independent risk factor for prognosis in patients with gastric cancer. Furthermore, the tumor-node-metastasis stage, perioperative transfusion, positive carcinoembryonic antigen findings, and major postoperative complications were factors associated with prognosis. The significant value of the SIRI and AFR for the early severe

postoperative complications and prognosis in gastric cancer patients can provide important insights for the future prevention and treatment of patients. However, we lacked an evaluation of postoperative SIRI and AFR dynamic changes in a relatively large cohort of gastric cancer patients. Therefore, larger multicenter prospective randomized controlled trials are needed to verify our conclusions. And, despite the fact that SIRI and AFR values are worthwhile and easily attainable routine blood parameters, the underlying biological and molecular mechanisms that account for their prognostic and predictive nature remain unclear.

Research conclusions

In this study, we created novel markers in the current study and evaluated their diagnostic and predictive potential to aid in the early identification and treatment of gastric cancer.

Research perspectives

Larger multicenter prospective randomized controlled trials are needed to verify our conclusions. Another, the underlying biological and molecular mechanisms that account for SIRI and AFR values prognostic and predictive nature remain unclear. Further research is needed to elucidate the specific pathways and interactions through which these indicators impact the postoperative outcomes in gastric cancer patients.

4%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|----------|--|-----------------|
| 1 | www.dovepress.com
<small>Internet</small> | 79 words — 1% |
| <hr/> | | |
| 2 | bmccancer.biomedcentral.com
<small>Internet</small> | 35 words — 1% |
| <hr/> | | |
| 3 | Chenjun Ma, Quan Liu, Chengyang Li, Jiwen Cheng, Deyun Liu, Zhanbin Yang, Haibiao Yan, Bo Wu, Yongxian Wu, Jiawen Zhao. "Novel Blood Indicators of Progression and Prognosis in Renal Cell Carcinoma: Red Cell Distribution Width-to-Lymphocyte Ratio and Albumin-to-Fibrinogen Ratio", Journal of Oncology, 2020
<small>Crossref</small> | 26 words — < 1% |
| <hr/> | | |
| 4 | www.science.gov
<small>Internet</small> | 23 words — < 1% |
| <hr/> | | |
| 5 | Yuanjie Qiu, Yan Wang, Nirui Shen, Qingting Wang et al. "Nomograms for Predicting Coexisting Cardiovascular Disease and Prognosis in Chronic Obstructive Pulmonary Disease: A Study Based on NHANES Data", Canadian Respiratory Journal, 2022
<small>Crossref</small> | 22 words — < 1% |
| <hr/> | | |
| 6 | docksci.com
<small>Internet</small> | 17 words — < 1% |

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 15 WORDS
EXCLUDE MATCHES < 15 WORDS