

Prognostic significance of preoperative and postoperative CK19 and CEA mess

全部 图片 视频 新闻 购物 地图 图书

找到约 3,280 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[PDF] [expression of ck-19 and cea mRNA in peripheral blood of gastric ...](#)

[exp-oncology.com.ua/wp/wp-content/uploads/.../882.pdf?...](#)

of **patients with gastric cancer** is poor, with an overall ... clinical and pathological **relevance** of detection of circulating **tumor cells** (CTC) in the **peripheral blood**.

[Detection of carcinoembryonic antigen messenger RNA in blood ...](#)

[https://translational-medicine.biomedcentral.com/.../1479-5876-8-107](#)

31 Oct 2010 ... The existence of circulating **tumor cells** (CTCs) in **peripheral blood** as an ... Blood samples were obtained from 123 **gastric carcinoma patients** at initial ... **CEA mRNA** expression significantly correlated with T stage and **postoperative** recurrence ... Real-time RT-PCR detection of **CEA mRNA levels** at initial ...

[Molecular detection of circulating tumor cells in the peripheral blood ...](#)

[www.ncbi.nlm.nih.gov/pubmed/16736329](#)

Fortunately, the presence of **tumor cells** in **peripheral blood** can be detected ... and is being regarded increasingly as a clinically relevant **prognostic** factor. ... of circulating **tumor cells** and evaluate their relationship to **postoperative** ... 72) of CRC **patients** were positive for hTERT, **CK-19**, CK-20, and **CEA mRNA** respectively.

[Molecular detection and prognostic value of epithelial markers ...](#)

[www.ncbi.nlm.nih.gov/pubmed/24934364](#)

(1)Clinic of Thoracic Oncology, National **Cancer** Institute of Mexico (INcan), ... **mRNA** expression **levels** in **peripheral blood** through three epithelial markers in ... METHODS: Seventy advanced NSCLC **patients** and ten healthy controls ... a **significant** statistical correlation between **levels** of CK-18, **CK-19** and **CEA mRNA**.

[PubPDF - Find Full Text Journal Articles Authored By Hisae linuma](#)

[www.pubpdf.com/search/author/Hisae+linuma](#)

Here, RNA-seq analysis of bone marrow from **gastric cancer patients** was ... as circulating

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 31014

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Prognostic significance of preoperative and postoperative CK19 and CEA messenger RNA levels in the peripheral blood of patients with gastric cardiac cancer

Yu-Feng Qiao, Chuan-Gui Chen, Jie Yue, Ming-Quan Ma, Zhao Ma, Zhen-Tao Yu

Abstract

Match Overview

1	Crossref 46 words Ru, Y., L. Zhang, Q. Chen, S. Gao, G. Wang, Z. Qu, T. Shan, N. Qian, and X. Feng. "Detection and Clinical Significa	1%
2	Crossref 41 words Wu, C. H., S. R. Lin, J. S. Hsieh, F. M. Chen, C. Y. Lu, F. J. Yu, T. L. Cheng, T. J. Huang, S. Y. Huang, and J. Y. Wang.	1%
3	Internet 34 words crawled on 10-Oct-2016 www.spandidos-publications.com	1%
4	Crossref 28 words Qiao, Y.-F., C.-G. Chen, J. Yue, Z. Ma, and Z.-T. Yu. "Clinic al significance of preoperative and postoperative cytoke	1%
5	Internet 28 words crawled on 28-Apr-2016 spandidos-publications.com	1%
6	Internet 27 words crawled on 04-Jul-2016 www.wjgnet.com	1%
7	Crossref 25 words Ming Jian Ge. "Detection of disseminated lung cancer ce lls in regional lymph nodes by assay of CK19 reverse tra	1%
8	Internet 17 words crawled on 01-Jul-2016 content.iospress.com	<1%

Prognostic significance of preoperative and postoperative CK19 and CEA mess

全部 图片 视频 新闻 购物 地图 图书

找到约 4,770 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[PDF] [Molecular detection of disseminated tumor cells in the peripheral ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/.../pdf/DM22-03-281315.pdf>

mRNA for the detection of circulating tumor cells in gastric cancer patients by a ... independent and significant predictor for postoperative recurrence/metastasis ($p = 0.032$). ... Patients with positive CEA mRNA expression in peripheral blood have a significantly higher ... alyzing mRNA molecular markers of hTERT, CK-19..

[Detection of carcinoembryonic antigen messenger RNA in blood ...](#)

<https://translational-medicine.biomedcentral.com/.../1479-5876-8-107>

31 Oct 2010 ... The existence of circulating tumor cells (CTCs) in peripheral blood as an ... Blood samples were obtained from 123 gastric carcinoma patients at initial diagnosis. mRNA ... Real-time RT-PCR detection of CEA mRNA levels at initial diagnosis ... mRNA expression levels of clinical relevance in cancer patients ...

[Circulating tumor cells in gastrointestinal malignancies - Weiss ...](#)

tgc.amegroups.com/article/view/1992/2834

Postoperative positive CTC are associated with decreased PFS and OS (22). ... Multimarker CEA/CK/CD133 mRNA assay was not prognostic in 735 patients post ... Bessa et al. found no significance to perioperative CTC detection via CEA mRNA ... In a recent study (82), pre-operative peripheral blood hTERT mRNA did not ...

[PDF] [Disseminated and circulating tumor cells in ... - HAL-Diderot](#)

<https://hal-univ-diderot.archives-ouvertes.fr/hal.../document>

24 Jun 2012 ... detected in the peripheral blood, which is supposed to have a ... Molecular detection of CTC/DTC relies on the detection of mRNA (by ... Target mRNA (CEA, MUC1, cytokeratins...) ... esophageal-cancer patients who had ≥ 2 CTCs/7.5 ml of blood, ... About half of these studies reported a significant negative ...

[PubPDF - Find Full Text Journal Articles Authored By Hisae Inuma](#)

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 3,650 条结果 (用时 0.96 秒)

Molecular detection of disseminated tumor cells in the peripheral ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/.../pdf/DM22-03-281315.pdf> - 翻译此页

作者: CH Wu - 2006 - 被引用次数: 50 - 相关文章

mRNA for the detection of circulating tumor cells in gastric cancer patients by a ... and significant predictor for postoperative recurrence/metastasis ($p = 0.032$). ... Patients with positive CEA mRNA expression in peripheral blood have a ... analyzing mRNA molecular markers of hTERT, CK-19, ... 60.2 years (range, 34–84). A 4-ml ...

Prognostic Value of Postoperative Detection of Blood Circulating ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: X Bessa - 2003 - 被引用次数: 59 - 相关文章

跳到 PATIENTS AND METHODS - Peripheral blood samples were obtained before and 24 ... targeting carcinoembryonic antigen (CEA) mRNA. ... No patient received preoperative chemoradiation ... and laboratory studies including serum CEA levels every 3 ... prognostic significance of circulating tumor cell detection, ...

Significant Detection of Circulating Cancer Cells in the Blood by ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: K Yamaguchi - 2000 - 被引用次数: 268 - 相关文章

From 52 patients with colorectal cancer, peripheral blood specimens were obtained ... Molecular detection of both CEA and cytokeratin 20 mRNA in mesenteric venous ... at the molecular level in bone marrow and even in circulating blood. of the corresponding clinical data and the postoperative therapeutic schedules.

KRT20 - keratin 20 - iHOP

www.ihop-net.org/UniPub/iHOP/gs/99885.html ▼ 翻译此页

Elevated CK20 or MUC2 mRNA levels were detected in 57 (95%) of the 61 LNs. ... Patients who strongly expressed Gal-3 were positive for CK-7 and negative for CK-20. numbers of cancer cells in gastric cancer are CK18, CK19, CK20, and CEA. ... Expression of cytokeratin 20 in thyroid