

World Journal of *Gastroenterology*

World J Gastroenterol 2017 December 28; 23(48): 8439-8678



**EDITORIAL**

- 8439 Serum levels of angiotensin converting enzyme as a biomarker of liver fibrosis

Miranda AS, Simões e Silva AC

MINIREVIEWS

- 8443 Mechanisms of autophagy activation in endothelial cell and their targeting during normothermic machine liver perfusion

Boteon YL, Laing R, Mergental H, Reynolds GM, Mirza DF, Afford SC, Bhogal RH

ORIGINAL ARTICLE**Basic Study**

- 8452 Human small intestine is capable of restoring barrier function after short ischemic periods

Schellekens DH, Hundscheid IH, Leenarts CA, Grootjans J, Lenaerts K, Buurman WA, Dejong CH, Derikx JP

- 8465 Stable gastric pentadecapeptide BPC 157 in treatment of colitis and ischemia and reperfusion in rats: New insights

Duzel A, Vlajnic J, Antunovic M, Malekinusic D, Vrdoljak B, Samara M, Gojkovic S, Krezic I, Vidovic T, Bilic Z, Knezevic M, Sever M, Lojo N, Kokot A, Kolovrat M, Drmic D, Vukojevic J, Kralj T, Kasnik K, Siroglavic M, Seiwerth S, Sikiric P

- 8489 Exploring pathogenesis of primary biliary cholangitis by proteomics: A pilot study

Deng CW, Wang L, Fei YY, Hu CJ, Yang YJ, Peng LY, Zeng XF, Zhang FC, Li YZ

- 8500 Influence of TBX21 T-1993C variant on autoimmune hepatitis development by Yin-Yang 1 binding

Sun W, Wu HY, Chen S

- 8512 Astragaloside IV inhibits pathological functions of gastric cancer-associated fibroblasts

Wang ZF, Ma DG, Zhu Z, Mu YP, Yang YY, Feng L, Yang H, Liang JQ, Liu YY, Liu L, Lu HW

Retrospective Cohort Study

- 8526 Prevalence and outcomes of pancreatic cystic neoplasms in liver transplant recipients

Liu K, Joshi V, van Camp L, Yang QW, Baars JE, Strasser SI, McCaughan GW, Majumdar A, Saxena P, Kaffes AJ

Retrospective Study

- 8533 Analysis of 12 variants in the development of gastric and colorectal cancers

Cavalcante GC, Amador MA, Ribeiro dos Santos AM, Carvalho DC, Andrade RB, Pereira EE, Fernandes MR, Costa DF, Santos NP, Assumpção PP, Ribeiro dos Santos Á, Santos S

- 8544** Childhood-onset inflammatory bowel diseases associated with mutation of Wiskott-Aldrich syndrome protein gene

Ohya T, Yanagimachi M, Iwasawa K, Umetsu S, Sogo T, Inui A, Fujisawa T, Ito S

- 8553** Comparison of totally laparoscopic total gastrectomy using an endoscopic linear stapler with laparoscopic-assisted total gastrectomy using a circular stapler in patients with gastric cancer: A single-center experience

Gong CS, Kim BS, Kim HS

- 8562** Prognostic significance of pretreatment serum carcinoembryonic antigen levels in gastric cancer with pathological lymph node-negative: A large sample single-center retrospective study

Xiao J, Ye ZS, Wei SH, Zeng Y, Lin ZM, Wang Y, Teng WH, Chen LC

- 8570** Fecal microbiota transplantation induces remission of infantile allergic colitis through gut microbiota re-establishment

Liu SX, Li YH, Dai WK, Li XS, Qiu CZ, Ruan ML, Zou B, Dong C, Liu YH, He JY, Huang ZH, Shu SN

- 8582** Prognostic value of lymph node metastasis in patients with T1-stage colorectal cancer from multiple centers in China

Sun ZQ, Ma S, Zhou QB, Yang SX, Chang Y, Zeng XY, Ren WG, Han FH, Xie X, Zeng FY, Sun XT, Wang GX, Li Z, Zhang ZY, Song JM, Liu JB, Yuan WT

Clinical Trial Study

- 8591** Association between acute pancreatitis and small intestinal bacterial overgrowth assessed by hydrogen breath test

Zhang M, Zhu HM, He F, Li BY, Li XC

Observational Study

- 8597** Endoscopic papillary large balloon dilatation with sphincterotomy is safe and effective for biliary stone removal independent of timing and size of sphincterotomy

Aujla UI, Ladep N, Dwyer L, Hood S, Stern N, Sturgess R

- 8605** Person-centered endoscopy safety checklist: Development, implementation, and evaluation

Dubois H, Schmidt PT, Creutzfeldt J, Bergenmar M

Randomized Clinical Trials

- 8615** Multicenter, randomized study to optimize bowel preparation for colon capsule endoscopy

Kastenberg D, Burch WC, Romeo DP, Kashyap PK, Pound DC, Papageorgiou N, Fernández-Urien Sainz I, Sokach CE, Rex DK

SYSTEMATIC REVIEWS

- 8626** *Fusobacterium's* link to colorectal neoplasia sequenced: A systematic review and future insights

Hussan H, Clinton SK, Roberts K, Bailey MT

- 8651** Psychiatric morbidity after surgery for inflammatory bowel disease: A systematic review

Zangenberg MS, El-Hussuna A

CASE REPORT

- 8660** Stricturing Crohn's disease-like colitis in a patient treated with belatacept

Bozon A, Jeantet G, Rivière B, Funakoshi N, Dufour G, Combes R, Valats JC, Delmas S, Serre JE, Bismuth M, Ramos J, Le Quintrec M, Blanc P, Pineton de Chambrun G

- 8666** Emphysematous pancreatitis associated with penetrating duodenal ulcer

Tana C, Silingardi M, Giamberardino MA, Cipollone F, Meschi T, Schiavone C

- 8671** Infiltrative xanthogranulomatous cholecystitis mimicking aggressive gallbladder carcinoma: A diagnostic and therapeutic dilemma

Nacif LS, Hessheimer AJ, Rodríguez Gómez S, Montironi C, Fondevila C

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Paola Iovino, MD, Associate Professor, Lecturer, Department of Medicine and Surgery, AOU San Giovanni di Dio e Ruggi di Aragona, Salerno 84131, Italy

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7th, 14th, 21st, and 28th each month. The *WJG* Editorial Board consists of 1375 experts in gastroenterology and hepatology from 68 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

World Journal of Gastroenterology (*WJG*) is now indexed in Current Contents[®]/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch[®]), Journal Citation Reports[®], Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2017 edition of Journal Citation Reports[®] cites the 2016 impact factor for *WJG* as 3.365 (5-year impact factor: 3.176), ranking *WJG* as 29th among 79 journals in gastroenterology and hepatology (quartile in category Q2).

FLYLEAF

I-IX Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Yu-Jie Ma*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Ze-Mao Gong*
Proofing Editorial Office Director: *Jin-Lei Wang*

NAME OF JOURNAL
World Journal of Gastroenterology

ISSN
ISSN 1007-9327 (print)
ISSN 2219-2840 (online)

LAUNCH DATE
October 1, 1995

FREQUENCY
Weekly

EDITORS-IN-CHIEF
Damian Garcia-Olmo, MD, PhD, Doctor, Professor, Surgeon, Department of Surgery, Universidad Autonoma de Madrid; Department of General Surgery, Fundacion Jimenez Diaz University Hospital, Madrid 28040, Spain

Stephen C Strom, PhD, Professor, Department of Laboratory Medicine, Division of Pathology, Karolinska Institutet, Stockholm 141-86, Sweden

Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach,

CA 90822, United States

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE
Jin-Lei Wang, Director
Ze-Mao Gong, Vice Director
World Journal of Gastroenterology
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
December 28, 2017

COPYRIGHT
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.f6publishing.com>

Serum levels of angiotensin converting enzyme as a biomarker of liver fibrosis

Aline Silva Miranda, Ana Cristina Simões e Silva

Aline Silva Miranda, Ana Cristina Simões e Silva, Interdisciplinary Laboratory of Medical Investigation, Faculty of Medicine, Federal University of Minas Gerais, Belo Horizonte, MG 30130-100, Brazil

Aline Silva Miranda, Laboratory of Neurobiology, Department of Morphology, Institute of Biological Sciences, Federal University of Minas Gerais, Belo Horizonte, MG 30130-100, Brazil

Ana Cristina Simões e Silva, Department of Pediatrics, Faculty of Medicine, Federal University of Minas Gerais, Belo Horizonte, MG 30130-100, Brazil

ORCID number: Aline Silva Miranda (0000-0003-2811-7924); Ana Cristina Simões e Silva (0000-0001-9222-3882).

Author contributions: Miranda AS and Simões e Silva AC designed the article, performed literature search, and wrote and reviewed the paper.

Supported by CNPq, No. 460334/2014-0; FAPEMIG, No. CDS - PPM-00555-15 (to Simoes e Silva AC); and 2016 NARSAD Young Investigator Grant Awardee from the Brain and Behavior Research Foundation, No. 25414 (to Miranda AS).

Conflict-of-interest statement: Miranda AS and Simões e Silva AC declare no conflict of interest related to this publication.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Ana Cristina Simões e Silva, MD, PhD, Professor, Department of Pediatrics, Interdisciplinary Laboratory of Medical Investigation, Faculty of Medicine, UFMG, Alfredo Balena Avenue, Number 190, 2nd floor, Room #281, Belo

Horizonte, MG 30130-100, Brazil. acssilva@hotmail.com
Telephone: +55-31-34098073
Fax: +55-31-34099770

Received: October 30, 2017
Peer-review started: October 31, 2017
First decision: November 22, 2017
Revised: December 4, 2017
Accepted: December 13, 2017
Article in press: December 13, 2017
Published online: December 28, 2017

Abstract

The renin angiotensin system (RAS) is classically conceived as a circulating hormonal system involved in blood pressure control and hydroelectrolyte balance. The discovery that RAS components are locally expressed in a wide range of organs and tissues, including the liver, pointed to a role for this system in the pathogenesis of several conditions including hepatic fibrosis and cirrhosis. It has been widely reported that the classical RAS axis composed by the angiotensin converting enzyme (ACE)-angiotensin (Ang) II-Ang type 1 (AT1) receptor mediates pro-inflammatory, pro-thrombotic, and pro-fibrotic processes. On the other hand, the alternative axis comprising ACE2-Ang-(1-7)-Mas receptor seems to play a protective role by frequently opposing Ang II action. Chronic hepatitis B (CHB) is one of the leading causes of liver fibrosis, accounting for the death of nearly one million people worldwide. Liver fibrosis is a key factor to determine therapeutic interventions for patients with CHB. However, the establishment of non-invasive and accurate methods to detect reversible stages of liver fibrosis is still a challenge. In an elegant study published in the 36th issue of the *World Journal of Gastroenterology*, Noguchi *et al* showed the predictive value of serum ACE levels in detecting not only advanced stages of liver fibrosis but also initial and

intermediate fibrotic stages. The serum levels of ACE might represent an accurate, non-invasive, widely available, and easy method to evaluate fibrosis related to CHB. Moreover, therapies involving the inhibition of the classical RAS axis components might be promising in the control of CHB-related liver fibrosis.

Key words: Renin angiotensin system; Angiotensin converting enzyme; Angiotensin II; Angiotensin-(1-7); Chronic hepatitis B; Hepatic cirrhosis; Liver fibrosis

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The therapeutic intervention for patients with chronic hepatitis B frequently relies on the pathological classification of liver fibrosis severity in biopsy. The establishment of non-invasive and accurate methods to detect reversible stages of liver fibrosis is still a challenge. High serum levels of angiotensin converting enzyme seem to better predict intermediate liver fibrosis than other classical fibrotic markers. Non-invasive methods to detect intermediate stages of liver fibrosis with very good accuracy may permit the introduction and/or evaluation of treatments during reversible stages of the disease. Further studies are urgently necessary to fully clarify the role of RAS components in liver disease.

Miranda AS, Simões e Silva AC. Serum levels of angiotensin converting enzyme as a biomarker of liver fibrosis. *World J Gastroenterol* 2017; 23(48): 8439-8442 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v23/i48/8439.htm> DOI: <http://dx.doi.org/10.3748/wjg.v23.i48.8439>

INTRODUCTION

The discovery of renin-angiotensin system (RAS) components locally expressed in several organs and tissues, including the kidney, brain, and liver, challenged the classical view of the RAS as a solely circulating hormonal system involved in blood pressure control and hydroelectrolyte balance^[1]. Currently, the RAS is considered a system formed by two opposing axes: the classical axis, which includes the angiotensin-converting enzyme (ACE)-angiotensin (Ang) II-Ang type 1 (AT₁) receptor, and the alternative axis, which comprises the ACE2-Ang-(1-7)-Mas receptor. In general, the classical arm seems to mediate pro-inflammatory, pro-thrombotic, and pro-fibrotic processes, mainly through the activation of AT₁ receptors^[2]. On the other hand, the heptapeptide Ang-(1-7) seems to play a protective role as a biologically active RAS mediator by frequently opposing the action of Ang II *via* stimulation of Mas receptors^[3,4]. In this scenario, over the past decades, an imbalance in the components of the RAS classical and alternative

axes has been implicated in the pathogenesis of a wide range of conditions such as atherosclerosis, obesity, insulin resistance, asthma, renal and liver diseases^[3-5]. Accordingly, many therapeutic strategies have been designed to inhibit ACE-Ang II-AT₁ receptor activity and to stimulate ACE2-Ang-(1-7)-Mas receptor activity^[4,5].

Chronic hepatitis B (CHB) is one of the major causes of liver fibrosis, which, along with hepatitis C, alcohol use, and obesity-related steatohepatitis, has resulted in a significant elevation in the occurrence of cirrhosis and in the mortality of at least 800000 individuals worldwide per year^[6]. Purnak *et al*^[7] detected high serum concentrations of ACE in patients with CHB and they considered this RAS enzyme as a marker of fibrosis. This finding is in line with a more recent study that supported the role of serum ACE level as a noninvasive marker for the prediction of necroinflammatory activity in CHB patients^[8]. Taken together, these studies point to a role of the RAS in liver injury in response to CHB, and pave the way for measuring components of this system as potential predictors of disease evolution.

It is worth mentioning that although the pathophysiology of hepatic fibrosis is still not totally clarified, current opinions have proposed that cirrhosis might be in theory reversible, above all in the compensated stage. Therefore, the evaluation of predictive biomarkers and of novel therapeutic targets is of utmost importance^[9].

STUDY ANALYSIS

In the 36th issue of the *World Journal of Gastroenterology*, Noguchi *et al*^[10] conducted an observational study to investigate the predictive value of serum ACE levels in CHB-associated fibrosis. A total of 100 patients diagnosed with CHB were enrolled in the study and underwent routine liver biopsy. Thirty patients with a history of hypertension, fatty liver, and alcohol abuse were excluded. The degree of hepatic fibrosis in the liver biopsy specimen was evaluated and classified based on the METAVIR score for chronic hepatitis, ranging from F0, no fibrosis, to F4, cirrhosis. The F2 degree (portal fibrosis with few septa) was considered significant liver fibrosis. Additionally, serum levels of ACE and well-known fibrotic markers including the number of platelets (PLT), the aspartate aminotransferase (AST)-to-platelet ratio index (APRI), the Mac-2 binding protein glycosylation isomer (M2BPGi) concentration, and the fibrosis index according to four factors (FIB-4) were also evaluated. For differentiating mild fibrosis (F0-F1) from substantial fibrosis (\geq F2), the 12.8 U/L cut-off value of ACE had a high sensitivity (91.7%) with a good specificity (75%). The receiver-operating characteristic (ROC) curve analysis showed that the area under the curve (AUC) value of ACE serum level measurements was 0.871. The AUC of serum ACE

was bigger than that of other tests for liver fibrosis, including APRI, FIB-4, M2BPGi, and PLT. Importantly, CHB patients in early stages of fibrosis (F0-1) had significantly lower serum levels of ACE than those with significant, advanced fibrosis and cirrhosis (F2-4). The authors concluded that serum levels of ACE might represent an accurate, non-invasive, widely available, and easy method to evaluate fibrosis related to CHB. This conclusion is particularly true for CHB patients without other associated conditions such as fatty liver and/or habitual alcoholic consumption.

The general severity of liver fibrosis influences therapeutic clinical decisions in CHB patients. Serum levels of ACE have been previously evaluated in CHB patients as a potential marker of hepatic fibrosis^[7,8]. For instance, Purnak *et al*^[7] reported higher serum levels of ACE in 22 patients with advanced liver fibrosis compared with 28 patients with mild fibrosis, indicating that the utilization of measurements of serum ACE levels for CHB patients may provide further prognostic data. A more recent study in 54 patients with severe fibrosis showed that serum ACE levels, together with hepatitis B virus deoxyribonucleic acid and serum transaminase levels, might be used as noninvasive markers for predicting necroinflammation in CHB patients^[8]. Even though these previous studies pave the way for the hypothesis that increased serum levels of ACE might be a marker of CHB-associated fibrosis, both included only patients at advanced stages of liver fibrosis for which treatment may not be as efficient as for earlier stages. The study of Noguchi *et al*^[10] was thus designed to overcome this limitation. The authors provide a more accurate classification of the degree of hepatic fibrosis, allowing the predictive value of serum ACE levels to be investigated not only for severe fibrosis but also for initial (F1) and intermediate (F2-F3) stages. Furthermore, Noguchi *et al*^[10] showed that measurement of ACE serum levels was better than other tests for detecting liver fibrosis. It should also be mentioned that the possibility of adopting a non-invasive yet accurate method to detect intermediate stages of liver fibrosis might allow the introduction and/or evaluation of treatments during reversible stages of the disease. Moreover, these findings also suggest that the therapeutic role of inhibitors of the classical RAS axis, such as ACE inhibitors and AT₁ receptor antagonists, should be more deeply investigated for controlling liver fibrosis in CHB patients.

PERSPECTIVES

The establishment of non-invasive and accurate methods to detect reversible stages of liver fibrosis remains a challenge. Many biomarkers have been investigated in this context. However, most of these biomarkers are accurate only for advanced stages of liver disease, when therapies are no longer efficient. Components of the RAS have a clear role in the

pathophysiology of liver disease. Therefore, the measurement of these molecules seems to be very reasonable not only for predicting the progression of liver disease but also for establishing its prognosis, and for being tested as therapeutic targets. In this regard, Noguchi *et al*^[10] clearly showed the ability of serum ACE levels to differentiate initial (F0 and F1) from intermediate (F2-F3) stages of liver fibrosis in CHB patients without other associated conditions. These findings open many different clinical and research approaches. First, measurement of serum ACE levels should be tested in larger samples of patients with CHB and other liver diseases. Second, serum levels of other RAS components, including serum levels of ACE2, Ang II, and Ang-(1-7), should be measured in patients with liver diseases to evaluate these molecules as biomarkers of liver fibrosis and/or disease prognosis. Third, randomized clinical trials with ACE inhibitors or AT₁ receptor antagonists should be conducted in patients in reversible stages of liver diseases. Fourth, the role of components that stimulate the alternative RAS axis (ACE2 activators, Ang-(1-7) analogs, and Mas receptor agonists) should be tested in phase I /phase II studies of liver diseases. In conclusion, the role of RAS components in liver diseases deserves further investigation.

REFERENCES

- 1 Paul M, Poyan Mehr A, Kreutz R. Physiology of local renin-angiotensin systems. *Physiol Rev* 2006; **86**: 747-803 [PMID: 16816138 DOI: 10.1152/physrev.00036.2005]
- 2 Kamo T, Akazawa H, Komuro I. Pleiotropic Effects of Angiotensin II Receptor Signaling in Cardiovascular Homeostasis and Aging. *Int Heart J* 2015; **56**: 249-254 [PMID: 25912907 DOI: 10.1536/ihj.14-429]
- 3 Simões e Silva AC, Silveira KD, Ferreira AJ, Teixeira MM. ACE2, angiotensin-(1-7) and Mas receptor axis in inflammation and fibrosis. *Br J Pharmacol* 2013; **169**: 477-492 [PMID: 23488800 DOI: 10.1111/bph.12159]
- 4 Rodrigues Prestes TR, Rocha NP, Miranda AS, Teixeira AL, Simoes-E-Silva AC. The Anti-Inflammatory Potential of ACE2/Angiotensin-(1-7)/Mas Receptor Axis: Evidence from Basic and Clinical Research. *Curr Drug Targets* 2017; **18**: 1301-1313 [PMID: 27469342 DOI: 10.2174/1389450117666160727142401]
- 5 Simões E Silva AC, Miranda AS, Rocha NP, Teixeira AL. Renin angiotensin system in liver diseases: Friend or foe? *World J Gastroenterol* 2017; **23**: 3396-3406 [PMID: 28596676 DOI: 10.3748/wjg.v23.i19.3396]
- 6 Kochanek KD, Murphy SL, Xu J, Tejada-Vera B. Deaths: Final Data for 2014. *Natl Vital Stat Rep* 2016; **65**: 1-122 [PMID: 27378572]
- 7 Purnak T, Beyazit Y, Oztas E, Yesil Y, Efe C, Torun S, Celik T, Tenlik I, Kurt M, Ozaslan E. Serum angiotensin-converting enzyme level as a marker of fibrosis in patients with chronic hepatitis B. *J Renin Angiotensin Aldosterone Syst* 2012; **13**: 244-249 [PMID: 22277254 DOI: 10.1177/1470320311434241]
- 8 Tay A, Albayrak F, Ozmen S, Albayrak A, Ozden K. Is Serum Angiotensin-Converting Enzyme Level Useful For Determining Necroinflammatory Activity In Chronic Hepatitis B Infection? *Genet Test Mol Biomarkers* 2017; **21**: 102-107 [PMID: 28207324 DOI: 10.1089/gtmb.2016.0067]
- 9 Kim G, Baik SK. Overview and recent trends of systematic reviews and meta-analyses in hepatology. *Clin Mol Hepatol* 2014;

- 20: 137-150 [PMID: 25032179 DOI: 10.3350/cmh.2014.20.2.137]
10 **Noguchi R**, Kaji K, Namisaki T, Moriya K, Kitade M, Takeda K, Kawaratani H, Okura Y, Aihara Y, Furukawa M, Mitoro A,

Yoshiji H. Serum angiotensin-converting enzyme level for evaluating significant fibrosis in chronic hepatitis B. *World J Gastroenterol* 2017; **23**: 6705-6714 [PMID: 29085215 DOI: 10.3748/wjg.v23.i36.6705]

P- Reviewer: Namisaki T **S- Editor:** Gong ZM
L- Editor: Wang TQ **E- Editor:** Ma YJ





Published by **Baishideng Publishing Group Inc**
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>



ISSN 1007-9327

