

## Animal care and use statement

Dear Editor,

**Title:** Optimal timing for the oral administration of Da-Cheng-Qi Decoction based on the pharmacokinetic and pharmacodynamic targeting of the pancreas in rats with acute pancreatitis

**Author:** Yu-mei Zhang, Lin Zhu, Xian-lin Zhao, Huan Chen, Hong-xin Kang, Jian-lei Zhao, Mei-hua Wan, Juan Li, Lv Zhu, Wenfu Tang\*

**Name of Journal:** *World Journal of Gastroenterology*

**ESPS Manuscript NO:** 35253

The animal protocol was designed to minimize pain or discomfort to the animals. The animals were acclimatized to laboratory conditions (22 ±2 °C, 65±10% relative humidity, ad libitum access to water and feeding with laboratory rodent chow (Chengdu, China)) for one week and fasted for 12 h prior to experimentation. Intragastric gavage administration was carried out with conscious animals, using straight gavage needles appropriate for the animal size (280~300 g body weight). All animals were euthanized by 10% chloral hydrate overdose (intraperitoneal injection, 5 mL/kg body weight).

Pro. Wen-fu Tang,



Pancreatic Diseases Research Group,

Department of Integrative Medicine,

West China Hospital, Sichuan University,

Chengdu, China.

Phone: 86 28 85422557

E-mail: tangwf@scu.edu.cn