

80434_Auto_Edited.docx

Name of Journal: *World Journal of Hepatology*

Manuscript NO: 80434

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Liver test abnormalities in asymptomatic and mild COVID-19 patients and their association with viral shedding time

Si-Yi Yu, Jiarong Xie, Junjun Luo, Hongpeng Lu, Lei Xu, Junjie Wang, Xue-Qin Chen

Abstract

BACKGROUND

Asymptomatic infections and mild symptoms are common in patients infected with the Omicron variant, and data on liver test abnormalities are rare.

AIM

¹ We evaluated the clinical characteristics of asymptomatic and mild COVID-19 patients with abnormal liver test results.

METHODS

This retrospective study included 661 Laboratory-confirmed asymptomatic and mild COVID-19 patients who were treated in two makeshift hospitals in Ningbo from April 5 to April 29, 2022. Clinical information and viral shedding time were collected, and univariate and multivariate logistic regression models were performed in statistical analyses.

RESULTS

Of the 661 patients, 83 (12.6%) had liver test abnormalities, and 6 (0.9%) had liver injuries. Abnormal liver tests revealed a reliable correlation with a history of liver

disease ($P<0.001$) and a potential correlation with male sex and obesity ($P<0.05$). Elevated alanine aminotransferase was reliably associated with obesity ($P<0.05$) and a history of liver disease ($P<0.001$). Elevated aspartate transaminase (AST) was reliably correlated with a history of liver disease ($P<0.001$), and potentially correlated with age over 30 years ($P<0.05$). There was a reliable correlation between $AST \geq 2 \times$ the upper limit of normal and a longer viral shedding time, especially in mild cases.

CONCLUSION

Obesity and a history of liver disease are risk factors for liver test abnormalities. Being male and an older age are potential risk factors. Attention should be given to liver tests in asymptomatic and mild COVID-19 patients, which has crucial clinical significance for evaluating the viral shedding time.

INTRODUCTION

1 Coronavirus disease 2019 (COVID-19) is a respiratory disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). This highly contagious disease poses a great threat to global public health. By September 2, 2022, the disease had resulted in 601,189,435 infections and 6,475,346 deaths [1]. In addition to respiratory symptoms and fever, 14-69% of patients with COVID-19 have abnormal liver function tests, mainly manifested by elevations of hypoalbuminemia, gamma-glutamyl transferase, alanine aminotransferase (ALT) and aspartate aminotransferase (AST) [2,3]. Patients with severe diseases are more likely to develop elevated liver tests, suggesting an association between liver injury and the severity of disease [2,4]. Possible mechanisms of COVID-19-related liver injury include direct SARS-CoV-2-induced cytopathic injury to hepatocytes and cholangiocytes, immune dysregulation and hypoxic liver injury, and drug-induced liver injury [5].

1 The Omicron B.1.1.529 (BA.1) variant was first discovered in South Africa on November 9, 2021 [6]. Omicron variants have rapidly replaced delta variants due to their

strong interpersonal infectivity, which has caused a worldwide pandemic. Compared with patients infected with previous variants, those infected with the Omicron variants are younger, have fewer comorbidities, and develop lower severity and mortality, and asymptomatic infections and mild symptoms are more common [7-9]. However, no studies have systematically focused on liver abnormalities in asymptomatic carriers or mild cases. Previous studies have revealed that liver test abnormalities are associated with prolonged hospitalization time and viral shedding time in COVID-19 patients [10,11], indicating a prognostic indicator of poor outcome. At present, no studies have examined the association between liver abnormalities and viral shedding time in asymptomatic and mild COVID-19 patients.

The epidemic prevention policies of different countries are established based on their own political, economic and health conditions, which are in line with the national circumstances and none is superior to others. This study was carried out under the policy of Chinese government, aiming to determine the clinical characteristics of liver test abnormalities in asymptomatic and mild COVID-19 patients infected with the Omicron BA2.2 variant [12] and their association with the viral shedding time. Our research provides suggestions for health policymakers and medical practitioners.

MATERIALS AND METHODS

Participants

In this cross-sectional study, we recruited patients with COVID-19 who were treated in Ningbo Makeshift Hospital and Dapengshan Makeshift Hospital in Ningbo, Zhejiang from April 5, 2022 to April 29, 2022. All patients were transferred from Shanghai. The inclusion criteria were as follows: (1) age over 14 years; (2) complete clinical information; and (3) asymptomatic carriers or mild cases of COVID-19. Asymptomatic carriers were those who had positive etiological tests of COVID-19 but with no clinical symptoms or imaging features of COVID-19. Mild cases were defined as confirmed cases with mild clinical symptoms and no pneumonia manifestation on imaging [13]. The exclusion criteria were as follows: (1) ordinary cases whose pneumonia

manifestation could be seen in imaging, or severe cases with respiratory distress ^[13]; (2) patients who were transferred to designated hospitals for further diagnosis and treatment due to underlying diseases or other reasons. This study complied with the Declaration of Helsinki and was approved by the Ethics Committee of Ningbo First Hospital (No. 2022RS069).

Measures

The clinical information of patients, including age, sex, height, weight, symptoms, medical history (hypertension, diabetes, and liver diseases), and laboratory tests (complete blood count, liver tests, chest CT scans, and nucleic acid tests for COVID-19), were retrospectively collected. Obesity was defined as a body mass index (BMI) ≥ 28 kg/m². Symptoms included fever, fatigue (weakness and muscle soreness), respiratory symptoms (sore throat, dry throat, cough, and chest stuffiness), and gastrointestinal symptoms (nausea, anorexia, and diarrhea). The patients' symptoms were collected from the electronic medical records, and were double-checked on the day of discharge. Liver diseases included chronic hepatitis B, alcoholic/nonalcoholic fatty liver disease (NAFLD) and other liver diseases. NAFLD was defined as hepatic steatosis detected by ultrasound or CT, excluding secondary causes and excessive alcohol consumption (>30 g/day for males, and >20 g/day for females). Hepatitis B was defined as positive serum hepatitis B surface antigen. Complete blood counts, liver and kidney tests, and chest CT scans were performed on the admission day, and nasal swabs for SARS-CoV-2 were tested every two days. All patients received Chinese medicine and symptomatic therapy (if necessary), but not antivirals or monoclonal antibodies.

Real-time reverse transcription PCR was used to detect SARS-CoV-2, and primers targeting the open reading frame 1ab (ORF1ab) and nucleocapsid protein N were used for amplification and detection. The corresponding sequences for ORF1ab were 5'-CCCTGTGGGTTTTACACTTAA-3'(F), 5'-ACGATTGTGCATCAGCTGA-3' (R), and 5'-CY3-CCGTCTGCGGTATGTGGGAAAGGTTATGG-BHQ1-3' (probe), and those for N were 5'-GGGGAAGTTCTCCTGCTAGAAT-3' (F), 5'-CAGACATTTTGCTCTCAAGCTG-30' (R), and 5'-FAM-TTGCTGCTGCTTGACAGATT-

TAMRA-3' (probe). The methods were based on the criteria provided by the National Health Commission of the People's Republic of China [14].

Liver test abnormalities were defined as elevations of the following liver enzymes in serum: ALT>50 U/L (male) or 40 U/L (female), AST>40 U/L (male) or 35 U/L (female), and total bilirubin (TBIL)>17.1 μ mol/L. Liver injury was defined as ALT and/or AST more than 3 \times the upper limit of normal (ULN) and/or TBIL more than 2 \times ULN [15].

Viral shedding was defined as the cycle threshold (Ct) values of both ORF1ab and N greater than 35. The viral shedding time was defined as the duration from the first positive nucleic acid test to the first negative result (two consecutive negatives). Patients were discharged with significantly recovered respiratory symptoms and two consecutive negative nucleic acid tests.

Statistical analysis

Data were analyzed by IBM SPSS (version 26.0). Continuous variables were described by the means \pm standard deviation (SD) or medians and interquartile range (IQR), while categorical variables were described by the frequency and percentage. For continuous variable comparison, independent sample T tests or one-way analysis of variance were used for normally distributed data, and Mann-Whitney U tests or Kruskal-Wallis tests were used for nonnormally distributed data. For categorical variable comparison, chi-square tests or Fisher's exact tests were applied. Ordinal logistic regression analyses were used to identify risk factors associated with liver test abnormalities, and multiple linear regression analyses were used to identify factors associated with the viral shedding time. $P<0.05$ (two-sided) was considered statistically significant. The results of statistical analyses can be re-verified if any party wishes to confirm the credibility.

RESULTS

Study design and participant criteria

The participant recruitment process is shown in Figure 1. In total, 429 patients in Ningbo Makeshift Hospital and 256 patients in Dapengshan Makeshift Hospital were selected according to the inclusion criteria. Excluding 17 ordinary cases and 7 patients who were transferred to designated hospitals (4 cases of pulmonary tuberculosis diagnosed with CT scan, one case of acute asthma attack, one case of acute appendicitis, and one case of persistent abdominal pain), 661 participants were included in the final sample.

As shown in Table 1, the median age was 33 years (IQR 27-44 years), 331 (50.1%) were male, and the median BMI was 22.8 kg/m² (IQR 20.7-25.2). A total of 130 (19.7%) had underlying diseases, of whom 57 (8.6%) had liver diseases. 45 (6.8%) had NAFLD, 11 (1.7%) had hepatitis B, and 1 (0.2%) had both. A total of 425 (64.3%) patients developed a fever, and 619 (93.6%) patients had mild cases. Eighty-three (12.6%) patients had liver test abnormalities and 6 (0.9%) had liver injuries. The number of patients with elevations in ALT, AST and TBIL was 53 (8.0%), 61 (9.2%), and 4 (0.6%), respectively, with a majority of mild liver test abnormalities (Supplementary Table 1). Some patients reported medicines taken before admission, including traditional Chinese medicine (382, 57.8%), nonsteroidal anti-inflammatory drugs (NSAIDs) (89, 13.5%), and other medicines for cold (91, 13.8%).

Clinical features of COVID-19 patients with liver test abnormalities

As shown by a univariate logistic regression model, liver test abnormalities were significantly associated with male sex [odds ratio (OR) 1.699, 95% confidence interval (CI) 1.061-2.721, $P=0.027$], obesity (OR 2.707, 95%CI 1.384-5.291, $P=0.004$) and a history of liver disease (OR 2.707, 95%CI 1.384-5.291, $P=0.004$) but not with age, disease type, clinical manifestations or medication history (Figure 2). The correlated factors in the univariate model were taken as key factors. A multivariate logistic regression model for key factors indicated that liver test abnormalities were only associated with a history of liver disease (OR 8.004, 95%CI 4.319-14.835, $P<0.001$). A multivariate logistic regression model for all factors showed that liver test abnormalities were significantly associated with the age of 30-49 years (compared with age 14-30 years, OR 1.970, 95%CI 1.073-

3.618, $P=0.029$), male sex (OR 1.728, 95%CI 1.005-2.971, $P = 0.048$), and a history of liver disease (OR 8.265, 95%CI 4.315-15.831, $P<0.001$). Based on the above models, liver test abnormalities had a reliable correlation with a history of liver disease, and a potential correlation with male sex and obesity.

As revealed by all three models in Supplementary Figure 1, elevated ALT had a firm correlation with obesity (univariate logistic regression: OR 3.82, 95%CI 1.84-7.94, $P<0.001$; multivariate logistic regression for key factors: OR 2.57, 95%CI 1.15-5.70, $P=0.021$; multivariate logistic regression for all factors: OR 3.31, 95%CI 1.37-7.98, $P=0.009$) and a history of liver disease (univariate logistic regression: OR 9.54, 95%CI 5.01-18.16, $P<0.001$; multivariate logistic regression for key factors: OR 8.30, 95%CI 4.21-16.36, $P<0.001$; multivariate logistic regression for all factors: OR 9.05, 95%CI 4.31-19.01, $P<0.001$). Therefore, ALT elevation was reliably correlated with obesity and a history of liver disease.

A univariate logistic regression model indicated that elevated AST was associated with a history of liver disease (OR 8.18, 95%CI 4.40-15.24, $P<0.001$, Supplementary Figure 2). A multivariate logistic regression model on all factors showed that AST elevation was associated with older age (age of 30-49 years vs. age of 14-29 years: OR 2.48, 95%CI 1.18-5.20, $P=0.016$; age over 50 years vs. age of 14-29 years: OR 2.81, 95%CI 1.14-6.97, $P=0.025$) and a history of liver disease (OR 7.84, 95%CI 3.89-15.78, $P<0.001$) (Supplementary Figure 2). Therefore, the AST elevation had a reliable correlation with a history of liver disease, and a potential correlation with age over 30 years.

In summary, obesity and a history of liver disease were risk factors for abnormal liver test results, and male sex and age over 30 years were potential risk factors. These findings are of great clinical significance for evaluating high-risk subjects with liver abnormalities among asymptomatic and mild COVID-19 patients.

Association between viral shedding time and liver test abnormalities

The association between the viral shedding time and abnormal liver test results was analyzed in 657 participants after excluding 4 subjects with missing data. As shown in Table 2, the viral shedding time of patients with $AST \geq 2 \times ULN$ were significantly

prolonged ($<2\times\text{ULN}$ vs. $\geq 2\times\text{ULN}$, viral shedding time: 10.51 ± 2.76 d vs. 12.40 ± 3.57 d, $P=0.033$). As suggested by three models in Figure 3, $\text{AST}\geq 2\times\text{ULN}$ had a reliable correlation with the viral shedding time (univariate linear regression: coefficient 1.893, 95%CI 0.158-3.628, $P=0.033$; multivariate linear regression of key factors: coefficient 1.862, 95%CI 0.140-3.584, $P=0.034$; multivariate linear regression for all factors: coefficient 2.778, 95%CI 0.615-4.942, $P=0.012$). In contrast, there was no significant correlation between ALT elevation and the viral shedding time. In addition, mild cases or female sex were associated with a longer viral shedding time (Table 2, Figure 3).

The correlations between the AST elevation or disease types and the viral shedding time were further analyzed. As shown in Supplementary Table 2, the viral shedding time of mild cases with $\text{AST}\geq 2\times\text{ULN}$ was significantly prolonged ($\text{AST}\geq 2\times\text{ULN}$ & mild cases vs. $\text{AST}<2\times\text{ULN}$ & mild cases vs. $\text{AST}<2\times\text{ULN}$ & asymptomatic carriers: 12.40 ± 3.57 d vs. 10.59 ± 2.71 d vs. 9.26 ± 3.16 d, $P=0.001$). Therefore, it is suggested that great attention should be given to the liver test results of asymptomatic and mild COVID-19 patients, which has crucial clinical significance for predicting the viral shedding time.

DISCUSSION

³ To the best of our knowledge, this is the first clinical study focusing on liver test abnormalities in asymptomatic and mild COVID-19 patients. In the current epidemic of the Omicron BA2.2 variant, 12.6% of COVID-19 patients developed liver test abnormalities, and 0.9% developed liver injuries. The majority of the liver test abnormalities were mild. The ratio of abnormal liver tests in our study was lower than previous data of 14-69% [23]. One possible reason for this was that our participants were all asymptomatic carriers or mild cases, among whom abnormal liver test results were uncommon. Previous studies revealed normal ALT and AST levels in asymptomatic patients, which were significantly lower than those with symptomatic infection [16]. In contrast, our study showed no differences in liver tests between asymptomatic carriers

and mild cases. The probable reason was that all of the symptomatic patients in our study were mild cases, which weakened the difference between the two groups.

As shown in our study, the risk factors for liver test abnormalities in asymptomatic and mild COVID-19 patients included obesity and a history of liver disease, and the potential risk factors included male sex and age over 30 years. Previous studies suggested that preexisting chronic liver diseases, such as NAFLD, chronic hepatitis B, and alcoholic liver disease, were high-risk factors for liver injury in COVID-19 [17-19]. Consistent with these findings, our study verified a strong correlation between abnormal liver test results and a history of liver disease. Virus-induced cytokine storms, impaired mitochondrial activity, or endoplasmic reticulum stress may aggravate the preexisting liver disease, which further leads to the progression of liver injury [20]. In addition, liver test abnormalities were more common in males, which was similar to a previous study [21]. The higher estrogen level in females may play a protective role [22]. In addition, elevated ALT was strongly associated with obesity. This may be due to a high ratio of NAFLD patients among obese individuals, and fatty liver diseases can result in elevated ALT levels [18]. There was a potential correlation between elevated AST and age over 30 years. Since AST is known to reflect the injury of organs, such as myocardium and skeletal muscle, these organic injuries may occur more frequently in elderly patients with COVID-19.

Medications for COVID-19 treatment, such as antibiotics and antiviral medications (e.g., lopinavir/ritonavir) have been reported to cause liver injury [10]. This study focused on the liver test features of COVID-19 patients in the early stage of the disease. These patients had rarely taken antibiotics or antiviral medications, while some had taken Chinese medicine, NSAIDs or other cold medicines for a short period. Importantly, no association was found between these medications and liver test abnormalities in COVID-19 patients.

As reported in hospitalized COVID-19 patients, abnormal liver test results at admission were associated with a prolonged hospitalization time and viral shedding time [10,11]. Early elevation of AST was closely related to mortality, suggesting that it

might be a predictor of poor prognosis for COVID-19 [23]. Similarly, we also found a reliable association between the increase in AST and a prolonged viral shedding time in asymptomatic and mild COVID-19 patients. We speculate that the lower immunity of patients with elevated AST levels might affect the viral clearance. The viral shedding time was even longer in mild cases with elevated AST. Similar results have been previously revealed in which the viral shedding duration was longer in symptomatic patients than in asymptomatic patients [16,24]. Lymphocyte-mediated immune responses may play a vital role [15].

Our study has two advantages. First, unlike studies concerning severe cases [2,4,15,23], we focused on the association between liver test results and viral shedding time, especially in patients infected with the Omicron BA2.2 variant, with a relatively high proportion of asymptomatic carriers and mild cases. The viral shedding time for patients with elevated AST were significantly longer, especially in mild patients. This provides crucial evidence for identifying high-risk patients with a prolonged viral shedding time. Second, we analyzed the risk factors for liver test abnormalities. Male sex, older age, obesity, and a history of liver disease may increase the risk of liver abnormalities, which can be used to identify high-risk patients with abnormal liver test results.

One limitation of our research is that we only measured the levels of ALT, AST, and TBIL, but failed to obtain other liver test results, such as γ -glutamyl transpeptidase and alkaline phosphatase, and we did not monitor the dynamic alterations of the liver tests. The absence of these indicators may have led to an incomplete assessment of liver abnormalities. Another limitation is that we only recruited participants from two makeshift hospitals in Ningbo, but not from multiple regions. This may have led to a limited sample size and underrepresentation of the sample.

CONCLUSION

1 Coronavirus disease 2019 (COVID-19) is a respiratory disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). This highly contagious

disease poses a great threat to global public health. By September 2, 2022, the disease had resulted in 601,189,435 infections and 6,475,346 deaths [1]. In addition to respiratory symptoms and fever, 14-69% of patients with COVID-19 have abnormal liver function tests, mainly manifested by elevations of hypoalbuminemia, gamma-glutamyl transferase, alanine aminotransferase (ALT) and aspartate aminotransferase (AST) [2,3]. Patients with severe diseases are more likely to develop elevated liver tests, suggesting an association between liver injury and the severity of disease [2,4]. Possible mechanisms of COVID-19-related liver injury include direct SARS-CoV-2-induced cytopathic injury to hepatocytes and cholangiocytes, immune dysregulation and hypoxic liver injury, and drug-induced liver injury [5].

¹ The Omicron B.1.1.529 (BA.1) variant was first discovered in South Africa on November 9, 2021 [6]. Omicron variants have rapidly replaced delta variants due to their strong interpersonal infectivity, which has caused a worldwide pandemic. Compared with patients infected with previous variants, those infected with the Omicron variants are younger, have fewer comorbidities, and develop lower severity and mortality, and asymptomatic infections and mild symptoms are more common [7-9]. However, no studies have systematically focused on liver abnormalities in asymptomatic carriers or mild cases. Previous studies have revealed that liver test abnormalities are associated with prolonged hospitalization time and viral shedding time in COVID-19 patients [10,11], indicating a prognostic indicator of poor outcome. At present, no studies have examined the association between liver abnormalities and viral shedding time in asymptomatic and mild COVID-19 patients.

The epidemic prevention policies of different countries are established based on their own political, economic and health conditions, which are in line with the national circumstances and none is superior to others. This study was carried out under the policy of Chinese government, aiming to determine the clinical characteristics of liver test abnormalities in asymptomatic and mild COVID-19 patients infected with the Omicron BA2.2 variant [12] and their association with the viral shedding time. Our research provides suggestions for health policymakers and medical practitioners.

ARTICLE HIGHLIGHTS

Research background

Data on liver test abnormalities in asymptomatic and mild COVID-19 patients are rare.

Research motivation

This study ¹ evaluated the clinical characteristics of asymptomatic and mild COVID-19 patients with abnormal liver test results.

Research objectives

We aimed to determine the liver test abnormalities in asymptomatic and mild COVID-19 patients and their association with the viral shedding time, providing suggestions for health policymakers and medical practitioners.

Research methods

Clinical information and viral shedding time were collected retrospectively from 661 Laboratory-confirmed asymptomatic and mild COVID-19 patients. Univariate and multivariate logistic regression models were performed in statistical analyses.

Research results

Elevated alanine aminotransferase was associated with obesity and a history of liver disease. Elevated aspartate transaminase (AST) was correlated with a history of liver disease age over 30 years. There was a correlation between $AST \geq 2 \times$ the upper limit of normal and a longer viral shedding time.

Research conclusions

Obesity and a history of liver disease are risk factors for liver test abnormalities. Liver test abnormality in asymptomatic and mild COVID-19 patients has clinical correlation with the viral shedding time.

Research perspectives

Attention should be given to liver tests in asymptomatic and mild COVID-19 patients, which has crucial clinical significance for evaluating the viral shedding time.

ORIGINALITY REPORT

4%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------|
| <div style="background-color: red; color: white; padding: 5px; display: inline-block; width: 30px; height: 30px; line-height: 30px; text-align: center;">1</div> | www.ncbi.nlm.nih.gov
<small>Internet</small> | 135 words — 3% |
| <hr/> | | |
| <div style="background-color: magenta; color: white; padding: 5px; display: inline-block; width: 30px; height: 30px; line-height: 30px; text-align: center;">2</div> | link.springer.com
<small>Internet</small> | 16 words — < 1% |
| <hr/> | | |
| <div style="background-color: purple; color: white; padding: 5px; display: inline-block; width: 30px; height: 30px; line-height: 30px; text-align: center;">3</div> | Li Qianqian, Chen Jing, Sun Kai, Song Ruixin et al.
"Safety and Immunogenicity of a Booster SARS-
Cov-2 Vaccination in Patients with Chronic Liver Disease", Cold
Spring Harbor Laboratory, 2022
<small>Crossref Posted Content</small> | 13 words — < 1% |
| <hr/> | | |
| <div style="background-color: teal; color: white; padding: 5px; display: inline-block; width: 30px; height: 30px; line-height: 30px; text-align: center;">4</div> | test.dovepress.com
<small>Internet</small> | 12 words — < 1% |

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES

EXCLUDE MATCHES

OFF

< 12 WORDS