



67383_Auto_Edited.docx

Quotes Excluded
Bibliography Excluded5%
SIMILAR**Name of Journal:** *World Journal of Clinical Cases***Manuscript NO:** 67383**Manuscript Type:** ORIGINAL ARTICLE*Observational Study***Real-world data on the infliximab biosimilar CT-P13 (Remsima®) in inflammatory bowel disease**

Jose María Huguet, Xavier Cortés, Marta Maia Bosca-Watts, Marian Aguas, Nuria Maroto, Lidia Martí, Cirilo Amorós, Jose María Paredes

Abstract

BACKGROUND

Match Overview

1	Internet 43 words crawled on 05-Oct-2020 www.researchgate.net	1%
2	Internet 38 words crawled on 02-Nov-2020 academic.oup.com	1%
3	Crossref 36 words Jacob E. Ollech, Inessa Normatov, Noam Peleg, Sushila R. D alal et al. "Efficacy of biosimilar infliximab CT-P13 among inj ...	1%
4	Internet 28 words crawled on 20-Sep-2019 link.springer.com	1%
5	Internet 25 words crawled on 28-Mar-2021 pubmed.ncbi.nlm.nih.gov	1%



ALL

IMAGES

VIDEOS

24,900 Results

Any time ▼

CT-P13 (Inflectra™, Remsima™) monitoring in patients with ...

<https://pubmed.ncbi.nlm.nih.gov/27435444>

The approval of **infliximab biosimilars Remsima™** and Inflectra™ (CT-P13) for patients with **inflammatory bowel disease** (IBD) is a promising step to reduce treatment costs. Since monitoring of Remicade™ serum trough levels and anti-Remicade™ immunogenicity hold an important significance in treatment modalities, no data about monitoring of drug serum trough ...

Cited by: 7**Author:** Kornelius Schulze, Nadine Koppka, F...**Publish Year:** 2016

Switching from Remicade® to Remsima® is well Tolerated ...

<https://pubmed.ncbi.nlm.nih.gov/27660339>

Abstract. Background and aims: A **biosimilar** version of **infliximab** [CT-P13/Remsima®] recently entered the European market. The clinical **data** on its use in **inflammatory bowel disease** [IBD] are sparse, especially on switching from the originator Remicade®. In this study, we aimed to prospectively investigate the feasibility, safety and immunogenicity of switching from **Remicade** to **Remsima** in a **real-life** ...

Cited by: 97**Author:** Lydia C. T. Buer, Lydia C. T. Buer, Bjør...**Publish Year:** 2016

Clinical experience with infliximab biosimilar Remsima (CT ...

<https://www.ncbi.nlm.nih.gov/pubmed/27134662>

Based on current **data**, **CT-P13** seems to be efficacious and generally well tolerated in **IBD** especially in patients who are naïve to biological therapy. Knowledge with regard to interchangeability between **CT-P13** and the originator **infliximab** is however, still rather sparse and more **data** are desired.

Cited by: 21**Author:** Jørgen Jahnsen, Jørgen Jahnsen**Publish Year:** 2016

Post-marketing analysis for biosimilar CT-P13 in ...

<https://pubmed.ncbi.nlm.nih.gov/33450057>

Background and aim: **CT-P13**, an **infliximab** (IFX) biosimilar, was approved for treatment of inflammatory bowel disease. However, no comparison with the originator IFX in this indication has

Real-world data on the infliximab biosimilar CT-P13 (Remsima®) in

 ALL IMAGES VIDEOS

26,500 Results

Any time ▼

Infliximab biosimilar CT-P13 is interchangeable with its ...

<https://pubmed.ncbi.nlm.nih.gov/31422647>

Background/aims: An interim analysis of post-marketing surveillance of **CT-P13**, an **infliximab biosimilar**, was performed to evaluate its safety and efficacy in Japanese patients with **inflammatory bowel disease**. Methods: Patients were prospectively enrolled between November 2014 and March 2017, after the launch of **CT-P13** in Japan, and case report forms of patients followed for at least 4 ...

Cited by: 4

Author: Tomoo Nakagawa, Taku Kobayashi, Kiyo...

Publish Year: 2019

CT-P13 (Infliximab TM, Remsima TM) was initiated in patients with

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

Show language tabs

7,780 Results

Any time ▾

[Infliximab biosimilar CT-P13 is interchangeable with its ...](https://pubmed.ncbi.nlm.nih.gov/31422647)

<https://pubmed.ncbi.nlm.nih.gov/31422647>

Background/aims: An interim analysis of post-marketing surveillance of **CT-P13**, an **infliximab biosimilar**, was performed to evaluate its safety and efficacy in Japanese patients with **inflammatory bowel disease**. Methods: Patients were prospectively enrolled between November 2014 and March 2017, after the launch of **CT-P13** in Japan, and case report forms of patients followed for at least 4 months ...

Cited by: 4

Author: Tomoo Nakagawa, Taku Kobayashi, Kiyohir...

Publish Year: 2019

[CT-P13 \(Inflectra™, Remsima™\) monitoring in patients with ...](https://pubmed.ncbi.nlm.nih.gov/27435444)

<https://pubmed.ncbi.nlm.nih.gov/27435444>

The approval of **infliximab biosimilars Remsima™** and **Inflectra™** (CT-P13) for patients with **inflammatory bowel disease** (IBD) is a promising step to reduce treatment costs. Since monitoring of **Remicade™** serum trough levels and anti-**Remicade™** immunogenicity hold an important significance in treatment modalities, no data about monitoring of drug serum trough levels or anti-drug antibody ...

Cited by: 9

Author: Kornelius Schulze, Nadine Koppka, Frederik...

Publish Year: 2016

[Clinical experience with infliximab biosimilar Remsima \(CT ...](https://www.ncbi.nlm.nih.gov/pubmed/27134662)

<https://www.ncbi.nlm.nih.gov/pubmed/27134662>

Based on current **data**, **CT-P13** seems to be efficacious and generally well tolerated in **IBD** especially in patients who are naïve to biological therapy. Knowledge with regard to interchangeability between **CT-P13** and the originator **infliximab** is however, still rather sparse and more **data** are desired.

Cited by: 21

Author: Jørgen Jahnsen, Jørgen Jahnsen

Publish Year: 2016

[Switching from infliximab to biosimilar in inflammatory ...](https://journals.sagepub.com/doi/full/10.1177/1756284819842748)

<https://journals.sagepub.com/doi/full/10.1177/1756284819842748>

Apr 15, 2019 · In 2017, the extension study of **PLANETRA** and **PLANETAS** showed that switching from **RP** to its **biosimilar CT-P13** is possible without negative effects on safety or efficacy in patients with AS.

65,66 Furthermore, switching from the **IFX RP** to **CT-P13** after 1 year of **IFX RP** treatment showed continued comparable efficacy, immunogenicity and safety, to maintenance of **CT-P13** treatment during the second year of the treatment. 66 In **PLANETRA**...