



23188-Review
BY YUJI SHINDO

Quotes Excluded
Bibliography Excluded

8%
SIMILAR

Name of journal: *World Journal of Gastrointestinal Endoscopy*

ESPS Manuscript NO: 23188

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Risk factors for postoperative bleeding after gastric endoscopic submucosal dissection in patients under antithrombotics

Yuji Shindo, Satohiro Matsumoto, Hiroyuki Miyatani, Yukio Yoshida, Hirosato Mashima

Match Overview

1	CrossCheck 84 words Sanomura, Yoji, Shiro Oka, Shinji Tanaka, Norifumi Numata, Makoto Higashiyama, Hiroyuki Kanao, Shigeto Yoshida	2%
2	CrossCheck 48 words Nakamura, Munetaka, Jun Nishikawa, Kouichi Hamabe, Junichi Nishimura, Masaaki Satake, Atsushi Goto, Shu Ki	1%
3	CrossCheck 47 words Fujimoto, Kazuma, Mitsuhiro Fujishiro, Mototsugu Kato, Kazuhide Higuchi, Ryulchi Iwakiri, Choitsu Sakamoto, S	1%
4	CrossCheck 46 words Matsumoto, Satohiro. "Selection of appropriate endoscopic therapies for duodenal tumors: An open-label study ..."	1%
5	Internet 28 words crawled on 15-Jan-2016 www.wjgnet.com	1%
6	CrossCheck 27 words Sanomura, Yoji, Shiro Oka, Shinji Tanaka, Makoto Higashiyama, Shigeto Yoshida, Koji Arihiro, Fumio Shimamoto	1%



Risk factors for postoperative bleeding after gastric endoscopic submucosal



学术搜索

找到约 161 条结果 (用时0.13秒)

我的著作引用情况



文章

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

我的图书馆

Antithrombotic drugs are risk factors for delayed postoperative bleeding after endoscopic submucosal dissection for gastric neoplasmsR Koh, K Hirasawa, S Yahara, H Oka... - ... *endoscopy*, 2013 - Elsevier... **postoperative bleeding**, a history of cardiovascular disease was the sole **factor** associated with ...CI, 1.596-6.587; P = .001) revealed them to be independent **risk factors** associated with ... Thedelayed **postoperative bleeding** rate was highest (11.6%) in lesions with a specimen size ...

被引用次数: 29 相关文章 所有 5 个版本 引用 保存

[PDF] Risk factors for bleeding after endoscopic submucosal dissection for gastric lesionsresearchgate.net 中的 **[PDF]**

Y Tsuji, K Ohata, T Ito, H Chiba, T Ohya... - World J ..., 2010 - researchgate.net

... in our study showed that use of these medicines was an independent **risk factor** for post ... Themedian interval between ESD and **postoperative** hemorrhage in L lesions was 1 d ... In conclusion,**risk factors** for **bleeding after gastric** ESD included beginner operators coagulating the ...

被引用次数: 74 相关文章 所有 9 个版本 引用 保存 更多

Risk factors for bleeding after endoscopic submucosal dissection of gastric epithelial neoplasm researchgate.net 中的 **[PDF]**M Higashiyama, S Oka, S Tanaka... - ... *Endoscopy*, 2011 - Wiley Online Library... Table 4. **Risk factors** for **bleeding after** ESD of **gastric** epithelial neoplasm in relation tosecond-look ... EMR) for 504 **gastric** tumors and studied the results, reporting that the only **risk factor**for **bleeding** ... for separation and therefore requires extra time, but in our high-risk cases, that is ...

被引用次数: 50 相关文章 所有 5 个版本 引用 保存

[HTML] The postoperative bleeding rate and its risk factors in patients on antithrombotic therapy biomedcentral.com 中的 **[HTML]****who undergo gastric endoscopic submucosal dissection**

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

 包括专利 包含引用



Risk factors for postoperative bleeding after gastric endoscopic submucos



全部 图片 新闻 视频 更多 搜索工具

找到约 19,700 条结果 (用时 0.77 秒)

Google 学术: Risk factors for postoperative bleeding after gastric endoscopic submucosal dissection in patients under antithrombotics

... postoperative bleeding after endoscopic submucosal ... - Koh - 被引用次数: 29

Risk factors for bleeding after endoscopic submucosal ... - Tsuji - 被引用次数: 74

Risk factors for bleeding after endoscopic submucosal ... - Higashiyama - 被引用次数: 50

小提示: 仅限搜索简体中文结果。您可以在设置中指定搜索语言

Risk factors for early and delayed post-operative bleeding ...

www.ncbi.nlm.nih.gov/pubmed/25280756 翻译此页

作者: T Matsumura - 2014 - 被引用次数: 1 - 相关文章

2014年10月3日 - Risk factors for early and delayed post-operative bleeding after endoscopic submucosal dissection of gastric neoplasms, including patients with continued use of antithrombotic agents. Matsumura T, Arai M(1), Maruoka D, ...

Antithrombotic drugs are risk factors for delayed ...



[全部](#) [图片](#) [新闻](#) [视频](#) [更多 ▾](#) [搜索工具](#)

找到约 19,800 条结果 (用时 0.95 秒)

Google 学术: Risk factors for postoperative bleeding after gastric endoscopic submucosal dissection in patients under antithrombotics

... postoperative bleeding after endoscopic submucosal ... - Koh - 被引用次数: 30

Risk factors for bleeding after endoscopic submucosal ... - Tsuji - 被引用次数: 75

Risk factors for bleeding after endoscopic submucosal ... - Higashiyama - 被引用次数: 50

小提示: 仅限搜索简体中文结果。您可以在设置中指定搜索语言

Risk factors for early and delayed post-operative bleeding ...

www.ncbi.nlm.nih.gov/pubmed/25280756 ▾ [翻译此页](#)

作者: T Matsumura - 2014 - 被引用次数: 1 - [相关文章](#)

2014年10月3日 - Risk factors for early and delayed post-operative bleeding after endoscopic submucosal dissection of gastric neoplasms, including patients with continued use of antithrombotic agents. Matsumura T, Arai M(1), Maruoka D, ...

Antithrombotic drugs are risk factors for delayed ...

www.ncbi.nlm.nih.gov/pubmed/23622974 ▾ [翻译此页](#)

作者: R Koh - 2013 - 被引用次数: 29 - [相关文章](#)

2013年4月25日 - Antithrombotic drugs are risk factors for delayed postoperative bleeding after endoscopic submucosal dissection for gastric neoplasms. ... to December 2010, we treated 1192 gastric neoplasms in 1032 consecutive patients.